Copyright © 2011 · Magnolia Press

Article



A new genus for *Libinia rostrata* Bell, 1835, with comments on the validity of *Libinia bellicosa* Oliveira, 1944 (Crustacea, Brachyura, Majoidea, Epialtidae)

MARCOS TAVARES^{1,3} & WILLIAM SANTANA²

¹Museum of Zoology, University of São Paulo, Av. Nazareth 481, Ipiranga 04263-000, São Paulo, SP, Brazil. E-mail: mdst@usp.br ²Postdoctoral Research Fellow, Museum of Zoology, University of São Paulo, Av. Nazareth 481, Ipiranga 04263-000, São Paulo, SP, Brazil. E-mail: william_santana@yahoo.com.br

³Corresponding author

Abstract

Stratiolibinia gen. nov., is established for *Libinia rostrata* Bell, 1835, its type species, and *Libinia bellicosa* Oliveira, 1944. The thoracic sternum, male first gonopods, abdomen, and pereiopods are distinctive, and easily distinguish the genus from *Libinia* Leach, 1815 sensu stricto.

Key words: New genus, Decapoda, Pisinae, Libinia, Western Atlantic, Eastern Pacific

Introduction

The rediscovery of the holotype female of *Libinia bellicosa* Oliveira, 1944, and the collection of several additional females and males from different localities in the Western Atlantic, prompted revisiting the validity of this species, sometimes considered to be identical with *Libinia rostrata* Bell, 1835, from the Eastern Pacific. The morphological differences between these two species are discussed and illustrated.

In addition, a comparison between *Libinia rostrata* and *L. bellicosa* with eight out of the 12 species currently placed in *Libinia* Leach, 1815 (Ng *et al.*, 2008), revealed that *L. rostrata* Bell, 1835, and *L. bellicosa* Oliveira, 1944, are no longer tenable in *Libinia*. *Stratiolibinia* **nov. gen.**, is proposed herein to receive both *L. rostrata* and *L. bellicosa*.

Abbreviations used are as follows: cl, carapace length (from the rostral notch to the posterior margin of the carapace); cw, carapace width (branchiostegal spines not included); G1, first male gonopod; mxp3, maxilliped 3; P2-5, pereiopod 2 to pereiopod 5 (P1 is cheliped); ThP17 and ThP18, thoracic pleurites 7 and 8, respectively; WA, Western Atlantic; EP, Eastern Pacific. The material studied belongs to the collections of the Museu Nacional, Universidade Federal do Rio de Janeiro (MNRJ); Museu de Zoologia, Universidade de São Paulo (MZUSP), and the National Museum of Natural History, Smithsonian Institution, Washington D.C. (USNM).

Systematics

Family Epialtidae MacLeay, 1838

Subfamily Pisinae Dana, 1851

Stratiolibinia gen. nov.

Type species. Libinia rostrata Bell, 1835, by present designation. Gender feminine.

Included species. *Stratiolibinia rostrata* (Bell, 1835) gen. nov., comb. nov., and *S. bellicosa* (Oliveira, 1944) gen. nov., comb. nov.