

Three new Nannastacidae (Crustacea: Cumacea) species from a Caribbean mesophotic ecosystem

IORGU PETRESCU¹, TAPAS CHATTERJEE^{2,4} & NIKOLAOS V. SCHIZAS³

¹'Grigore Antipa' National Museum of Natural History, Kiseleff 1, Bucharest 011341, Romania. Email: iorgup@antipa.ro

²Department of Biology, Indian School of Learning, ISM Annexe, P.O. – ISM, Dhanbad 826004, Jharkhand, India.

E-mail: drtchatterjee@yahoo.co.in

³Department of Marine Sciences, University of Puerto Rico at Mayagüez Call Box 9000, Mayagüez, PR 00681, USA.

E-mail: nschizas@gmail.com

⁴Corresponding author

Abstract

Examination of substrata from the mesophotic reefs of Mona Island, Puerto Rico yielded 3 new species of cumaceans, all from the family Nannastacidae: *Cumella achimae* sp. nov., *C. victoriae* sp. nov. and *Nannastacus craciuni* sp. nov. The 3 new species bring the total of new cumacean taxa described from the mesophotic reefs of US Caribbean to 9, highlighting the potential of mesophotic reefs as a biodiversity hotspot. For the first time we report the genus *Nannastacus* from the Caribbean Sea.

Key words: Cumacea, new species, mesophotic cruise, tri-mix rebreathers.

Introduction

Near the declining shallow water coral reefs of the Caribbean there are thriving deeper water coral communities, named mesophotic coral ecosystems (MCEs). These reefs, similarly to their shallow water counterparts, are characterized by macroalgae, sponges, and corals (Lesser *et al.* 2009, Locker *et al.*, 2010; Sherman *et al.*, 2010). Mesophotic coral ecosystems found on the insular and continental slopes of Caribbean islands between 30 and 100 m (Locker *et al.*, 2010). Even at the lower depth range (70–100 m), there is enough light for photosynthesis to take place enhancing the growth of several scleractinian corals (e.g. *Agaricia*, *Montastraea*). These corals are foundation species of the MCEs providing habitat for commercially important species (e.g. groupers, snappers) (Bejarano, 2013) but also for a highly diverse macrofauna and meiofauna. Our knowledge of commercially important fish and corals observed in MCEs is improving but no systematic effort to study the macrofaunal and meiofaunal biodiversity exists.

Previous work on Cumacea from Puerto Rico resulted in a total of 6 new species and 1 new genus from this area, all of them belonging to the family Nannastacidae (Petrescu, Chatterjee & Schizas, 2012, 2013, 2014 in press).

In this paper, we are describing 2 new species of the genus *Cumella* and 1 new species of the genus *Nannastacus* collected during the first of 3 mesophotic cruises (2010–2012) organized by the University of Puerto Rico at Mayagüez (UPRM), the Caribbean Coral Reef Institute (CCRI) and the Department of Marine Sciences (DMS) of UPRM.

Material and methods

All 3 specimens were collected during the first mesophotic cruise of UPRM-CCRI-DMS from southeast Mona Island (18°02'43.9080" N, 67°52'42.7200" W), Puerto Rico, Caribbean Sea, 73.1 m depth, wash from the coral

References

- Băcescu, M. (1971) New Cumacea from the littoral waters of Florida (Caribbean Sea). *Travaux Museum d'Histoire naturelle "Grigore Antipa"*, 11, 5–24.
- Băcescu, M. & Muradian, Z. (1972) Nouvelles espèces de Nannastacidae (Crustacés Cumacés) dans les eaux sahariennes de l'Atlantique. *Revue Travaux Institut Pêches maritime*, 36 (3), 255–269.
- Băcescu, M. (1992) Deux espèces nouvelles de *Cumella* (Crustacea, Cumacea) des grottes sous - marines de Bermuda. *Travaux Museum d'Histoire naturelle "Grigore Antipa"*, 32, 257–262.
- Băcescu, M. & Petrescu, I. (1999) Ordre de Cumacés (Cumacea Krøyer, 1846). *Mémoires de l'Institut océanographique, Monaco*, 19, 391–428.
- Bejarano, I. (2013) *Deep Reef Fishes off La Parguera Insular Slope, Puerto Rico, and their Connectivity with Shallow Reefs, PhD Dissertation, Department of Marine Sciences*. University of Puerto Rico at Mayaguez, Mayaguez, PR, 115 pp.
- Calman, W.T. (1911) On new and rare Crustacea of the Order Cumacea from the collection of the Copenhagen Museum - Part II. The Families Nannastacidae and Diastylidae. *Transaction Zoological Society London*, 18, 341–400.
- Corbera, J. & Sorbe, J.C. (1999) The problematic cumacean *Schizotrema atlanticum* from the Eastern Atlantic: redescription and ecological notes. *Journal of Crustacean Biology*, 19 (1), 123–130.
<http://dx.doi.org/10.2307/1549554>
- Hansen, H.J. (1895) III. Ordnung: Cumacea. In: Isopoden, Cumaceen und Stomatopoden der Plankton Expedition N Atlantic. *Ergebnisse Atlantlantic Ocean Plankton Expedition*, 2, 52–63.
- Lesser, M.P., Slattery, M. & Leichter, J.J. (2009) Ecology of mesophotic coral reefs. *Journal of Experimental Marine Biology and Ecology*, 375, 1–8.
- Locke, S., Armstrong R., Battista, T., Rooney, J., Sherman, C. & Zawada, D. (2010) Geomorphology of mesophotic coral ecosystems: current perspectives on morphology, distribution, and mapping strategies. *Coral Reefs*, 29, 329–345.
<http://dx.doi.org/10.1007/s00338-010-0613-6>
- Petrescu, I. (2002) Cumacea (Crustacea: Peracarida) from Belize. *Travaux du Museum national d'Histoire naturelle "Grigore Antipa"*, 44, 141–203.
- Petrescu, I. (2004) New mentions of Cumaceans (Crustacean: Cumacea) from Cuba. *Travaux National du Museum d'Histoire naturelle "Grigore Antipa"*, 47, 89–95.
- Petrescu, I. & Sterrer, W. (2001) Cumacea (Crustacea) from shallow waters of Bermuda, *Annalen Naturhistorisches Museum Wien*, 103 B, 89–128.
- Petrescu, I. & Heard, R.W. (2004) Cumaceans (Crustacea, Peracarida) from Costa Rica. *Zootaxa*, 721, 1–12.
- Petrescu, I., Chatterjee, T. & Schizas, N.V. (2012) New genus and new species of Cumacea (Crustacea: Peracarida) from the mesophotic coral ecosystem of SW Puerto Rico, Caribbean. *Zootaxa*, 3476, 55–61.
- Petrescu, I., Chatterjee, T. & Schizas, N.V. (2013) Two new species of the genus *Cumella* (Crustacea: Cumacea: Nannastacidae) associated with mesophotic reefs of Puerto Rico and St. Croix, Caribbean Sea. *Cahiers de Biologie Marine*, 54, 257–262.
- Petrescu, I., Chatterjee, T. & Schizas, N.V. (2014) New species of *Cumella* (Crustacea: Cumacea: Nannastacidae) from mesophotic habitats of Mona Island, Puerto Rico, Caribbean Sea. *Cahiers de Biologie Marine*, 55 (2), in Press.
- Petrescu, I., Iliffe, T. & Sarbu, S. (1994) Contributions to the knowledge of the cumaceans (Crustacea) from Jamaica II. Five new species of the genus *Cumella*. *Travaux du Museum d'Histoire naturelle "Grigore Antipa"*, 34, 347–367.
- Sherman, C., Nemeth, M., Ruiz, H., Bejarano, I., Appeldoorn, R., Pagán, F., Schäfer, M. & Weil, E. (2010) Geomorphology and benthic cover of mesophotic coral ecosystems of the upper insular slope of southwest Puerto Rico. *Coral Reefs*, 29, 347–360.
<http://dx.doi.org/10.1007/s00338-010-0607-4>