

<http://dx.doi.org/10.11646/zootaxa.3936.4.4>
<http://zoobank.org/urn:lsid:zoobank.org:pub:26655EB2-0352-4D4F-BBD6-44F89D05DC41>

Description of a new species of the *Miniopterus aelleni* group (Chiroptera: Miniopteridae) from upland areas of central and northern Madagascar

STEVEN M. GOODMAN^{1,2,9}, BEZA RAMASINDRAZANA^{2,3,4,5},
KATE M. NAUGHTON⁶ & BELINDA APPLETON^{7,8}

¹Field Museum of Natural History, 1400 South Lake Shore Drive, Chicago, Illinois 60605, USA. E-mail: sgoodman@fieldmuseum.org

²Association Vahatra, BP 3972, Antananarivo (101), Madagascar. E-mails: sgoodman@yahatra.mg, ramasindrazana@gmail.com

³Département de Biologie Animale, Faculté des Sciences, Université d'Antananarivo, BP 906, Antananarivo (101), Madagascar

⁴Centre de Recherche et de Veille sur les maladies émergentes dans l'Océan Indien, 2 rue Maxime Rivière, 97490 Sainte Clotilde, La Réunion, France

⁵Institut de Recherche pour le Développement, 97492 Sainte Clotilde, La Réunion, France

⁶Marine Invertebrates, Sciences Department, Museum Victoria, GPO Box 666, Melbourne 3001, Victoria, Australia.
E-mail: kmnaughton@gmail.com

⁷Department of Genetics, University of Melbourne, Victoria 3010, Australia. E-mail: b.appleton@unimelb.edu.au

⁸Life and Environmental Science, Deakin University, Victoria, 3216, Australia. E-mail: belinda.appleton@deakin.edu.au

⁹Corresponding author

Abstract

Recent molecular genetic work, combined with morphological comparisons, on Malagasy members of the bat genus *Miniopterus* (Family Miniopteridae), has uncovered a number of cryptic species. Based on recently collected specimens and associated tissues, we examine patterns of variation in *M. aelleni*, the holotype of which comes from Ankarana in northern Madagascar. Using molecular genetic (mitochondrial cytochrome *b*) and morphological characters we describe a new species, *M. ambohitrensis* sp. nov. In northern Madagascar, *M. ambohitrensis* and *M. aelleni* are allopatric, but occur in relatively close geographical contact (approximately 40 km direct line distance) with *M. ambohitrensis* found at Montagne d'Ambre in montane humid forest and *M. aelleni* sensu stricto at Ankarana in dry deciduous forest. Morphologically, this new taxon is differentiated from *M. aelleni* based on pelage coloration, external measurements, craniodental differences, and tragus shape. Comparisons using 725 bp of cytochrome *b* found a divergence of 1.1% within *M. aelleni* sensu stricto, 0.8% within *M. ambohitrensis*, and 3.3% between these two clades. The two sister species do not demonstrate acoustical differences based on recordings made in a flight cage. *Miniopterus ambohitrensis* is known from four localities in the northern and central portions of Madagascar, all from montane regions and across an elevational range from about 800 to 1600 m; its calculated “Extent of occurrence” is 15,143 km². It is possible that this species is at least partially migratory.

Key words: taxonomy, morphology, molecular genetics, new species, montane, humid formation forest, cryptic species

Résumé

Les études morphologiques et moléculaires récemment entreprises sur les espèces du genre *Miniopterus* (Famille des Miniopteridae) ont révélé la présence de diverses espèces cryptiques. Dans la présente étude, nous avons analysé des spécimens récemment collectés et les tissus associés afin de déterminer la variation de *M. aelleni* dont l'holotype vient du Parc National d'Ankarana, situé dans la partie Nord de Madagascar. En se basant sur les données moléculaires (cytochrome *b*) et morphologiques, nous avons décrit une nouvelle espèce, *M. ambohitrensis* sp. nov. Dans la partie nord malgache, *M. aelleni* et *M. ambohitrensis* vivent en allopatrie. Ces deux espèces sont réparties dans des zones géographiquement proches (distantes d'environ 40 km à vol d'oiseau) avec *M. ambohitrensis* dans des zones de forêts humides de montagne du Parc National de la Montagne d'Ambre et *M. aelleni* sensu stricto, dans des zones dominées par des forêts sèches caudicifoliées du Parc National d'Ankarana. Du point de vue morphologique, cette nouvelle espèce de *Miniopterus* se distingue de *M. aelleni* par la couleur du pelage, les mensurations externes et crano-dentaires ainsi que la forme du tragus. L'analyse des 725 pb du cytochrome *b* montre une divergence génétique de 1,1 % chez *M. aelleni* sensu stricto et de 0,8

months (June through August) ranged from 9.7 to 11.2°C (Langrand 1995). We have found bats at the site in torpor, but no evidence of deep hibernation.

In a more practical sense, the species described herein, *M. ambohitrensis*, can be separated from its sister species, *M. aelleni*, by a number of morphological characters, including size, coloration, tragus shape, and craniodental features. These two species have allopatric distributions. As has been shown for several genetically distinct cryptic species of small to moderate size Malagasy *Miniopterus*, tragus shape can be diagnostic to distinguish taxa and can be used to diagnose different clades (Goodman *et al.* 2007, 2008, 2009a, 2009b, 2010, 2011). *Miniopterus ambohitrensis* possesses a number of mensural and phenotypic characters, including tragus shape, which allows it to be distinguished from *M. aelleni*. A detailed study overlaying clade allocation with a variety of different genetic markers and morphological variation, specifically tragus shape, as well as bioacoustic characteristics is needed to resolve the affinities of animals currently placed in the P-group.

Acknowledgements

Conservation International (CABS), The John T. and Catherine D. MacArthur Foundation, and the Volkswagen Foundation have generously supported field research associated with this paper. We are grateful to the Direction du Système des Aires Protégées, Direction Générale de l'Environnement et des Forêts, and Madagascar National Parks (ex. Association National pour la Gestion des Aires Protégées or ANGAP) for issuing permits to conduct faunal surveys and aid in numerous ways. Yann Gomard, Corrie M. Schoeman, and Peter Taylor helped with fieldwork. For access to specimens under their care, we are indebted to Kristofer Helgen, USNM; Jean-Marc Pons, MNHN; and Daniel Rakotondravony, UADBA. Herivololona Mbola Rakotondratsimba kindly prepared Figure 1 and John Weinstein Figures 4 and 5. BR received a postdoctoral fellowship from the RunEmerge project funded by the European Frame work program FP7 Capacities/Regpot, and post-doctoral grants from Fonds de Coopération Régionale of the Préfecture de La Réunion and from the Dr. Ralph and Marian Falk Medical Research Trust to The Field Museum of Natural History, Chicago. We would like to thank an anonymous reviewer for their detailed comments on an earlier version of the ms.

References

- Appleton, B.R., McKenzie, J.A. & Christidis, L. (2004) Molecular systematics and biogeography of the bent-wing bat complex *Miniopterus schreibersii* (Kuhl, 1817) (Chiroptera: Vespertilionidae). *Molecular Phylogenetics and Evolution*, 31, 431–439.
<http://dx.doi.org/10.1016/j.ympev.2003.08.017>
- Battistini, R. (1965) Problèmes géomorphologiques de l'Extrême Nord de Madagascar. *Madagascar; Revue de Géographie*, 7, 1–61.
- Bernard, R.T.F. (1980) Reproductive cycles of *Miniopterus schreibersi natalensis* (Kuhl, 1819) and *M. fraterculus* Thomas and Schwann, 1906. *Annals of the Transvaal Museum*, 32, 55–64.
- Bilgin, R., Gürün, K., Maraci, O., Furman, A., Hulva, P., Çoraman, E., Lučan, R.K., Bartonička, T. & Horáček, I. (2012) Syntopic occurrence in Turkey supports separate species status for *Miniopterus schreibersii schreibersii* and *M. schreibersii pallidus* (Mammalia: Chiroptera). *Acta Chiropterologica*, 14, 279–289.
<http://dx.doi.org/10.3161/150811012X661611>
- Brenon, P. (1972) The geology of Madagascar. In: Battistini, R. & Richard-Vindard, G. (Eds.), *Biogeography and ecology in Madagascar*. Dr. W. Junk, The Hague, pp. 27–86.
- Cardinal, B.R. & Christidis, L. (2000) Mitochondrial DNA and morphology reveal three geographically distinct lineages of the large bentwing bat (*Miniopterus schreibersii*) in Australia. *Australian Journal of Zoology*, 48, 1–19.
<http://dx.doi.org/10.1071/ZO99067>
- Christidis, L., Goodman, S.M., Naughton, K. & Appleton, B. (2014) Insights into the evolution of a cryptic radiation of bats: dispersal and ecological radiation of Malagasy *Miniopterus* (Chiroptera: Miniopteridae). *PLoS ONE*, 9 (3), e92440.
<http://dx.doi.org/10.1371/journal.pone.0092440>
- Donque, G. (1975) *Contribution à l'étude du climat de Madagascar*. Nouvelle Imprimerie des Arts Graphiques, Antananarivo, 478 pp.
- Furman, A., Oztunc, T. & Coraman, E. (2010) On the phylogeny of *Miniopterus schreibersii schreibersii* and *Miniopterus schreibersii pallidus* from Asia Minor in reference to other *Miniopterus* taxa (Chiroptera: Vespertilionidae). *Acta Chiropterologica*, 12, 61–72.

- <http://dx.doi.org/10.3161/150811010X504581>
- Gummel, N.J. & Akiyama, S. (1996) An efficient method for the extraction of DNA from vertebrate tissues. *Trends in Genetics*, 12, 338–339.
[http://dx.doi.org/10.1016/S0168-9525\(96\)80005-9](http://dx.doi.org/10.1016/S0168-9525(96)80005-9)
- Goodman, S.M. (2011) *Les chauves-souris de Madagascar*. Association Vahatra, Antananarivo, 129 pp.
- Goodman, S.M. & Ramasindrazana, B. (2013) Bats or the Order Chiroptera. In: Goodman, S.M. & Raherilalao, M.J. (Eds.), *Atlas of selected land vertebrates of Madagascar*. Association Vahatra, Antananarivo, pp. 169–209.
- Goodman, S.M., Ryan, K.E., Maminirina, C.P., Fahr, J., Christidis, L. & Appleton, B. (2007) The specific status of populations on Madagascar referred to *Miniopterus fraterculus* (Chiroptera: Vespertilionidae), with description of a new species. *Journal of Mammalogy*, 88, 1216–1229.
<http://dx.doi.org/10.1644/06-MAMM-A-285R1.1>
- Goodman, S.M., Bradman, H.M., Maminirina, C.P., Ryan, K.E., Christidis, L. & Appleton, B. (2008) A new species of *Miniopterus* (Chiroptera: Miniopteridae) from lowland southeastern Madagascar. *Mammal Biology*, 73, 199–213.
<http://dx.doi.org/10.1016/j.mambio.2007.12.003>
- Goodman, S.M., Maminirina, C.P., Weyeneth, N., Bradman, H.M., Christidis, L., Ruedi, M. & Appleton, B. (2009a) The use of molecular and morphological characters to resolve the taxonomic identity of cryptic species: the case of *Miniopterus manavi* (Chiroptera, Miniopteridae). *Zoologica Scripta*, 38, 339–363.
<http://dx.doi.org/10.1111/j.1463-6409.2008.00377.x>
- Goodman, S.M., Maminirina, C.P., Bradman, H.M., Christidis, L. & Appleton, B. (2009b) The use of molecular phylogenetic and morphological tools to identify cryptic and paraphyletic species: examples from the diminutive long-fingered bats (*Miniopterus*: Miniopteridae: Chiroptera) on Madagascar. *American Museum Novitates*, 3669, 1–33.
- Goodman, S.M., Maminirina, C.P., Bradman, H.M., Christidis, L. & Appleton, B. (2010) Patterns of morphological and genetic variation in the endemic Malagasy bat *Miniopterus gleni* (Chiroptera: Miniopteridae), with the description of a new species, *M. griffithsi*. *Journal of Zoological Systematics and Evolutionary Research*, 48, 75–86.
<http://dx.doi.org/10.1111/j.1439-0469.2009.00524.x>
- Goodman, S.M., Ramasindrazana, B., Maminirina, C.P., Schoeman, M.C. & Appleton, B. (2011) Morphological, bioacoustical, and genetic variation in *Miniopterus* bats from eastern Madagascar, with the description of a new species. *Zootaxa*, 2880, 1–19.
- Guindon, S. & Gascuel, O. (2003) A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Systematic Biology*, 52, 696–704.
<http://dx.doi.org/10.1080/10635150390235520>
- Harper, G.J., Steininger, M.K., Tucker, C.J., Juhn, D. & Hawkins, F. (2007) Fifty years of deforestation and forest fragmentation in Madagascar. *Environmental Conservation*, 34, 325–333.
<http://dx.doi.org/10.1017/S0376892907004262>
- Huelsenbeck, J.P. & Ronquist, F. (2001) MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics*, 17, 754–755.
<http://dx.doi.org/10.1093/bioinformatics/17.8.754>
- Hurlbert, S.H. (1984) Pseudoreplication and the design of ecological field experiments. *Ecological Monographs*, 54, 187–211.
<http://dx.doi.org/10.2307/1942661>
- Irwin, D.M., Kocher, T.D. & Wilson, A.C. (1991) Evolution of the cytochrome b gene of mammals. *Journal of Molecular Evolution*, 32, 128–144.
<http://dx.doi.org/10.1007/BF02515385>
- IUCN (2012) *IUCN Red List categories and criteria: version 3.1. Second Edition*. International Union for the Conservation of Nature, Gland, Switzerland and Cambridge, 32 pp.
- Kimura, M. (1980) A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *Journal of Molecular Evolution*, 16, 111–120.
<http://dx.doi.org/10.1007/BF01731581>
- Koopman, K.F. (1994) Chiroptera: Systematics. In: Niethammer, J., Schliemann, H. & Starck, D. (Eds) *Handbook of Zoology*. Vol. 8. Walter de Gruyter, Berlin, pp. 1–217.
- Langrand, O. (1995) The effects of forest fragmentation on bird species in Madagascar: A case study from Ambohitantely Forest Reserve on the Central High Plateau. Master of Science, Department of Zoology and Entomology, University of Natal, Pietermaritzburg. [total page number unknown]
- Miller-Butterworth, C.M., Eick, G., Jacobs, D.S., Schoeman, M.C. & Harley, E.H. (2005) Genetic and phenotypic differences between South African long-fingered bats, with a global miniopterine phylogeny. *Journal of Mammalogy*, 86, 1121–1135.
<http://dx.doi.org/10.1644/05-MAMM-A-021R1.1>
- Monadjem, A., Goodman, S.M., Stanley, W.T. & Appleton, B. (2013) A cryptic new species of *Miniopterus* from south-eastern Africa based on molecular and morphological characters. *Zootaxa*, 3746 (1), 123–142.
<http://dx.doi.org/10.11646/zootaxa.3746.1.5>
- Obrist, M.K. (1995) The influence of individual, habitat and conspecifics on sonar signal design. *Behavioral Ecology and Sociobiology*, 36, 207–219.
<http://dx.doi.org/10.1007/BF00177798>
- Onyango, D.W., Gachoka, J.M., Otiang'a-Owiti, G.E. & Hendrickx, A.G. (1995) Seasonally dependent testicular apoptosis in

- the tropical long-fingered bat (*Miniopterus inflatus*). *Zeitschrift für Säugetierkunde*, 60, 206–214.
- Peterson, R.L., Eger, J.L. & Mitchell, L. (1995) Chiroptères. Vol. 84. Faune de Madagascar. Muséum national d'Histoire naturelle, Paris, 204 pp.
- Posada, D. (2008) jModelTest: Phylogenetic Model Averaging. *Molecular Biology and Evolution*, 25, 1253–1256.
<http://dx.doi.org/10.1093/molbev/msn083>
- Puechmaille, S.J., Allegrini, B., Benda, P., Gürün, K., Šrámek, J., Ibañez, C., Juste, J. & Bilgin, R. (2014) A new species of the *Miniopterus schreibersii* species complex (Chiroptera: Miniopteridae) from the Maghreb Region, North Africa. *Zootaxa*, 3794 (1), 108–124.
<http://dx.doi.org/10.11646/zootaxa.3794.1.4>
- Rabearivony, J., Thorstrom, R., Rene de Roland, L.-A., Rakotondratsima, M., Andriamalala, T.R.A., Sam, T.S., Razafimanjato, G., Rakotondravony, D., Raselimanana, A.P. & Rakotoson, M. (2010) Protected area surface extension in Madagascar: Do endemism and threatened species remain useful criteria for site selection? *Madagascar Conservation and Development*, 5, 35–47.
- Ramasindrazana, B. & Goodman, S.M. (2014) Documented record of *Triaenops menamena* (Family Hipposideridae) in the Central Highlands of Madagascar. *African Bat Conservation News*, 35, 6–7.
- Ramasindrazana, B., Goodman, S.M., Schoeman, M.C. & Appleton, B. (2011) Identification of cryptic species of *Miniopterus* bats (Chiroptera: Miniopteridae) from Madagascar and the Comoros using bioacoustics overlaid on molecular genetic and morphological characters. *Biological Journal of the Linnean Society*, 104, 284–302.
<http://dx.doi.org/10.1111/j.1095-8312.2011.01740.x>
- R Core Team. (2013) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna. Available from: <http://www.R-project.org/> (accessed 18 February 2015)
- Ronquist, F., Teslenko, M., van der Mark, P., Ayres, D.L., Darling, A., Höhna, S., Larget, B., Liang, L., Suchard, M.A. & Huelsenbeck, J.P. (2012) MrBayes 3.2: Efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology*, 61, 539–542.
<http://dx.doi.org/10.1093/sysbio/sys029>
- Rossi, G. (1974) Review of J.P. Karche. 1972. Contribution à l'étude géologique de la Montagne d'Ambre et des régions voisines du Nord de Madagascar. Thèse d'Etat (Sciences), Université de Besançon. Madagascar. *Revue de Géographie*, 24, 95–96.
- Schatz, G. (2000) Endemism in the Malagasy flora. In: Lourenço, W.R. & Goodman, S.M. (Eds.), *Diversité et endémisme à Madagascar*. Mémoires de la Société de Biogéographie, Paris, pp. 1–8.
- Schnitzler, H.-U. & Kalko, E.K.V. (2001) Echolocation by insect-eating bats. *Bioscience*, 51, 557–569.
[http://dx.doi.org/10.1641/0006-3568\(2001\)051\[0557:EBIEB\]2.0.CO;2](http://dx.doi.org/10.1641/0006-3568(2001)051[0557:EBIEB]2.0.CO;2)
- Schoeman, M.C. & Jacobs, D.S. (2008) The relative influence of competition and prey defenses on the phenotypic structure of insectivorous bat ensembles in southern Africa. *PLoS ONE*, 3 (11), e3715.
<http://dx.doi.org/10.1371/journal.pone.0003715>
- Schoeman, M.C., Goodman, S.M., Ramasindrazana, B. & Kabinouva, D. (2015) Species interactions during diversification and community assembly in Malagasy *Miniopterus* bats. *Evolutionary Ecology*, 29, 17–47.
<http://dx.doi.org/10.1007/s10682-014-9745-4>
- Siemers, B.M. (2004) Bats in the field and in a flight cage: recording and analysis of their echolocation calls and behavior. In: Brigham, R.M., Kalko, E.K.V., Jones, G., Parsons, S. & Limpens, H.J.G.A. (Eds.), *Bat echolocation research: tools, techniques and analysis*. Bat Conservation International, Austin, Texas, pp. 107–113.
- Smith, M.F. & Patton, J.L. (1991) Variation in mitochondrial cytochrome *b* sequence in natural populations of South American akodontine rodents (Muridae: Sigmodontinae). *Molecular Biology and Evolution*, 8, 85–103.
- Šrámek, J., Gvoždík, V. & Benda, P. (2013) Hidden diversity in bent-winged bats (Chiroptera: Miniopteridae) of the western Palaearctic and adjacent regions: implications for taxonomy. *Zoological Journal of the Linnean Society*, 167, 165–190.
<http://dx.doi.org/10.1111/j.1096-3642.2012.00870.x>
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M. & Kumar, S. (2011) MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology and Evolution*, 28, 2371–2739.
<http://dx.doi.org/10.1093/molbev/msr121>
- Toews, D.P.L. & Brelsford, A. (2012) The biogeography of mitochondrial and nuclear discordance in animals. *Molecular Ecology*, 21 (16), 3907–3930.
<http://dx.doi.org/10.1111/j.1365-294X.2012.05664.x>
- van der Merwe, M. (1975) Preliminary study on the annual movements of the Natal clinging bat. *South African Journal of Science*, 71, 237–241.
- van der Merwe, M. (1979) Growth of ovarian follicles in the Natal clinging bat. *South African Journal of Science*, 14, 111–117.
- Voigt, C.C., Helbig-Bonitz, M., Kramer-Schadt, S. & Kalko, E.K.V. (2014) The third dimension of bat migration: Evidence for elevational movements of *Miniopterus natalensis* along the slopes of Mount Kilimanjaro. *Oecologia*, 174, 751–764.
<http://dx.doi.org/10.1007/s00442-013-2819-0>
- Zwickl, D.J. (2006) *Genetic algorithm approaches for the phylogenetic analysis of large biological sequence datasets under the maximum likelihood criterion*. PhD thesis, University of Texas, Austin.