



RESEARCH ARTICLE

A revised checklist species of the Curculionoidea (Coleoptera, excluding Scolytinae) of Rostov Oblast and Kalmykia, the southern part of European Russia

Yuri G. Arzanov¹

¹*Institute of Arid Zones of RAS, Chekhov str., 41, Rostov-on-Don, 344006, Russia. E-mail: arz99@mail.ru*

Abstract: This work provides a list of 703 species of weevils (Coleoptera: Curculionoidea, excluding Scolytinae) recorded in two adjacent administrative regions in the southern part of European Russia: Rostov Oblast and the Republic of Kalmykia. The species diversity of the ten recorded families is as follows: Nemonychidae (1), Anthribidae (10), Rhynchitidae (12), Attelabidae (2), Apionidae (70), Nanophyidae (12), Brachyceridae (2), Dryophthoridae (4), Erihinidae (13), Curculionidae (577, excluding Scolytinae). In total, 151 species are added to the faunal list of the southern European Russia.

Key words: Coleoptera, Curculionoidea, southern European Russia, Rostov Oblast, Kalmykia.

Introduction

The recently published two weevil volumes of the Catalogue of Palaearctic Coleoptera (Löbl & Smetana 2011, 2013) significantly updated numerous regional faunas. However, unlike the other treated countries, both Russia and China were regionally subdivided into smaller units. The subregions referring to the southern part of European Russia (coded as ST in Löbl & Smetana 2011, 2013) is rather heterogeneous in its physical and climatic characteristics, with the landscapes varying from deserts and steppes to broadleaf forests and alpine meadows.

This paper focuses on the faunal list of weevils (Curculionoidea, excluding Scolytinae) of two adjacent administrative regions of southern Russia: Rostov Oblast and the Republic of Kalmykia (=Kalmykia). They both belong to the same unit of “Russia: South European territory” as adopted in the aforementioned catalogue (Löbl & Smetana 2011,

2013) (Fig. 1). The region is predominantly flat, lacking significant mountains, and is heavily modified for agriculture. The original vegetation was predominantly of steppe formation (=grasslands) on the west (Rostov Oblast) and semi deserts or deserts in the east (Kalmykia), with broadleaf forests almost exclusively restricted to river valleys. Relatively few previous attempts to document weevil diversity of the region focussed mainly on the pest species, and all are now outdated (Kiseritzky 1912; Lukjanovitsh 1926; Dobrovolsky 1951; Arzanov 1988, 1990).




Figure 1. Map of Rostov Oblast and Kalmykia showing borders of ten internal regions. Subregion abbreviations are: Upper Don (UD), Donets Ridge (DR), Donets Elevation (DE), Lower Don (LD), Priazovia (P), Manych River Valley (MV), Salsk Steppe (SS) Ergeni Elevation (E), Sarpinsk Lowland (SL) and Black Lands (BL). Total number of species of Curculionoidea (excepting Scolytinae) is given inside each subregion.

Material and methods

The present list is predominantly based on specimens that have been annually collected by the authors in both Rostov Oblast and Kalmykia since 1975. Identifications were made by comparing specimens with the vouchers deposited in the collection of the

Zoological Institute in St. Petersburg, Russia. Classification and taxonomic arrangement follow that of Alonso-Zarazaga & Lyal (1999, 2002), which was partly adopted in the weevil volumes of the Catalogue of Palaearctic Coleoptera (Löbl & Smetana 2011, 2013), plus a few subsequent modifications (Korotyeav & Fridman 2011; Lyal 2014; Osella & Bello 2010). Except for the families and subfamilies, all taxa of the same rank are arranged alphabetically.

For the purposes of the present checklist, the entire territory of Rostov Oblast and Kalmykia is further subdivided in ten non-administrative regions to represent discrete biological and climatic entities (Fig. 1): Upper Don (UD), Donets Ridge (DR), Donets Elevation (DE), Lower Don (LD), Priazovia (P), Manych River Valley (MV), Salsk Steppe (SS) Ergeni Elevation (E), Sarpinsk Lowland (SL) and Black Lands (BL).

Results

The following 151 species are added to the faunal list of the southern European Russia region, as delimited in the recent “Catalogue of Palaearctic Coleoptera” (Löbl & Smetana 2011, 2013): **APIONIDAE**: **Apionini**: *Apion cruentatum* Walton, 1844, *A. haematodes* Kirby, 1808; **Aplemonini**: *Perapion marchicum* (Herbst, 1797), *P. myochroum* (Schilsky, 1902); **Exapiini**: *Exapion corniculatum* (Germar, 1817), *E. difficile* (Herbst, 1797); **Oxystomatini**: *Cyanapion afer* (Gyllenhal, 1833), *Betulapion simile simile* (Kirby, 1811); **Piezotrachelini**: *Protapion ruficrus* (Germar, 1817). **NANOPHYIDAE**: *Dieckmanniellus gracilis* (Redtenbacher, 1847), *D. nitidulus* (Gyllenhal, 1838), *Nanomimus hemisphaericus* (Olivier, 1807), *Nanophyes brevis brevis* Boheman, 1845. **BRACHYCYERIDAE**: *Brachycerus sinuatus* Olivier, 1807. **ERIRHINIDAE**: **Arthostenini**: *Arthostenus fullo* Boheman, 1836; **Erirhinini**: *Bagoopsis volgensis* Faust, 1881, *B. globicollis* (Fairmaire, 1863), *Icaris sparganii* (Gyllenhal, 1835), *Notaris acridulus* (Linnaeus, 1758), *N. scirpi* (Fabricius, 1792), *Picia sinuatocollis* (Faust, 1885), *Procas picipes steveni* (Krynicki, 1832), *Thryogenes festucae* (Herbst, 1795), *Th. nereis* (Paykull, 1800), *Tournotaris bimaculata* (Fabricius, 1787). **Tanysphyrini**: *Tanysphyrus lemnae* (Paykull, 1792). **CURCULIONIDAE**: **Curculioninae** **Anthonomini**: *Anthonomus pinivorax* Silfverberg, 1977, *A. bituberculatus* C. G. Thomson, 1868, *A. pedicularius* (Linnaeus, 1758), *Bradybatus kellneri* Bach, 1854, *B. seriesetosus* Petri, 1912; **Cionini**: *Cionus hortulanus* (Geoffroy, 1785), *C. olivieri* Rosenschoeld, 1838, *C. scrophulariae* (Linnaeus, 1758), *C. tuberculosus* (Scopoli, 1763); **Curculionini** Latreille, 1802, *Curculio elephas* (Gyllenhal, 1835); **Ellescini**: *Dorytomus dejeani* Faust, 1883, *D. filirostris* Gyllenhal, 1835, *D. ictor* (Herbst, 1795), *D. longimanus* (Forster, 1771), *D. melanophthalmus* (Paykull, 1792), *D. nebulosus* Gyllenhal, 1835, *D. tremulae* (Fabricius, 1787), *D. suratus* (Gyllenhal, 1835), *Ellescus infirmus* (Herbst, 1795); **Itini**: *Ita kirschii* Tournier, 1878; **Mecinini**: *Cleopomiarus distinctus* (Boheman, 1845), *Gymnetron melanarium* (Germar, 1821), *G. niloticum* Kirsch, 1881, *G. rotundicolle* Gyllenhal, 1838, *G. veronicae* (Germar, 1821), *G. villosum* Gyllenhal, 1838, *G. vittipenne* Marseul, 1876, *G. terminassianae* Smreczyński, 1975, *Mecinus collaris* Germar, 1821, *M. heydenii* Wencker, 1866, *M. labilis* (Herbst, 1795), *M. plantaginis* (Eppelsheim, 1875), *Miarus ajugae* (Herbst, 1795), *Rhinusa antirrhini* (Paykull, 1800), *Rh. asellus* (Gravenhorst, 1807), *Rh. florum* (Rübsaamen, 1895), *Rh. linariae* (Panzer, 1795), *Rh. melas* Boheman, 1838, *Rh. neta* (Germar, 1821), *Rh. pilosa* (Gyllenhal, 1838); **Rhamphini**: *Orchestes alni* (Linnaeus, 1758), *O. mutabilis* Boheman, 1843, *Pseudorchestes circumvistulanus* (Bialooki, 1997), *P. horioni* (Dieckmann, 1958); **Smicronychini**: *Smicronyx coecus* (Reich, 1797), *S. jungermanniae* (Reich, 1797), *S. nebulosus* Tounier, 1874, *S. reichii* (Gyllenhal, 1835), *S. smreczynskii* (F. Solari, 1952); **Storeini**: *Pachytychius transcaucasicus* Pic, 1913; **Tychiini**: *Lignyodes bischoffi* (Blatchley,

1916); *Sibinia bipunctata* Kirsch, 1871, *S. phalerata* Gyllenhal, 1835, *S. primita* Herbst, 1795, *S. variata* Gyllenhal, 1835, *Tychius argentatus* Chevrolat, 1859, *T. bedeli* Faust, 1885, *T. flavus* Becker, 1864, *T. hauseri* Faust, 1889, *T. pusillus* Germar, 1842, *T. rufirostris* Schoenherr, 1832, *T. squamulatus* Gyllenhal, 1835, *T. sulphureus* Faust, 1881, *T. trivialis* Boheman, 1843, *T. uralensis* Pic, 1902; **Bagoinae:** *Bagous argillaceus* Gyllenhal, 1836, *B. biimpressus* Fåhraeus, 1845, *B. diglyptus* Boheman, 1845, *B. frivaldszkyi* Tournier, 1874, *B. glabrirostris* (Herbst, 1795), *B. limosus* (Gyllenhal, 1827), *B. lutosus* (Gyllenhal, 1813), *B. lutulentus* (Gyllenhal, 1813), *B. petro* (Herbst, 1795), *B. puncticollis* Boheman, 1845, *B. subcarinatus* Gyllenhal, 1836, *B. tubulus* Caldara & O'Brien, 1994; **Baridinae:** *Elasmobaris alboguttata* (H. Brisout de Barnevile, 1870), *E. signifera* (Faust, 1881), *Melanobaris atramentaria* (Boheman, 1836), *M. nigritarsis* (Boheman, 1844); **Ceutorhynchinae:** **Ceutorhynchini:** *Ceutorhynchus canaliculatus* C. N. F. Brisout de Barnevile, 1869, *C. coarctatus* Gyllenhal, 1837, *C. constrictus* (Marsham, 1802), *C. difficilis* Schultze, 1898, *C. sulcatus* C. N. F. Brisout de Barnevile, 1869, *Datonychus angulosus* (Boheman, 1845), *D. arquata* (Herbst, 1795), *Glocianus lethierryi* (C. N. F. Brisout de Barnevile, 1866), *G. moelleri* (C. G. Thomson, 1868), *G. polystriatus* (Schultze, 1898), *Mogulones abbreviatulus* (Fabricius, 1792), *M. albosignatus* (Gyllenhal, 1837), *M. dimidiatus* (J. Frivaldszky, 1865), *M. raphani* (Fabricius, 1792), *M. venedicus* (Weise, 1879), *Phrydiuchus topiarius* (Germar, 1824), *Platygasteronyx solskyi* (Faust, 1885), *Poophagus hopffgarteni* Tournier, 1873, *Prisistus caucasicus caucasicus* (Kirsch, 1879), *P. obsoletus* (Germar, 1824), *Thamiocolus sahlbergi* (C. R. Sahlberg, 1845), *Rhinoncus castor* (Fabricius, 1792); **Hypurini:** *Aphytobius sphaerion* Boheman, 1845; **Cyclominae:** *Gronops inaequalis* Boheman, 1842, *Entomoderus phrygius* (K. Daniel & J. Daniel, 1903); **Entiminae:** **Brachyderini:** *Brachyderes incanus* (Linnaeus, 1758); **Naupactini:** *Mesagroicus poriventris* Reitter, 1903; **Otiorhynchini:** *Otiorhynchus simulans* Stierlin, 1877; **Polydrusini:** *Liophloeus tessulatus* (O. F. Müller, 1776), *Polydrusus viridicinctus* Gyllenhal, 1834; **Sitonini:** *Sitona waterhousei* Walton, 1846; **Hyperinae:** *Hypera arundinis* (Paykull, 1792), *H. rogenhoferi* (Ferrari, 1866); **Lixinae:** **Cleonini:** *Coniocleonus hollbergii* (Fåhraeus, 1842), *Pleurocleonus sollicitus* (Gyllenhal, 1834), *Pseudocleonus grammicus* (Panzer, 1789), *Rhabdorrhynchus echii* (Brahm, 1790), *Xanthochelus nomas* (Pallas, 1771); **Lixini:** *Larinus idoneus* Gyllenhal, 1835, *L. sibiricus* Gyllenhal, 1835; **Mesoptiliinae:** *Magdalis phlegmatica* (Herbst, 1797); **Molytinae:** **Cryptorhynchini:** *Cryptorhynchus lapathi* (Linnaeus, 1758); **Pissodini:** *Pissodes pini pini* (Linnaeus, 1758); **Orobitinae:** *Orobitis cyaneus* (Linnaeus, 1758). The following 15 species previously wrongly cited for either Rostov Oblast or the Republic of Kalmykia should be excluded from the faunal lists of both regions: *Aulacobaris picicornis* (Marsham, 1802), *Aulacobaris prasina* (Boheman, 1836), *Conorhynchus nigrivittis* (Pallas, 1781), *Elasmabaris signifera* (Faust, 1881), *Liocleonus clathratus* (Olivier, 1807), *Lixus albomarginatus* Bohemann, 1842, *Minyops minutus rufus* Ménétriés, 1849, *Otiorhynchus (Nubidanus) juvenilis* Schoenherr, 1832, *Pholicodes (Pseudopholicodes) albidus* (Boheman, 1840), *Pholicodes inauratus arzanovi* Davidian, 1992, *Pseudocleonus marginicollis* (Fåhraeus, 1842), *Rhinocyllus conicus* (Frölich, 1792), *Entomoderus phrygius* (K. Daniel & J. Daniel, 1903), *Maximus aemulus* (Faust, 1894), *Xanthochelus nomas* (Pallas, 1771).

The following 15 species were erroneously included in the faunal lists of either Rostov Oblast or Kalmykia:

Aulacobaris picicornis (Marsham, 1802): Dobrovolsky (1951), Arzanov (1990); species presence is not confirmed;

Aulacobaris prasina (Boheman, 1836): Dobrovolsky (1951), (1990); species presence is not confirmed;

Conorhynchus nigrivittis (Pallas, 1781): Arzanov (1990); misidentification of *Conorhynchus kindermanni* Faust, 1904;
Elasmabar is signifera (Faust, 1881): Arzanov (1990); misidentification of *Elasmabar is annularis* (Faust, 1881);
Liocleonus clathratus (Olivier, 1807): Arzanov (1988); likely mislabelled specimen;
Lixus albomarginatus Bohemann, 1842: Kiseritzky (1912), Dobrovolsky (1951); misidentification of *Lixus ascanii* Linnaeus, 1767 (Dieckmann, 1980);
Minyops minutus rufus Ménétríés, 1849: Arzanov (1990); misidentification of *Minyops carinatus* Linnaeus, 1767 (Osella, Bello, 2009);
Otiorhynchus (Nubidanus) juvenilis Schoenherr, 1832: Arzanov (1990); misidentification of *Otiorhynchus* sp. pr. *unctuosus* Germar, 1824;
Pholicodes (Pseudopholicodes) albidus (Boheman, 1840): Arzanov (1990); misidentification of *Pholicodes* sp. (Davidian, 1992);
Pholicodes inauratus arzanovi Davidian, 1992: Arzanov (1990); misidentification of *Pholicodes trivialis* Boheman, 1834 (Davidian, 1992);
Pseudocleonus marginicollis (Fähraeus, 1842): Lukjanovitsh (1926), Arzanov (1988, 1990); species presence is not confirmed;
Rhinocyllus conicus (Frölich, 1792): Arzanov (1988); misidentification of *Rhinocyllus oblongus* Capiomont, 1873;
Entomoderus phrygius (K. Daniel & J. Daniel, 1903): Arzanov (1990); misidentification of *Rhyti rrhinus gibbus* (Kolenati, 1858);
Maximus aemulus (Faust, 1894): Arzanov (1988); likely mislabelled specimen;
Xanthochelus nomas (Pallas, 1771): Arzanov (1990); misidentification of *Xanthoprochelus eversmanni* (Fahraeus, 1842).

The known fauna of Curculionoidea (excluding Scolytinae) of both Rostov Oblast and Kalmykia (Table 1) consists of ten families and 700 species: Nemonychidae (1), Anthribidae (7), Rhynchtidae (12), Attelabidae (2), Apionidae (70), Nanophyidae (12), Brachyceridae (2), Dryophthoridae (4), Erirhinidae (13) and Curculionidae (577, excluding Scolytinae). Rostov Oblast has 620 recorded species, while 301 species were recorded in Kalmykia.

Table 1. Revised checklist of 700 species of Curculionoidea (excluding Scolytinae) of Rostov Oblast and Kalmykia. Species not cited for southern part of European Russia in the Catalogue of Palaearctic Coleoptera (Löbl & Smetana 2011, 2013) are marked with asterisk (*). Subregionals abbreviations are explained in Fig. 1.

NEMONYCHIDAE Bedel, 1882						
NEMONYCHINAE Bedel, 1882						
<i>Nemonyx</i> L. Redtenbacher, 1845						
<i>lepturoides</i> (Fabricius, 1801)	UD	DR	DE	LD	P	
ANTHRIBIDAE Billberg, 1820						
ANTHRIBINAE Billberg, 1820						
Anthribini Billberg, 1820						
<i>Anthribus</i> Geoffroy, 1762						
<i>nubelosus</i> Forster, 1770	UD	DR	DE	LD	P	
<i>Platyrhinini</i> Imhoff, 1856						

<i>Platyrhinus</i> Clairville, 1798										
<i>resinosus</i> (Scopoli, 1763)	UD	DR	DE	LD	P					
<i>Platystomini</i> Pierce, 1916										
<i>Platystomos</i> D. H. Schneider, 1791										
<i>albinus</i> (Linnaeus, 1758)	UD	DR	DE	LD	P					
<i>Tropiderini</i> Lacordaire, 1865										
<i>Gonotropis</i> LeConte, 1876										
<i>dorsalis</i> (Gyllenhal, 1813)			DE	LD						
<i>Tropideres</i> Schoenherr, 1823										
<i>albirostris</i> (Schaller, 1783)	UD	DR	DE	LD	P					
<i>Zygaenodini</i> Lacordaire, 1865										
<i>Dissoleucas</i> Jordan, 1925										
<i>niveirostris</i> (Fabricius, 1798)	UD	DR	DE	LD	P					
<i>Exechesops</i> Schoenherr, 1847										
<i>leucopis</i> (Jordan, 1928)	UD	DR	DE							
<i>URODONTINAE</i> C.G. Thomson, 1859										
<i>Bruchela</i> Dejean, 1821										
<i>pygmaea</i> (Gyllenhal, 1833)	UD	DR	DE	LD	P	MV	SS	E	SL	BL
<i>rufipes rufipes</i> (Olivier, 1790)			DE	LD	P	MV	SS	E	SL	BL
<i>suturalis</i> (Fabricius, 1792)	UD	DR	DE	LD	P	MV	SS	E		
RHYNCHITIDAE Gistel, 1848										
<i>RHYNCHITINAE</i> Gistel, 1848										
<i>Byctiscini</i> Voss, 1923										
<i>Bystiscus</i> C. G. Thomson, 1859										
<i>betulae</i> (Linnaeus, 1758)	UD	DR	DE	LD	P	MV	SS	E		
<i>populi</i> (Linnaeus, 1758)	UD	DR								
<i>Rhinchitini</i> Gistel, 1848										
<i>Epirhynchites</i> Voss, 1969										
<i>auratus</i> (Scopoli, 1763)	UD	DR	DE	LD	P	MV	SS	E		
<i>giganteus</i> (Schoenherr, 1832)	UD	DR	DE	LD						
<i>Involvulus</i> Schrank, 1798										
<i>cupreus</i> (Linnaeus, 1758)	UD	DR	DE	LD						
<i>Lasiorhynchites</i> Jekel, 1860										
<i>sericeus</i> (Herbst, 1797)	UD	DR								
<i>Neocoenorhinus</i> subgen. <i>Neocoenorhinidius</i> Legalov, 2003										
<i>pauxillus</i> (Germar, 1824)	UD	DR	DE	LD	P	MV				
<i>Neocoenorhinus</i> subgen. <i>Neocoenorhinus</i> Voss, 1952										
<i>germanicus</i> (Herbst, 1797)	UD	DR								
<i>Rhynchites</i> D. H. Schneider, 1791	UD	DR	DE	LD	P			E		
<i>bacchus</i> (Linnaeus, 1758)										
<i>Tatianaerhynchites</i> Legalov, 2002										

<i>aequatus</i> (Linnaeus, 1767)	UD	DR	DE	LD						
<i>Teretriorhynchites</i> Voss, 1938										
<i>caeruleus</i> (DeGeer, 1775)	UD	DR	DE	LD	P					
<i>pubescens</i> (Fabricius, 1775)							SS	E		
ATTELABIDAE Billberg, 1820										
<i>Apoderus</i> Olivier, 1807										
<i>coryli</i> (Linnaeus, 1758)	UD	DR	DE							
<i>Attelabus</i> Linnaeus, 1758										
<i>nitens</i> (Scopoli, 1763)	UD	DR	DE							
APIONIDAE Schoenherr, 1823										
Apionini Schoenherr, 1823										
<i>Apion</i> Herbst, 1797										
* <i>cruentatum</i> Walton, 1844	UD	DR	DE	LD						
<i>frumentarium</i> Linnaeus, 1758	UD	DR	DE	LD	P	MV				
* <i>haematodes</i> <i>haematodes</i> Kirby, 1808	UD	DR	DE	LD						
<i>rubiginosum</i> Grill, 1893				LD						
Aplemonini Kissenger, 1968										
<i>Perapion</i> subgen. <i>Eroosapion</i> Ehret, 1994										
<i>lemoroi</i> (C. N. F. Brisout de Barneville, 1880)		DR		LD						
<i>Perapion</i> subgen. <i>Hemiperapion</i> Wagner, 1910										
<i>jacobsoni</i> (Wagner, 1910)										BL
<i>Perapion</i> (subgen. <i>Perapion</i>) Wagner, 1907										
<i>affine</i> (Kirby, 1808)				LD	P					
<i>connexum</i> (Schilsky, 1902)	UD	DR	DE	LD	P					
<i>curtiostre</i> (Germar, 1817)				LD	P					
* <i>marchicum</i> (Herbst, 1797)				LD						
* <i>myochroum</i> (Schilsky, 1902)										BL
<i>oblongum</i> Gyllenhal, 1839				LD						
<i>Pseudaplemonus</i> Wagner, 1930										
<i>artemisiae</i> (Moravitz, 1861)							E	SL	BL	
<i>externepunctatus</i> (Desbrochers des Loges, 1875)								SL	BL	
Aspidapiini Alonso-Zarazaga, 1990										
<i>Alocentron</i> subgen. <i>Alocentron</i> Schilsky, 1901										
<i>curviostre</i> (Gyllenhal, 1833)	UD	DR	DE	LD	P					
<i>Aspidapion</i> subgen. <i>Aspidapion</i> Schilsky, 1901										
<i>radiolus radiolus</i> (Marsham, 1802)	UD	DR	DE	LD	P					
<i>validum</i> (Germar, 1817)	UD	DR	DE	LD	P	MV	SS			
<i>Aspidapion</i> subgen. <i>Koestlinia</i> Alonso-Zarazaga, 1990										
<i>aeneum</i> (Fabricius, 1775)				LD						

	UD	DR	DE	LD	P	MV					
Ceratapiini Alonso-Zarazaga, 1990											
<i>Ceratapion</i> subgen. <i>Acanephodus</i> Alonso-Zarazaga, 1990											
<i>onopordi onopordi</i> (Kirby, 1808)	UD	DR	DE	LD	P	MV					
<i>Ceratapion</i> subgen. <i>Angustapion</i> Wanat, 1995					LD						
<i>transsylvanicum</i> (Schilsky, 1906)				LD							
<i>Ceratapion</i> subgen. <i>Ceratapion</i> Schilsky, 1901											
<i>armatum</i> (Gerstaecker, 1854)		DR									
<i>carduorum</i> (Kirby, 1808)	UD	DR	DE	LD	P	MV	SS	E	SL	BL	
<i>gibbirostre</i> (Gyllenhal, 1813)	UD	DR	DE	LD	P						
<i>Ceratapion</i> subgen. <i>Clementiellus</i> Alonso-Zarazaga, 1990											
<i>orientale</i> (Gerstaecker, 1854)	UD	DR	DE	LD							
<i>Ceratapion</i> subgen. <i>Echinostroma</i> Alonso-Zarazaga, 1990											
<i>penetrans penetrans</i> (Germar, 1817)		DR	DE	LD	P						
<i>Diplapion</i> Reitter, 1916											
<i>confluens</i> (Kirby, 1808)			DE	LD	P	MV					
<i>detritum</i> (Mulsant & Rey, 1858)		DR	DE	LD							
<i>sareptanum</i> (Desbrochers des Loges, 1867)						MV		E	SL		
<i>stolidum</i> (Germar, 1817)	UD	DR	DE	LD	P	MV					
<i>Omphalapion</i> Schilsky, 1901											
<i>hookerorum</i> (Kirby, 1808)				LD	P	MV	SS				
<i>Taphrotopium</i> Reitter, 1916											
<i>steveni</i> (Gyllenhal, 1839)			DE				SS	E			
<i> sulcifrons</i> (Herbst, 1797)				LD	P						
Exapiini Alonso-Zarazaga, 1990											
<i>Exapion</i> Bedel, 1887											
i corniculatum (Germar, 1817)		DR	DE	LD							
i difficile (Herbst, 1797)	UD	DR	DE	LD							
Kalcapiini Alonso-Zarazaga, 1990											
<i>Kalkapion</i> Schilsky, 1906											
<i>pallipes</i> (Kirby, 1808)				LD							
<i>Squamapion</i> Bokor, 1923											
<i>elongatum</i> (Germar, 1817)	UD	DR	DE	LD							
<i>Taeniapion</i> Schilsky, 1906											
<i>urticarium urticarium</i> (Herbst, 1784)	UD	DR	DE	LD	P						
Malvapiini Alonso-Zarazaga, 1990											
<i>Malvapion</i> A. Hoffmann, 1958											
<i>malvae</i> (Fabricius, 1775)	UD	DR									
<i>Pseudapion</i> Schilsky, 1906											
<i>fulvirostre</i> (Gyllenhal, 1833)	UD	DR	DE	LD	P	MV	SS	E	SL	BL	

		DR		LD				E	SL	BL
<i>rufirostre</i> (Fabricius, 1775)										
<i>Rhopalapion</i> Schilsky, 1906										
<i>longirostre</i> (Olivier, 1807)	UD	DR	DE	LD	P	MV	SS	E		
Oxystomatini Alonso-Zarazaga, 1990										
<i>Catapion</i> Schilsky, 1906										
<i>seniculus</i> (Kirby, 1808)	UD	DR	DE	LD						
<i>Cyanapion</i> subgen. <i>Bothryorrhynchapion</i> Borok, 1923										
* <i>afer</i> (Gyllenhal, 1833)	UD		DE							
<i>Cyanapion</i> subgen. <i>Cyanapion</i> Borok, 1923										
<i>colymbinum</i> (Germar, 1817)			DE	LD		MV				
<i>Eutrichapion</i>										
<i>mystriophorum</i> Alonso-Zarasaga, 1994				LD						
<i>viciae</i> (Paykull, 1800)	UD	DR	DE	LD	P					
<i>Hemitrichapion</i> subgen. <i>Dimesomyops</i> Voss, 1959										
<i>pavidum</i> (Germar, 1817)	UD	DR	DE	LD	P					
<i>Hemitrichapion</i> subgen. <i>Tinocysba</i> Alonso- Zarazaga, 1990										
<i>reflexum</i> (Gyllenhal, 1833)				LD						
<i>Holotrichapion</i> subgen. <i>Apiops</i> Alonso- Zarazaga, 1990										
<i>pisi</i> (Fabricius, 1801)			DE	LD						
<i>pullum</i> (Gyllenhal, 1833)		DR	DE	LD	P	MV	SS	E	SL	BL
<i>Isochnopterapion</i> subgen. <i>Isochnopterapion</i> Bokor, 1923										
<i>loti</i> (Kirby, 1808)	UD	DR	DE	LD	P	MV				
<i>Loborhynchapion</i> Györffy, 1956										
<i>amethystinum</i> (Miller, 1857)				LD			SS			
<i>Mesotrichapion</i> Györffy, 1956										
<i>punctirostre</i> (Gyllenhal, 1839)			DE	LD		MV				
<i>Oxystoma</i> Duméril, 1805										
<i>cerdo</i> (Gerstaecker, 1854)				LD						
<i>pomonae</i> (Fabricius, 1798)			DE	LD						
<i>Stenopterapion</i> subgen. <i>Stenopterapion</i> Bokor, 1923										
<i>intermedium</i> (Eppelsheim, 1875)			DE	LD	P					
<i>meliloti</i> (Kirby, 1808)	UD	DR	DE	LD	P	MV	SS	E		
<i>tenue</i> (Kirby, 1808)	UD	DR	DE	LD	P	MV	SS	E		
<i>Betulapion</i> Ehret, 1994										
* <i>simile simile</i> (Kirby, 1811)				LD						
Piezotrachelini Voss, 1959										
<i>Protapton</i> Schilsky, 1908										
<i>apricans</i> (Herbst, 1797)	UD		DE	LD						
<i>assimile assimile</i> (Kirby, 1808)	UD	DR	DE	LD						
<i>filiostre</i> (Kirby, 1808)			DE	LD	P			E	SL	
<i>fulvipes</i> (Geoffroy, 1785)		DR	DE	LD						

<i>nigritarse</i> (Kirby, 1808)	UD	DR	DE	LD		MV			
<i>ononidis</i> (Gyllenhal, 1827)			DE	LD					
* <i>ruficrus</i> (Germar, 1817)	UD	DR	DE						
<i>trifolii</i> (Linnaeus, 1768)				LD	P				
<i>Pseudoprotapion</i> Ehret, 1990									
<i>astragali astragali</i> (Paykull, 1800)	UD				P				
<i>elegantulum</i> (Germar, 1818)		DR	DE	LD		MV			
<i>ergenense</i> (Becker, 1864)			DE	LD					
NANOPHYIDAE Gistel, 1848									
CORIMALIINAE Alonso-Zarazaga, 1989									
<i>Allomalia</i> Alonso-Zarazaga, 1989									
<i>setulosa</i> (Tournier, 1868)							SL		
<i>Corimalia</i> Gozis, 1885									
<i>fausti fausti</i> (Reitter, 1890)						E	SL	BL	
<i>heleneae</i> Korotyaev & Zherikhin, 1996					SS	E	SL	BL	
<i>Hypophyes</i> Reitter, 1916						SS	E	SL	BL
<i>minutissimus</i> (Tournier, 1868)									
<i>Titanomalia</i> Alonso-Zarazaga, 1989									
<i>komaroffi</i> (Faust, 1877)								BL	
NANOPHYINAE Gistel, 1848									
<i>Dieckmanniellus</i> Alonso-Zarazaga, 1989									
* <i>gracilis</i> (Redtenbacher, 1847)			DE				E	SL	BL
* <i>nitidulus</i> (Gyllenhal, 1838)								SL	
<i>Nanomimus</i> Alonso-Zarazaga, 1989									
<i>circumscriptus</i> (Aubé, 1864)		DE	LD						
* <i>hemisphericus</i> (Olivier, 1807)			LD						
<i>Nanophyes</i> Schoenherr, 1838									
* <i>brevis brevis</i> Boheman, 1845	DR	DE	LD						
<i>globiformis</i> Kiesenwetter, 1864			LD						
<i>marmoratus marmoratus</i> (Goeze, 1777)	UD	DR	DE	LD	P	MV			
BRACHYCERIDAE Billberg, 1820									
<i>Brachycerus</i> Olivier, 1789									
<i>foveicollis</i> Gyllenhal, 1833			DE	LD					
* <i>sinuatus</i> Olivier, 1807			DE	LD	P	MV			
DRYOPHTHORIDAE Schoenherr, 1825									
SPHENOPHORINAE Schoenherr, 1833									
Litosomini Lacordaire, 1865									
<i>Sitophilus</i> Schoenherr, 1838									
<i>granarius</i> (Linnaeus, 1758)	UD	DR	DE	LD	P	MV	SS	E	SL
<i>oryzae</i> (Linnaeus, 1763)	UD	DR	DE	LD	P	MV	SS	E	SL
Sphenophorini Lacordaire, 1865									
<i>Sphenophorus</i> Schoenherr, 1838									

<i>abbreviatus</i> (Fabricius, 1787)	UD	DR	DE	LD	P	MV	SS	E	SL	BL
<i>striatopunctatus</i> (Goeze, 1777)	UD			LD						
ERIRHINIDAE Schoenherr, 1826										
Arthrostenini Reitter, 1913										
<i>Arthrostenus</i> Schoenherr, 1826										
* <i>fullo</i> Boheman, 1836	UD	DR	DE				SS	E	SL	BL
Erirhinini Schoenherr, 1825										
<i>Bagoopsis</i> Faust, 1881										
* <i>globicollis</i> (Fairmaire, 1863)			DE	LD						
* <i>volgensis</i> Faust, 1881							SS	E		
<i>Icaris</i> Tournier, 1874										
* <i>sparganii sparganii</i> (Gyllenhal, 1835)			DE	LD		MV	SS	E	SL	BL
<i>Lepidonotaris</i> Zumpt, 1929										
<i>petax</i> (C. R. Sahlberg, 1823)			DE		P			E	SL	BL
<i>Notaris</i> Germar, 1817										
* <i>acridulus acridulus</i> (Linnaeus, 1758)	UD		DE	LD						
* <i>scirpi</i> (Fabricius, 1792)	UD	DR	DE	LD	P	MV	SS	E	SL	BL
<i>Picia</i> Tournier, 1895										
* <i>sinuatocollis</i> (Faust, 1885)									SL	
<i>Procas</i> Stephens, 1831										
* <i>picipes steveni</i> (Krynicki, 1832)		DR	DE	LD	P	MV				
<i>Thryogenes</i> Bedel, 1884										
* <i>festucae</i> (Herbst, 1795)	UD	DR	DE	LD			SS	E		BL
* <i>nereis</i> (Paykull, 1800)		DR	DE							
<i>Tournotaris</i> Alonso-Zarazaga & Lyal, 1999										
* <i>bimaculata</i> (Fabricius, 1787)	UD	DR	DE	LD						
Tanysphyrini Gistel, 1817										
<i>Tanysphyrus</i> Germar, 1817										
* <i>lemnae</i> (Paykull, 1792)			DE	LD					SL	BL
CURCULIONIDAE Latreille, 1802										
CURCULIONINAE Latreille, 1802										
Acalyptini Thomson, 1859										
<i>Acalyptus</i> Schoenherr, 1833										
* <i>carpini</i> (Fabricius, 1792)			DE							
<i>Acentrus</i> Desmarest, 1839										
<i>histrio</i> (Boheman, 1837)		DR		LD	P	MV	SS	E	SL	BL
Anthonomini C. G. Thomson, 1859										
<i>Anthonomus</i> subgen. <i>Anthomorphus</i> Weise, 1883										
* <i>pinivorax</i> Silfverberg, 1977				LD						
<i>Anthonomus</i> subgen. <i>Anthonomidius</i> Reitter, 1915										
<i>germanicus</i> Dieckmann, 1968		DR	DE	LD						

<i>rubripes</i> Gyllenhal, 1835		DR	DE	LD		MV	SS		
<i>Anthonomus</i> subgen. <i>Anthonomus</i> Germar, 1817									
* <i>bituberculatus</i> C. G. Thomson, 1868			DE						
* <i>pedicularius</i> (Linnaeus, 1758)			DE		P				
<i>pomorum</i> (Linnaeus, 1758)	UD	DR	DE	LD	P				
<i>rubi</i> (Herbst, 1795)	UD	DR	DE	LD	P	MV			
<i>ulmi</i> (DeGeer, 1775)		DR							
<i>Anthonomus</i> subgen. <i>Furcipes</i> Desbrochers des Loges, 1868									
<i>rectirostris</i> (Linnaeus, 1758)	UD								
<i>Bradybatus</i> subgen. <i>Bradybatus</i> Germar, 1824									
* <i>kellneri</i> Bach, 1854	UD	DR	DE						
* <i>seriesetosus</i> Petri, 1912				LD					
Cionini Schoenherr, 1825									
<i>Cionus</i> Clairville, 1798									
* <i>hortulanus</i> (Geoffroy, 1785)	UD	DR	DE	LD	P	MV			
* <i>olivieri</i> Rosenschoeld, 1838	UD	DR	DE	LD	P	MV			
* <i>scrophulariae</i> (Linnaeus, 1758)		DR	DE						
<i>thapsus thapsus</i> (Fabricius, 1792)			DE	LD					
* <i>tuberculosus</i> (Scopoli, 1763)	UD			LD					
Curculionini Latreille, 1802									
<i>Archarius</i> subgen. <i>Archarius</i> Gistel, 1856									
<i>pyrrhoceras</i> (Marsham, 1802)	UD	DR	DE	LD					
<i>salicivorus</i> (Paykull, 1792)			DE	LD					
<i>Curculio</i> Linnaeus, 1758									
<i>betulae</i> (Stephens, 1831)				LD					
* <i>elephas</i> (Gyllenhal, 1835)	UD	DR	DE	LD					
<i>glandium</i> Marsham, 1802	UD	DR	DE	LD	P	MV	SS		
<i>nucum</i> Linnaeus, 1758		DR	DE						
<i>rubidus</i> Gyllenhal, (1835)			DE	LD					
<i>venosus venosus</i> (Gravenhorst, 1807)				LD					
<i>villosus</i> Fabricius, 1781			DE						
Ellescini C.G. Thomson, 1859									
<i>Dorytomus</i> subgen. <i>Dorytomus</i> Germar, 1817									
* <i>dejeani</i> Faust, 1883			DE	LD		SS			
* <i>filirostris</i> (Gyllenhal, 1835)			DE	LD					
<i>hirtipennis</i> Bedel, 1884			DE	LD				BL	
* <i>ictor</i> (Herbst, 1795)			DE	LD	P	MV			
* <i>longimanus</i> Forster, 1771			DE	LD					
* <i>melanophthalmus</i> (Paykull, 1792)		DR	DE						
* <i>nebulosus</i> (Gyllenhal, 1835)	UD		DE	LD					
* <i>suratus</i> (Gyllenhal, 1835)			DE	LD					
* <i>tremulae</i> (Fabricius, 1787)	UD	DR	DE	LD					

<i>Ellescus</i> Dejean, 1821											
* <i>infirmus</i> (Herbst, 1795)			DE								
<i>scanicus</i> (Paykull, 1792)				LD							
Itini Reitter, 1913											
<i>Ita</i> Tournier, 1878											
* <i>kirschii</i> Tournier, 1878								E			
<i>Geranorhinus</i> Chevrolat, 1860											
<i>virens</i> Faust, 1885								E	SL	BL	
Mecinini Gistel, 1848											
<i>Cleopomiarus</i> Pierse, 1919											
* <i>distinctus</i> (Boheman, 1845)		DR									
<i>graminis</i> (Gyllenhal, 1813)	UD	DR	DE	LD							
<i>Gymnetron</i> Schoenherr, 1825											
<i>hoferi</i> Desbrochers des Loges, 1869			DE			MV	SS	E		BL	
* <i>melanarium</i> (Germar, 1821)	UD	DR	DE	LD	P	MV					
<i>niloticum</i> Kirsch, 1881		DR	DE	LD							
* <i>rotundicolle</i> (Gyllenhal, 1838)	UD			LD	P	MV					
<i>sauromatum</i> Arzanov, 2006		DR					SS				
* <i>terminassianae</i> Smreczyński, 1975					P						
* <i>veronicae</i> (Germar, 1821)			DE	LD							
* <i>villosum</i> Gyllenhal, 1838			DE	LD		MV	SS	E			
* <i>vittipenne</i> Marseul, 1876				LD							
<i>Mecinus</i> Germar, 1821											
* <i>collaris</i> Germar, 1821		DR	DE	LD		MV	SS			BL	
<i>henrici</i> (Arzanov, 1991)									SL		
* <i>heydenii</i> Wencker, 1866		DR	DE	LD	P						
<i>ictericus</i> (Gyllenhal, 1838)	UD	DR	DE	LD							
<i>janthinus</i> Germar, 1821	UD	DR	DE	LD	P	MV	SS	E	SL	BL	
* <i>labilis</i> (Herbst, 1795)	UD	DR	DE	LD	P	MV	SS	E	SL	BL	
<i>marina</i> (Korotyaev, 1984)						MV		E	SL	BL	
<i>pascuorum</i> (Gyllenhal, 1813)	UD	DR	DE	LD	P	MV	SS	E			
<i>pirazzolii</i> (Stierlin, 1867)	UD	DR	DE	LD							
* <i>plantaginis</i> (Eppelsheim, 1875)	UD	DR	DE	LD	P						
<i>pyraster</i> (Herbst, 1795)			DE	LD	P	MV	SS	E	SL	BL	
<i>tanaiticus</i> Arzanov, 2000			DE								
<i>Miarus</i> Schoenherr, 1826											
* <i>ajugae</i> (Herbst, 1795)				LD							
<i>Rhinusa</i> Stephens, 1829											
* <i>antirrhini</i> (Paykull, 1800)			DE	LD				E	SL	BL	
* <i>asellus</i> (Gravenhorst, 1807)		DR	DE	LD							
<i>collina</i> (Gyllenhal, 1813)			DE	LD							
* <i>florum</i> (Rübsaamen, 1895)		DR					SS	E			
* <i>linariae</i> (Panzer, 1795)	UD	DR	DE	LD	P	MV	SS				
* <i>melas</i> Boheman, 1838			DE	LD							
* <i>neta</i> (Germar, 1821)	UD	DR	DE	LD	P	MV	SS				

<i>*pilosa</i> (Gyllenhal, 1838)			DE	LD						
<i>tetra</i> (Fabricius, 1792)	UD	DR	DE	LD	P	MV	SS			
Rhamphini Rafinesque, 1815										
<i>Isochnus</i> C. G. Thomson, 1859										
<i>sequensi</i> (Stierlin, 1894)		DR	DE			MV				
<i>Orchestes</i> subgen. <i>Alyctus</i> C. G. Thomson, 1859										
<i>testaceus</i> (O. F. Müller, 1776)	UD			LD						
<i>Orchestes</i> subgen. <i>Orchestes</i> Illiger, 1798										
<i>*alni</i> (Linnaeus, 1758)	UD									
<i>hortorum</i> (Fabricius, 1792)			DE							
<i>*mutabilis</i> Boheman, 1843	UD						SS			
<i>quedenfeldtii</i> Gerharrdt, 1865				LD						
<i>quercus</i> (Linnaeus, 1758)				LD						
<i>Pseudorchemes</i> Bedel, 1894										
<i>asrtacanicus</i> (Tournier, 1873)										BL
<i>cinereus</i> (Fåhraeus, 1843)			DE	LD						
<i>*circumvistulanus</i> (Białooki, 1997)	UD		DE	LD	P					
<i>ermischii</i> (Dieckmann, 1958)	UD	DR	DE	LD						
<i>*horioni</i> (Dieckmann, 1958)	UD		DE	LD						
<i>*smreczynskii</i> (Dieckmann, 1958)	UD	DR	DE	LD						
<i>Rhamphus</i> Clairville, 1798										
<i>oxyacanthae</i> (Marsham, 1802)	UD			LD		MV	SS			
<i>pulicarius</i> (Herbst, 1795)			DE	LD						
<i>Tachyerges</i> Schoenherr, 1825										
<i>salicis</i> (Linnaeus, 1758)				LD						
<i>stigma</i> (Germar, 1821)			DE	LD						
Smicronychini Seidliz, 1891										
<i>Smicronyx</i> Schoenherr, 1843										
<i>*coecus</i> (Reich, 1797)		DR	DE	LD		MV	SS			
<i>*ungermanniae</i> (Reich, 1797)	UD	DR	DE	LD	P	MV	SS	E	SL	BL
<i>kubanicus</i> Reitter, 1888		DR	DE	LD		MV	SS			
<i>*nebulosus</i> Tounier, 1874							SS			
<i>*reichii</i> (Gyllenhal, 1835)				LD			SS			
<i>scops</i> Tournier, 1874								E		
<i>*smreczynskii</i> F. Solari, 1952	UD	DR	DE	LD	P	MV	SS	E	SL	BL
<i>syriacus</i> Faust, 1887		DR	DE	LD		MV	SS			
Storeini Lacordaire, 1863										
<i>Pachytychius</i> Jekel, 1861										
<i>sparsutus</i> (Olivier, 1807)		DR	DE	LD	P	MV				
<i>*transcaucasicus</i> Pic, 1913				LD		MV				
Styphlini Jekel, 1861										
<i>Paraphilernus</i> Desbrochers des Loges, 1892										
<i>bilunulatus</i> (Desbrochers des Loges,		DR	DE	LD	P	MV				

1870)											
<i>Philernus</i> Schoenherr, 1835											
<i>farinosus</i> (Gebler, 1830)						MV	SS	E	SL	BL	
<i>ponticus</i> Korotyaev, 1979						MV			SL	BL	
<i>Pseudostyphlus</i> Tournier, 1874											
<i>pillumus</i> (Gyllenhal, 1835)				LD		MV	SS				
Tychiini C. G. Thomson, 1859											
<i>Lignyodes</i> Dejean, 1835											
* <i>bischoffi</i> (Blatchley, 1916)				LD							
<i>enucleatus</i> (Panzer, 1798)				LD			SS				
<i>suturatus</i> (Fairmaire, 1860)	UD		DE	LD			SS				
<i>Sibinia</i> subgen. <i>Dichotychius</i> Bedel, 1885											
<i>beckeri</i> Desbrochers des Loges, 1873		DR						E	SL	BL	
* <i>bipunctata</i> Kirsch, 1871										BL	
<i>planiuscula planiuscula</i> (Desbrochers des Loges, 1873)								E	SL	BL	
<i>staticis</i> (Becker, 1864)									SL	BL	
<i>subirrorata</i> Faust, 1885						MV		E	SL	BL	
<i>Sibinia</i> subgen. <i>Sibinia</i> Germar, 1817											
<i>femoralis</i> Gyllenhal, 1824										BL	
<i>pellucens</i> (Scopoli, 1772)	UD	DR	DE	LD	P	MV	SS	E	SL	BL	
* <i>phalerata</i> Gyllenhal, 1835			DE	LD		MV					
* <i>primita</i> (Herbst, 1795)			DE								
<i>subelliptica</i> Desbrochers des Loges, 1873	UD	DR	DE	LD	P	MV	SS	E	SL	BL	
<i>tibialis</i> Gyllenhal, 1835			DE								
<i>unicolor</i> Fåhraeus, 1843			DE	LD	P	MV	SS	E	SL	BL	
* <i>variata</i> Gyllenhal, 1835			DE	LD							
<i>viscariae</i> (Linnaeus, 1760)		DR	DE	LD	P	MV		E			
<i>Tychius</i> Germar, 1817											
<i>affinis</i> Becker, 1864				LD	P						
<i>albilaterus</i> Stierlin, 1864			DE	LD	P	MV					
<i>albolineatus</i> Motschulsky, 1860	UD	DR	DE	LD							
* <i>argentatus</i> Chevrolat, 1859			DE					E			
<i>astragali</i> Becker, 1862			DE		P	MV					
<i>aureolus</i> Kiesenwetter, 1852	UD	DR	DE	LD	P	MV	SS	E	SL	BL	
* <i>bedeli</i> Faust, 1885									SL	BL	
<i>breviusculus</i> Desbrochers des Loges, 1873		DR	DE	LD	P	MV					
<i>crassirostris</i> Kirsch, 1871		DR	DE	LD				E			
* <i>flavus</i> Becker, 1864	UD	DR	DE	LD	P	MV	SS	E	SL		
* <i>hauseri</i> Faust, 1889								E		BL	
<i>junceus</i> (Reich, 1797)		DR	DE	LD	P	MV	SS				
<i>karkaralensis</i> Bajtenov, 1974										BL	
<i>medicaginis</i> C. N. F. Brisout de Barneville, 1863	UD	DR	DE	LD	P	MV	SS	E	SL	BL	
<i>meliloti</i> Stephens, 1831	UD	DR	DE	LD	P	MV	SS	E	SL	BL	
<i>moleslus</i> Faust, 1891			DE	LD							

<i>morawitzi</i> Becker, 1864								E	SL	BL
<i>picrostris</i> (Fabricius, 1787)	UD	DR	DE	LD	P	MV	SS	E	SL	BL
<i>polylineatus</i> (Germar, 1824)			DE							
* <i>pusillus</i> Germar, 1842			DE	LD						
<i>quinquepunctatus</i> <i>quinquepunctatus</i> (Linnaeus, 1758)	UD	DR	DE	LD	P	MV	SS			
* <i>rufirostris</i> Schoenherr, 1832								E		
* <i>squamulatus</i> Gyllenhal, 1835			DE	LD						
<i>stephensi</i> Gyllenhal, 1835		DR	DE	LD						
<i>subsulcatus</i> Tournier, 1874	UD	DR	DE		P	MV				
* <i>sulphureus</i> Faust, 1881									SL	
<i>tridentinus</i> Peneke, 1922				LD	P		SS			
* <i>trivialis</i> Boheman, 1843			DE			MV	SS			
* <i>uralensis</i> Pic, 1902		DR	DE	LD	P					
<i>winkleri</i> (Franz, 1940)								E		
BAGOINAE Thomson, 1859										
<i>Bagous</i> Germar, 1817										
<i>alismatis</i> (Marsham, 1802)	UD	DR	DE	LD	P	MV	SS	E	SL	BL
* <i>argillaceus</i> Gyllenhal, 1836	UD	DR	DE	LD	P	MV	SS	E	SL	BL
* <i>biimpressus</i> Fähræus, 1845								E		
<i>collignensis</i> (Herbst, 1797)				LD						
* <i>diglyptus</i> Boheman, 1845										BL
<i>elegans</i> (Fabricius, 1801)										BL
* <i>frivaldszkyi</i> Tournier, 1874				LD						
* <i>glabrirostris</i> (Herbst, 1795)										BL
* <i>limosus</i> (Gyllenhal, 1827)										BL
* <i>lutosus</i> Gyllenhal, 1813			DE	LD						
* <i>lutulentus</i> (Gyllenhal, 1813)						MV				
<i>minutissimus</i> Faust, 1887								E		
<i>nodulosus</i> Gyllenhal, 1836	UD	DR	DE	LD	P	MV				
<i>peregrinus</i> Gratschev, 1993			DE							
* <i>petro</i> (Herbst, 1795)										BL
* <i>puncticollis</i> Boheman, 1845										BL
<i>robustus</i> H. Brisout de Barneville, 1863				LD						
* <i>subcarinatus</i> Gyllenhal, 1836		DR	DE	LD				E	SL	BL
* <i>tubulus</i> Caldara & O'Brien, 1994		DR	DE	LD						
<i>validus</i> Rosenhauser, 1847	UD	DR	DE			MV				
BARIDINAE Schoenherr, 1836										
Apostasimerini Schoenherr, 1844										
<i>Limnobaris</i> Bedel, 1885										
<i>dolorosa</i> (Goeze, 1777)	UD	DR	DE	LD	P	MV	SS	E	SL	BL
<i>t-album</i> (Linnaeus, 1758)	UD	DR	DE	LD	P	MV	SS	E	SL	BL
Baridini Schoenherr, 1836										
<i>Aulacobaris</i> Desbrochers des Loges, 1892										
<i>angusta angusta</i> (Brullé, 1832)		DR	DE	LD	P	MV				
<i>chlorizans</i> (Germar, 1824)				LD						

<i>coerulescens coerulescens</i> (Scopoli, 1763)	UD	DR	DE	LD	P	MV	SS	E		
<i>concinna</i> (Bohemian, 1844)		DR	DE	LD						
<i>janthina</i> (Bohemian, 1836)	UD	DR	DE	LD	P	MV	SS	E	SL	BL
<i>lepidii</i> (Germar, 1824)			DE	LD	P	MV	SS	E		
<i>violaceomicans violaceomicans</i> (A. & F. Solaro, 1905)		DR	DE	LD				E	SL	
<i>Baris</i> Germar, 1817										
<i>analis analis</i> (Olivier, 1791)	UD	DR	DE	LD	P	MV	SS	E	SL	BL
<i>artemisiae</i> (Herbst, 1794)	UD	DR	DE	LD	P	MV	SS	E		
<i>convexicollis</i> (Bohemian, 1836)			DE	LD	P			E	SL	
<i>limbata</i> (H. Brisout de Barneville, 1870)								E	SL	
<i>memnonia memnonia</i> (Bohemian, 1836)						MV	SS			
<i>nesapia</i> Faust, 1887		DR	DE							
<i>spitzyi</i> Hochhuth, 1847	UD	DR	DE	LD	P	MV		E		
<i>sulcata</i> (Bohemian, 1836)		DR		LD		MV				
<i>Cosmobaris</i> Casey, 1920										
<i>scolopacea</i> (Germar, 1819)	UD	DR	DE	LD	P	MV	SS	E	SL	BL
<i>Elasmobaris</i> Reitter, 1913										
* <i>alboguttata alboguttata</i> (H. Brisout de Barneville, 1870)										BL
* <i>signifera</i> (Faust, 1881)								E	SL	BL
<i>Eremobaris</i> Zaslavskij, 1956										
<i>picturata</i> (Ménétrier, 1849)			DE							
<i>Labiaticola</i> Alonso-Zarazaga & Lyal, 1999										
<i>atricolor</i> (Bohemian, 1844)		DR								
<i>melas</i> (Bohemian, 1836)	UD	DR	DE	LD	P	MV	SS	E	SL	BL
<i>sibiricus</i> (Faust, 1890)			DE	LD			SS			
<i>Malvaevora</i> Zaslavskij, 1956										
<i>timida timida</i> (Rossi, 1792)	UD	DR	DE	LD	P	MV	SS	E	SL	BL
<i>Melanobaris</i> Alonso-Zarazaga & Lyal, 1999										
* <i>atramentaria</i> (Bohemian, 1836)			DE	LD						
<i>carbonaria</i> (Bohemian, 1836)				LD	P		SS			
<i>dalmatina</i> (H. Brisout de Barneville, 1870)			DE							
<i>hochhuthi</i> (Faust, 1882)		DR	DE	LD				E		
* <i>nigritarsis</i> (Bohemian, 1836)			DE	LD	P					
<i>semistriata</i> (Bohemian, 1836)	UD	DR	DE	LD	P	MV	SS	E	SL	BL
<i>Ulobaris</i> Reitter, 1895										
<i>loricata</i> (Bohemian, 1836)	UD		DE	LD				E	SL	BL
CEUTORHYNCHINAE Gistel, 1848										
Amalini H. Wagner, 1936										
<i>Amalus</i> Schoenherr, 1825										
<i>scortillum</i> (Herbst, 1795)			DE	LD	P	MV	SS	E	SL	BL
Ceutorhynchini Gistel, 1848										
<i>Boragosiocalus</i> Deickmann, 1975										
<i>rinderae</i> (Becker, 1864)				LD						

<i>Calosirus</i> C. G. Thomson, 1859											
<i>ovulum</i> (Schultze, 1897)				LD							
<i>Ceutorhynchus</i> Germar, 1824											
<i>aeneicollis</i> Germar, 1824					P	MV	SS	E	SL	BL	
<i>arator</i> Gyllenhal, 1837			DE								
<i>arnoldii</i> (Korotyaev, 1980)			DE	LD							
<i>assimilis</i> (Paykull, 1792)			DE	LD	P						
<i>buniadis</i> Penecke, 1928				LD							
* <i>canaliculatus</i> C. N. F. Brisout de Barneville, 1869			DE								
<i>cardariae</i> Korotyaev, 1992	DR	DE									
<i>carinatus</i> Gyllenhal, 1837							SS				
<i>chalybaeus</i> Germar, 1824			DE	LD	P	MV	SS				
<i>chlorophanus</i> Rouget, 1857	UD	DR	DE	LD							
<i>circassicus</i> Schultze, 1900				LD							
* <i>coarctatus</i> Gyllenhal, 1837											BL
<i>confusus</i> Schultze, 1903			DE	LD		MV	SS				
* <i>constrictus</i> (Marsham, 1802)			DE								
<i>contractus</i> Marsham, 1802.			DE	LD	P	MV	SS				
* <i>difficilis</i> Schultze, 1898					P						
<i>erysimi</i> (Fabricius, 1787)	UD	DR	DE	LD	P	MV	SS	E	SL	BL	
<i>fabrilis</i> Faust, 1887			DE			MV					
<i>fallax</i> Boheman, 1845		DR	DE	LD							
<i>gallorhenanus</i> F. Solari, 1949		DR	DE	LD	P						
<i>granulicollis</i> C. G. Thomson, 1865				LD		MV					
<i>griseus</i> C. N. F. Brisout de Barneville, 1869											BL
<i>hampei</i> C. N. F. Brisout de Barneville, 1869				LD				E	SL	BL	
<i>ignitus</i> Germar, 1824		DR	DE	LD	P						
<i>inaffectatus</i> Gyllenhal, 1837		DR	DE	LD		MV					
<i>languidus</i> Schultze, 1902								E	SL	BL	
<i>lothari</i> Kryzhanovskaya, 1993		DR	DE	LD							
<i>merkli</i> Korotyaev, 2001				LD							
<i>nanus</i> Gyllenhal, 1837				LD		MV	SS				
<i>niyazii</i> Hoffmann, 1957		DR		LD							
<i>pallidactylus</i> (Marsham, 1802)		DR		LD							
<i>pervicax</i> Weise, 1883				LD			SS				
<i>piceolatus</i> (C. N. F. Brisout de Barneville, 1883)			DE	LD							
<i>pictarsis</i> Gyllenhal, 1837		DR	DE	LD							
<i>posthumus</i> Germar, 1824				LD							
<i>psoropygus</i> Iablokoff-Khnzorian, 1971											BL
<i>pulvinatus</i> Gyllenhal, 1837	UD	DR	DE	LD	P	MV	SS	E	SL	BL	
<i>rapae</i> Gyllenhal, 1837		DR	DE	LD	P						
<i>rhenanus</i> (Schultze, 1895)				LD							
<i>roberti</i> Gyllenhal, 1837	UD	DR	DE	LD							
<i>sisymbrii</i> (Dieckmann, 1966)			DE	LD	P	MV	SS	E	SL	BL	
<i>sophiae</i> Gyllenhal, 1837	UD	DR	DE	LD	P	MV	SS	E	SL	BL	

<i>*sulcatus</i> C. N. F. Brisout de Barneville, 1869				LD						
<i>sulcicollis</i> (Paykull, 1800)			DE	LD						
<i>syrites</i> Germar, 1824		DR		LD			SS			
<i>talickyi</i> Korotyaev, 1980		DR	DE	LD			SS			
<i>turbatus</i> Schultze, 1903		DR	DE	LD				E		
<i>typhae</i> (Herbst, 1795)	UD	DR	DE	LD	P	MV	SS	E	SL	BL
<i>viridanus</i> Gyllenhal, 1837		DR	DE	LD						
<i>Coeliastes</i> Weise, 1883										
<i>lamii</i> (Fabricius, 1792)		DR	DE	LD						
<i>Datonychus</i> H. Wagner, 1944										
<i>*angulosus</i> (Boheman, 1845)	UD									
<i>*arquata</i> (Herbst, 1795)			DE							
<i>paszlawskyi</i> (Kuthy, 1890)				LD						
<i>Glocianus</i> Reitter, 1916										
<i>distinctus</i> (C. N. F. Brisout de Barneville, 1870)	UD	DR	DE	LD						
<i>herbsti</i> (Faust, 1895)			DE		P					
<i>*lethierryi</i> (C. N. F. Brisout de Barneville, 1866)					P					
<i>*moelleri</i> (C. G. Thomson, 1868)						MV				
<i>pilosellus</i> (Gyllenhal, 1837)				LD						
<i>*polystriatus</i> (Schultze, 1898)				LD		MV	SS			
<i>punctiger</i> (C. R. Sahlberg, 1835)		DR	DE	LD						
<i>steveni</i> (Boheman, 1845)						MV				
<i>Hadropontus</i> C. G. Thomson, 1859										
<i>trimaculatus</i> (Fabricius, 1775)				LD						
<i>Microplontus</i> H. Wagner, 1944										
<i>rugulosus</i> (Herbst, 1795)		DR	DE	LD						
<i>Mogulones</i> Reitter, 1916										
<i>*abbreviatulus</i> (Fabricius, 1792)	UD									
<i>*albosignatus</i> (Gyllenhal, 1837)			DE	LD	P	MV				
<i>amplipennis</i> (Schultze, 1897)				LD						
<i>asperifoliarum</i> (Gyllenhal, 1813)			DE	LD	P					
<i>austriacus</i> (C. N. F. Brisout de Barneville, 1869)		DR	DE	LD	P	MV				
<i>crucifer</i> (Pallas, 1771)	UD	DR		LD		MV				
<i>*dimidiatus</i> (Frivaldszky, 1865)				LD						
<i>*raphani</i> (Fabricius, 1792)		DR	DE	LD	P					
<i>trisignatus</i> (Gyllenhal, 1837)	UD		DE	LD						
<i>*venedicus</i> (Weise, 1879)				LD						
<i>Nedyus</i> Schoenherr, 1825										
<i>quadrimaculatus</i> (Linnaeus, 1758)	UD	DR	DE	LD	P					
<i>Neoglocianus</i> Dieckmann, 1972										
<i>maculaalba</i> (Herbst, 1795)		DR								
<i>Neoprohinus</i> Colonnelli, 2004										
<i>cinnamomeus</i> (Schultze, 1897)				LD						
<i>Phrydiuchus</i> Gozis, 1885										
<i>augusti</i> Colonnelli, 2003				LD						

<i>*topiarius</i> (Germar, 1824)		DR		LD						
<i>Platygasteronyx</i> Reitter, 1913										
<i>*solskyi</i> (Faust, 1885)		DR								
<i>Poophagus</i> Schoenherr, 1837										
<i>*hopffgarteni</i> Tournier, 1873		DR	DE							
<i>robustus</i> Faust, 1881			DE							
<i>sisymbrii</i> (Fabricius, 1777)			DE							
<i>Prisistus</i> Reitter, 1916										
<i>*caucasicus caucasicus</i> (Kirsch, 1878)								E		
<i>*obsoletus</i> (Germar, 1824)					MV	SS	E	SL	BL	
<i>suturalba</i> (Schultze, 1903)		DE								
<i>Ranunculiphilus</i> Dieckmann, 1970										
<i>faeculentus</i> (Gyllenhal, 1837)		DR		LD		MV				
<i>Sirocalodes</i> Voss, 1958										
<i>depressicollis</i> (Gyllenhal, 1813)	UD	DR	DE							
<i>quercicola</i> (Paykull, 1792)				LD		MV				
<i>Stenocarus</i> C.G. Thomson, 1859										
<i>cardui</i> Herbst, 1784									BL	
<i>ruficornis</i> (Stephens, 1831)				LD	P	MV	SS			
<i>Thamiocolus</i> Colonnelli, 1979										
<i>nubeculosus</i> (Gyllenhal, 1837)		DR	DE	LD		MV				
<i>*sahlbergi</i> (Sahlberg, 1845)		DR	DE	LD						
<i>signatus</i> (Gyllenhal, 1837)		DR	DE	LD		MV	SS			
<i>sinapis</i> (Desbrochers des Loges, 1893)	UD	DR	DE	LD	P	MV				
<i>uniformis</i> (Gyllenhal, 1837)	UD	DR	DE	LD	P	MV	SS	E	SL	BL
<i>viduatus</i> (Gyllenhal, 1813)		DR	DE		P					
<i>virgatus</i> (Gyllenhal, 1837)	UD	DR	DE	LD	P	MV	SS	E	SL	BL
<i>Trichosirocalus</i> Colonnelli, 1979										
<i>troglodytes</i> (Fabricius, 1787)	UD	DR	DE	LD	P	MV	SS	E		
<i>Zacladus</i> Reitter, 1913										
<i>asperatus</i> (Gyllenhal, 1837)				LD						
<i>exiguus</i> (Olivier, 1807)						MV				
<i>gerani</i> (Paykull, 1800)	UD	DR	DE	LD	P					
Hypurini Schultze, 1902										
<i>Aphytobius</i> H. Wagner, 1937										
<i>*sphaerion</i> (Boheman, 1845)					P					
Mononychini LeConte, 1876										
<i>Mononychus</i> Germar, 1824										
<i>punctumalbum</i> (Herbst, 1784)	UD	DR	DE	LD	P	MV	SS	E	SL	BL
Phytobiini Gistel, 1848										
<i>Neophytobius</i> H. Warner, 1936										
<i>granatus</i> (Gyllenhal, 1835)		DR						E	SL	BL
<i>quadrinodosus</i> (Gyllenhal, 1813)				LD						
<i>Pelenomus</i> C. G. Thomson, 1859										

<i>quadrituberculatus</i> (Fabricius, 1787)		DR	DE	LD				E	SL	
<i>waltoni</i> (Bohemian, 1843)									SL	
<i>Phytobius</i> Schoenherr, 1833										
<i>leucogaster</i> (Marsham, 1802)	UD	DR						E	SL	BL
<i>Rhinoncus</i> Schoenherr, 1825										
<i>bosnicus</i> Schultze, 1900			DE	LD		MV				
<i>bruchoides</i> (Herbst, 1784)		DR			P			E		
* <i>castor</i> (Fabricius, 1792)		DR	DE	LD	P			E	SL	
<i>inconspicetus</i> (Herbst, 1795)			DE	LD	P	MV				
<i>pericarpinus</i> (Linnaeus, 1758)	UD	DR	DE	LD	P	MV	SS	E	SL	BL
<i>perpendicularis</i> (Reich, 1797)	UD		DE	LD		MV	SS			
Scleropterini Schultze, 1902										
<i>Homorosoma</i> J. Frivaldzky, 1894										
<i>validirostre</i> (Gyllenhal, 1837)				LD						
<i>Tapinotus</i> Schoenherr, 1826										
<i>sellatus</i> (Fabricius, 1794)	UD	DR		LD	P	MV				
CONODERINAE Schoenherr, 1833										
<i>Coryssomerus</i> Schoenherr, 1825										
<i>capucinus</i> (Becker, 1817)	UD	DR	DE	LD	P	MV	SS	E	SL	BL
COSSONINAE Schoenherr, 1825										
Cossonini Schoenherr, 1825										
<i>Cossonus</i> subgen. <i>Caenocossonus</i> Voss, 1955										
<i>cylindricus</i> Sahlberg, 1835	UD	DR	DE	LD		MV	SS	E		
<i>Cossonus</i> subgen. <i>Cossonus</i> Clairville, 1798										
<i>parallelepipedus</i> (Herbst, 1795)	UD		DE	LD	P					
<i>Mesites</i> Schoenherr, 1838										
<i>cunipes</i> (Bohemian, 1838)				LD						
Onychlipini Wollaston, 1873										
<i>Hexarthrum</i> Wollaston, 1860										
<i>exiguum</i> (Bohemian, 1838)		DR	DE	LD						
Rhyncolini Gistel, 1848										
<i>Phloeophagus</i> Schoenherr, 1838										
<i>Ignarius</i> Marsham, 1802	UD	DR	DE	LD						
<i>Rhyncolus</i> subgen. <i>Rhyncolus</i> Germar, 1817										
<i>ater ater</i> (Linnaeus, 1758)				LD						
<i>elongatus</i> (Gyllenhal, 1827)			DE							
<i>nefarius</i> Faust, 1885				LD						
CYCLOMINAE Schoenherr, 1826										
<i>Gronops</i> subgen. <i>Asperogronops</i> F. Solari, 1940										
* <i>inaequalis</i> Boheman, 1842		DR								
<i>Entomoderus</i> subgen. <i>Byrsopidius</i> F. Solari,										

1940											
* <i>phrygius</i> (K. & J. Daniel &, 1903)								E	SL		
ENTIMINAE Schoenherr, 1823											
Alophini LeConte, 1874											
<i>Graptus</i> Schoenherr, 1823											
<i>steppensis</i> Davidian, Arzanov & Korotyaev, 2004			DE	LD		MV	SS				
<i>triguttatus triguttatus</i> (Fabricius, 1775)		DR	DE	LD	P						
Brachyderini Schoenherr, 1826											
<i>Brachyderes</i> Schoenherr, 1826											
<i>*incanus</i> (Linnaeus, 1758)		DR	DE	LD							
<i>Pholicodes</i> subgen. <i>Pholicodes</i> Schoenherr, 1826											
<i>inauratus arzanovi</i> Davidian, 1992		DR	DE	LD	P						
<i>Pholicodes</i> subgen. <i>Pseudopholicodes</i> Davidian, 1992											
<i>albidus</i> (Boheman, 1840)					P		SS	E			
<i>Strophomorphus</i> Seidlitz, 1867											
<i>porcellus</i> (Schoenherr, 1832)							SS				
<i>Strophosoma</i> subgen. <i>Pelletierius</i> Alonso-Zarazaga & Lyal, 1999											
<i>albosignatum</i> (Boheman, 1840)		DR	DE								
<i>Strophosoma</i> subgen. <i>Strophosoma</i> Billberg, 1820											
<i>capitatum</i> (DeGeer, 1775)		DR									
Cneorhinini Lacordaire, 1863											
<i>Attactogenus</i> Tournier, 1876											
<i>albinus</i> Boheman, 1833	UD	DR	DE	LD	P						
Cyphicerini Lacordaire, 1863											
<i>Chloebius</i> Schoenherr, 1826											
<i>immeritus</i> Schoenherr, 1826							SS	E	SL	BL	
<i>Ptochus</i> Schoenherr, 1826											
<i>porcellus</i> Boheman, 1834	UD	DR	DE	LD	P	MV	SS	E	SL	BL	
Myorhinini Marseul, 1863											
<i>Apsis</i> Germar, 1820											
<i>albolineata</i> (Fabricius, 1792)	UD	DR	DE	LD	P	MV	SS	E	SL	BL	
Nastini Reitter, 1913											
<i>Nastus</i> Schoenherr, 1842											
<i>goryi sareptanus</i> Faust, 1883				LD	P	MV	SS				
Naupactini Gistel, 1848											
<i>Mesagroicus</i> Schoenherr, 1840											
<i>*poriventris</i> Reitter, 1903				LD			SS	E	SL	BL	

Omiini Shuckard, 1840											
<i>Humeromima</i> Podlussány, 1998											
<i>nitida</i> (Boheman, 1842)		DR	DE						E		
<i>Omias</i> Germar, 1817											
<i>murinus</i> (Boheman, 1842)		DR	DE	LD	P	MV	SS				
<i>puberulus</i> (Boheman, 1834)	UD	DR	DE	LD	P	MV	SS	E	SL		
<i>verruca</i> (Boheman, 1834)	UD	DR	DE	LD	P	MV	SS	E	SL	BL	
<i>Urometopus</i> Formánek, 1904											
<i>nemorum</i> Arnol'di, 1965	UD	DR	DE	LD							
<i>strigifrons</i> (Gyllenhal, 1834)		DR	DE	LD							
Otiorhynchini Schoenherr, 1826											
<i>Otiorhynchus</i> subgen. <i>Amosilnus</i> Reitter, 1912											
* <i>simulans</i> Stierlin, 1877				LD							
<i>Otiorhynchus</i> subgen. <i>Choilisanus</i> Reitter, 1912											
<i>balcanicus</i> Stierlin, 1861				LD							
<i>brunneus</i> Steven, 1909	UD	DR	DE	LD	P	MV	SS	E	SL		
<i>pilosus pilosus</i> Gyllenhal, 1834	UD	DR	DE								
<i>raucus</i> (Fabricius, 1777)	UD	DR	DE	LD							
<i>velutinus</i> Germar, 1824	UD	DR	DE	LD	P	MV	SS	E	SL	BL	
<i>Otiorhynchus</i> subgen. <i>Cryphiphorus</i> Magnano, 1998											
<i>ligustici</i> (Linnaeus, 1758)	UD	DR	DE	LD	P	MV	SS	E	SL		
<i>Otiorhynchus</i> subgen. <i>Duphuna</i> Reitter, 1914											
<i>aurosparsus</i> Germar, 1824		DR									
<i>Otiorhynchus</i> subgen. <i>Melasemnus</i> Reitter, 1912											
<i>smreczynskii</i> Cmoluch, 1968		DR	DE	LD							
<i>Otiorhynchus</i> subgen. <i>Nubidamus</i> Reitter, 1912											
<i>juvenilis</i> Schoenherr, 1832								E	SL	BL	
<i>Otiorhynchus</i> subgen. <i>Otiolehus</i> Reitter, 1914											
<i>tristis</i> (Scopoli, 1763)	UD	DR	DE	LD							
<i>Otiorhynchus</i> subgen. <i>Pendragon</i> Gozis, 1885											
<i>ovatus ovatus</i> (Linnaeus, 1758)	UD	DR	DE	LD	P	MV	SS				
<i>Otiorhynchus</i> subgen. <i>Podoropelmus</i> Reitter, 1912											
<i>albidus</i> Stierlin, 1861				LD							
<i>fullo</i> Schrank, 1781	UD	DR	DE	LD							
<i>scopularis</i> Hochhuth, 1847	UD	DR	DE	LD			SS				
<i>Otiorhynchus</i> subgen. <i>Pontotiorhynchus</i> Yunakov, 2003											
<i>asphaltinus</i> Germar, 1824		DR	DE	LD							
<i>Otiorhynchus</i> subgen. <i>Pseudocryphiphorus</i> Magnano, 1998											
<i>chrysostictus</i> Gyllenhal, 1834	UD	DR	DE	LD	P	MV	SS	E			
<i>Otiorhynchus</i> subgen. <i>Zelotomelus</i> Reitter,											

1912											
<i>erinaceus</i> Stierlin, 1876		DR									
<i>Stomodes</i> Schoenherr, 1826											
<i>tolutarius</i> Schoenherr, 1826				LD		MV	SS				
Phyllibiini Schoenherr, 1826											
<i>Phyllobius</i> subgen. <i>Alsus</i> Motschulsky, 1845											
<i>brevis</i> Gyllenhal, 1834	UD	DR	DE	LD	P	MV	SS	E	SL	BL	
<i>Phyllobius</i> subgen. <i>Dieletus</i> Reitter, 1916											
<i>argentatus argentatus</i> (Linnaeus, 1758)	UD	DR	DE								
<i>Phyllobius</i> subgen. <i>Metaphyllobius</i> Smirnov, 1913											
<i>glaucus</i> (Scopoli, 1763)	UD	DR									
<i>pomaceus</i> Gyllenhal, 1834		DR									
<i>Phyllobius</i> subgen. <i>Nanoschetus</i> Reitter, 1916											
<i>cylindricollis</i> Gyllenhal, 1834			DE	LD		MV	SS				
<i>Phyllobius</i> subgen. <i>Nemoicus</i> Dillwyn, 1829											
<i>oblongus</i> (Linnaeus, 1758)	UD	DR	DE	LD	P	MV	SS				
<i>Phyllobius</i> subgen. <i>Phyllobius</i> Germar, 1824											
<i>betulinus betulinus</i> (Bechstein & Scharfenberg, 1805)		DR	DE	LD							
<i>pyri</i> (Linnaeus, 1758)	UD	DR	DE	LD	P	MV					
<i>thalassinus</i> Gyllenhal, 1834	UD	DR	DE	LD							
<i>Phyllobius</i> subgen. <i>Pterygorrhynchus</i> Pesarini, 1969											
<i>contemptus</i> Schoenherr, 1832	UD	DR	DE	LD							
<i>maculicornis</i> Germar, 1824								E			
<i>Phyllobius</i> subgen. <i>Subphyllobius</i> Schilsky, 1911											
<i>virideaeoris virideaeoris</i> (Laicharting, 1781)										BL	
Polydrusini Schoenherr, 1823											
<i>Liophloeus</i> Germar, 1817											
* <i>tessulatus</i> (O. F. Müller, 1776)		DR	DE								
<i>Polydrusus</i> subgen. <i>Eudipnus</i> C. G. Thomson, 1859											
<i>mollis</i> (Strøm, 1768)	UD	DR	DE								
<i>Polydrusus</i> subgen. <i>Eurodrusus</i> Korotyaev & Meleshko, 1997											
<i>confluens</i> Stephens, 1831		DR		LD							
<i>Polydrusus</i> subgen. <i>Eustolus</i> C.G. Thomson, 1859											
<i>impressifrons</i> Gyllenhal, 1834	UD	DR	DE	LD						BL	
<i>pterygomalis</i> Boheman, 1840		DR	DE		P	MV	SS				
<i>flavipes</i> (DeGeer, 1775)	UD	DR	DE								
<i>Polydrusus</i> subgen. <i>Leucodrusus</i> Stierlin, 1884											
<i>mariae</i> (Faust, 1882)				LD							
<i>Polydrusus</i> subgen. <i>Poecilodrosus</i> Korotyaev											

& Meleshko, 1997										
* <i>viridicinctus</i> Gyllenhal, 1834	UD	DR	DE							
<i>Polydrusus</i> subgen. <i>Polydrusus</i> Germar, 1817										
<i>ferrugineus</i> Boheman, 1840				LD						
<i>Polydrusus</i> subgen. <i>Scythodrusus</i> Korotyaev & Meleshko, 1997										
<i>inustus</i> Germar, 1824	UD	DR	DE	LD	P	MV	SS	E	SL	
Psallidiini Lacordaire, 1863										
<i>Psallidium</i> Herbst, 1795										
<i>maxillosum maxillosum</i> (Fabricius, 1792)	UD	DR	DE	LD	P	MV	SS	E	SL	
Sciaphilini Sharp, 1891										
<i>Archeophloeus</i> Iablokoff-Khnzorian, 1959										
<i>inermis</i> (Boheman, 1842)			DE			MV	SS			
<i>Brachysomus</i> Schoenherr, 1823										
<i>echinatus</i> (Bonsdorff, 1785)	UD			LD						
<i>Eusomatus</i> Krynicki, 1834										
<i>taeniatus</i> Krynicki, 1834		DR	DE	LD	P					
<i>virens</i> (Boheman, 1883)	UD						SS	E	SL	
<i>Eusomostrophus</i> Tournier, 1878										
<i>acuminatus</i> (Boheman, 1840)	UD	DR	DE	LD	P	MV	SS	E	SL	BL
<i>Eusomus</i> Germar, 1824										
<i>ovulum</i> Germar, 1824	UD	DR	DE	LD	P	MV	SS	E	SL	BL
<i>Exomias</i> Bedel, 1883										
<i>lebedevi</i> (Roubal, 1926)			DE							
<i>pellucidus pellucidus</i> (Boheman, 1834)				LD						
<i>Foucartia</i> Jacquelin du Val, 1854										
<i>liturata</i> Stierlin, 1884			DE							
<i>squamulata</i> (Herbst, 1795)	UD	DR	DE	LD	P	MV	SS			
<i>Sauromates</i> Korotyaev, 1992										
<i>arnoldii</i> Korotyaev, 1992		DR	DE	LD	P					
<i>Sciaphilus</i> Schoenherr, 1823										
<i>asperatus</i> (Bonsdorff, 1785)	UD									
<i>Sciaphobus</i> K. Daniel, 1904										
<i>ningnidus</i> (Germar, 1824)		DR								
<i>squalidus squalidus</i> (Gyllenhal, 1834)	UD	DR	DE	LD	P					
Sitonini Gistel, 1848										
<i>Schelopius</i> Desbrochers des Loges, 1872										
<i>planifrons planifrons</i> Fåhraeus, 1840								SL	BL	
<i>Sitona</i> Germar, 1817										
<i>callosus</i> Gyllenhal, 1834	UD	DR	DE	LD	P	MV	SS	E	SL	BL
<i>cylindricollis cylindricollis</i> (Fåhraeus, 1840)	UD	DR	DE	LD	P	MV	SS	E	SL	BL
<i>hispidulus</i> (Fabricius, 1777)		DR	DE	LD	P					
<i>humeralis</i> Stephens, 1831	UD	DR	DE	LD	P	MV	SS	E	SL	BL

	UD	DR	DE	LD	P	MV	SS	E	SL	BL
<i>inops</i> Schoenherr, 1832				LD						
<i>languidus</i> Gyllenhal, 1834				LD						
<i>lateralis</i> Gyllenhal, 1834	UD		DE	LD	P					
<i>lineatus</i> (Linnaeus, 1758)	UD	DR	DE	LD	P	MV	SS	E		
<i>longulus</i> Gyllenhal, 1834			DE	LD	P					
<i>macularius macularius</i> (Marsham, 1802)	UD	DR	DE	LD	P	MV	SS	E		
<i>obsoletus obsoletus</i> (Gmelin, 1790)		DR	DE	LD						
<i>puncticollis</i> Stephens, 1831	UD	DR	DE	LD	P	MV	SS	E	SL	BL
<i>striatellus</i> Gyllenhal, 1834			DE	LD			SS			
<i>sulcifrons</i> (Thunberg, 1798)		DR	DE	LD						
<i>suturalis suturalis</i> Stephens, 1831	UD	DR	DE	LD	P	MV	SS	E		
* <i>waterhousei</i> Walton, 1846				LD						
Tanymecini Lacordaire, 1863										
<i>Chlorophanus</i> C. R. Sahlberg, 1823										
<i>flavescens</i> (Fabricius, 1787)	UD		DE	LD						
<i>micans</i> Krynicki, 1832			DE	LD			SS	E	SL	BL
<i>vittatus</i> Schoenherr, 1832									SL	
<i>Cycloderes</i> C. R. Sahlberg, 1823										
<i>pilosulus</i> (Herbst, 1795)	UD	DR	DE	LD	P	MV	SS	E	SL	BL
<i>Megamecus</i> subgen. <i>Acercomecus</i> Reitter, 1903										
<i>argentatus</i> Gyllenhal, 1834								E	SL	BL
<i>Megamecus</i> subgen. <i>Megamecus</i> Reitter, 1903										
<i>urbanus</i> (Gyllenhal, 1834)								E	SL	BL
<i>variegatus</i> (Gebler, 1830)								E	SL	BL
<i>Phacephorus</i> Schoenherr, 1840										
<i>argyrostomus</i> (Gyllenhal, 1840)									SL	BL
<i>nebulosus</i> (Fähraeus, 1840)							SS	E	SL	BL
<i>Tanymecus</i> subgen. <i>Episomecus</i> Reitter, 1903										
<i>dilaticollis</i> Gyllenhal, 1834			DE	LD	P	MV	SS	E		
<i>Tanymecus</i> subgen. <i>Tanymecus</i> Germar, 1817										
<i>palliatus</i> (Fabricius, 1787)	UD	DR	DE	LD	P	MV	SS	E	SL	
<i>Xylinophorus</i> Faust, 1885										
<i>scobinatus</i> (Ménétriés, 1858)							SS	E		
Trachyphloeini Gistel, 1848										
<i>Cathormiocerus</i> Schoenherr, 1842										
<i>aristatus</i> (Gyllenhal, 1827)				LD		MV				
<i>Trachyphloeus</i> Faust, 1887										
<i>alternans</i> Gyllenhal, 1834		DR	DE	LD	P	MV	SS	E		
<i>scabriculus</i> (Linnaeus, 1771)				LD						
<i>spinimanus</i> Germar, 1824				LD		MV				
HYPERINAE Marseul, 1863										
<i>Adonus</i> Zaslavskij, 1999										

<i>asiaticus</i> (Ménétriés, 1849)						MV	SS	E	SL	BL
<i>Coniatus</i> Germar, 1817										
<i>splendidulus</i> (Fabricius, 1781)			LD		MV	SS	E	SL	BL	
<i>Hypera</i> subgen. <i>Dapalinus</i> Capiomont, 1868										
<i>fornicata</i> (Peneki, 1928)		DE	LD	P	MV	SS				
<i>meles</i> (Fabricius, 1792)		DE	LD	P						
<i>Hypera</i> subgen. <i>Eririnomorphus</i> Capiomont, 1868										
* <i>arundinis</i> (Paykull, 1792)		DR	DE		P	MV				
<i>connaculata</i> (Herbst, 1795)		DR	DE	LD						
<i>interruptovittatus</i> (Desbrochers des Loges, 1875)						MV	SS			
* <i>rogenhoferi</i> (Ferrari, 1866)						MV				
<i>ruminis</i> (Linnaeus, 1758)	UD	DR	DE	LD	P	MV	SS	E		
<i>Hypera</i> subgen. <i>Hypera</i> Germar, 1817										
<i>denominanda</i> (Capiomont, 1868)			DE	LD	P	MV				
<i>farinosa</i> (Boheman, 1840)								E	SL	BL
<i>melancholica</i> (Fabricius, 1792)			DE							
<i>miles</i> (Paykull, 1792)		DR	DE	LD	P			E	SL	
<i>nigrirostris</i> (Fabricius, 1775)				LD	P					
<i>plantaginis</i> (DeGeer, 1775)		DR	DE	LD	P	MV	SS			
<i>postica</i> (Gyllenhal, 1813)	UD	DR	DE	LD	P	MV	SS	E	SL	BL
<i>transsylvanica</i> (Petri, 1901)			DE	LD	P	MV		E		
<i>viciae</i> (Gyllenhal, 1813)		DR	DE	LD	P	MV	SS			
<i>Hypera</i> subgen. <i>Kippenbergia</i> Alonso-Zarazaga & Lyal, 2005										
<i>arator</i> (Linnaeus, 1758)		DR		LD						BL
<i>Limobius</i> Schoenherr, 1843										
<i>borealis borealis</i> (Paykull, 1792)	UD		DE	LD	P					
<i>Metadonus</i> Capiomont, 1868										
<i>anceps</i> (Boheman, 1842)	UD	DR	DE	LD	P	MV	SS	E	SL	BL
<i>curtus</i> (Boheman, 1842)								E	SL	BL
<i>distinguendus</i> (Boheman, 1842)		DR	DE	LD		MV	SS	E	SL	
<i>Pachyperra</i> Capiomont, 1868										
<i>spissa</i> (Boheman, 1842)				LD	P	MV	SS	E	SL	
LIXINAE Schoenherr, 1823										
Cleonini Schoenherr, 1826										
<i>Adosomus</i> Faust, 1904										
<i>roridus</i> (Pallas, 1781)		DR		LD						
<i>Asproparthenis</i> Gozis, 1886										
<i>carinata</i> (Zoubkoff, 1829)								E	SL	BL
<i>carinicollis</i> (Gyllenhal, 1834)				LD		MV	SS	E	SL	BL
<i>foveocollis</i> (Gebler 1834)										BL
<i>punctiventris</i> (Germar, 1824)	UD	DR	DE	LD	P	MV	SS	E	SL	BL
<i>vexata</i> (Gyllenhal, 1834)						MV	SS	E	SL	BL
<i>Bothynoderes</i> Schoenherr, 1823										
<i>affinis</i> (Schrank, 1781)	UD	DR	DE	LD	P	MV	SS	E		

	UD	DR	DE	LD	P	MV	SS	E	SL	BL
<i>declivis</i> (Olivier, 1807)										
<i>Chromonotus</i> subgen. <i>Chevrolatius</i> Arzanov, 2006										
<i>bipunctatus</i> (Zoubkoff, 1829)									SL	BL
<i>hirsutulus</i> (Faust, 1883)								E		
<i>pilosellus</i> (Fåhraeus, 1842)										BL
<i>Chromonotus</i> subgen. <i>Chromonotus</i> Motschulsky, 1860										
<i>pictus</i> (Pallas, 1771)								E	SL	BL
<i>vittatus</i> (Zoubkoff, 1829)								E	SL	BL
<i>Cleonis</i> Dejean, 1821										
<i>pigra</i> (Scopoli, 1763)	UD	DR	DE	LD	P	MV	SS	E	SL	BL
<i>Coniocleonus</i> subgen. <i>Augustecleonus</i> Arzanov, 2006										
* <i>hollbergi</i> (Fåhraeus, 1842)				LD	P		SS			
<i>Conorhynchus</i> Motschulsky, 1860										
<i>lacerta</i> Chevrolat, 1873						MV	SS	E	SL	BL
<i>nigrivittis</i> (Pallas, 1781)								E	SL	BL
<i>pulverulentus</i> (Zoubkoff, 1829)								E	SL	BL
<i>Cyphocleonus</i> Motschulsky, 1860										
<i>achates</i> (Fåhraeus, 1842)	UD		DE	LD				E		
<i>adumbratus</i> (Gebler, 1833)				LD						
<i>cenchrus</i> (Pallas, 1781)								E		
<i>dealbatus</i> (Gmelin, 1790)	UD	DR	DE	LD	P	MV	SS	E	SL	BL
<i>trisulcatus</i> (Herbst, 1795)	UD	DR	DE	LD	P					
<i>Eumecops</i> Hochhuth, 1851										
<i>kittayi</i> (Hochhuth, 1851)				LD		MV				
<i>Leucomigus</i> Motschulsky, 1860										
<i>candidatus candidatus</i> (Pallas, 1771)	UD	DR	DE	LD	P	MV	SS	E	SL	BL
<i>Leucophyes</i> Marshall, 1946										
<i>pedestris</i> (Poda, 1761)	UD	DR	DE	LD		MV				
<i>Maximus</i> Alonso-Zarazaga & Lyal, 2009										
<i>strabus</i> (Gyllenhal, 1834)			DE	LD	P	MV	SS	E	SL	BL
<i>Mecaspis</i> Schoenherr, 1823										
<i>alternans</i> (Herbst, 1795)		DR	DE	LD						
<i>Pachycerus</i> Schoenherr, 1823										
<i>segnis</i> (Germar, 1824)	UD	DR	DE	LD	P	MV	SS	E	SL	BL
<i>Pleurocleonus</i> Motschulsky, 1860										
<i>quadrivittatus</i> (Zoubkoff, 1829)			DE	LD		MV	SS	E		
* <i>sollicitus</i> (Gyllenhal, 1834)		DR		LD				E	SL	
<i>Pseudocleonus</i> subgen. <i>Neopseudocleonus</i> Arzanov, 2005										
* <i>grammicus</i> (Panzer, 1789)				LD						
<i>marginicollis</i> (Fåhraeus, 1842)			DE							
<i>Pseudocleonus</i> subgen. <i>Pseudocleonus</i> Chevrolat, 1872										
<i>cinereus</i> (Schrank, 1781)		DR	DE	LD	P	MV	SS			
<i>Rhabdorrhynchus</i> Motschulsky, 1860										
* <i>echii</i> (Brahm, 1780)		DR		LD			SS			

<i>Scaphomorphus</i> Motschulsky, 1860											
<i>vibex</i> (Pallas, 1781)											BL
<i>Stephanocleonus</i> subgen. <i>Sanzia</i> Alonso-Zarazaga & Lyal, 1999											
<i>microgrammus</i> (Gyllenhal, 1834)	DR	DE									
<i>tetragrammus</i> (Pallas, 1781)		DE	LD		MV	SS					
<i>Temnorhinus</i> subgen. <i>Massimocleonus</i> Arzanov, 2006											
<i>elongatus</i> (Gebler, 1844)								E	SL		
<i>Temnorhinus</i> subgen. <i>Temnorhinus</i> Chevrolat, 1872											
<i>hololeucus</i> (Pallas, 1781)								E	SL	BL	
<i>vereundus</i> (Faust, 1883)									SL		
<i>Xanthochelus</i> Chevrolat, 1872											
<i>*nomas</i> (Pallas, 1771)					MV						
Lixini Schoenherr, 1823											
<i>Hypolixus</i> Desbrochers des Loges, 1883											
<i>astrachanicus</i> (Faust, 1883)										BL	
<i>Lachnaeus</i> Schoenherr, 1826											
<i>crinitus</i> (Boheman, 1826)	UD	DR	DE	LD	P	MV	SS	E	SL	BL	
<i>Larinus</i> subgen. <i>Larinus</i> Dejean, 1821											
<i>cynarae</i> (Fabricius, 1787)			DE	LD							
<i>*idoneus</i> Gyllenhal, 1835	UD	DR	DE								
<i>inaequalicollis</i> Capiomont, 1874	UD	DR	DE	LD	P	MV					
<i>latus</i> (Herbst, 1783)				LD							
<i>onopordi</i> (Fabricius, 1787)	UD	DR	DE								
<i>pollinis</i> (Laicharting, 1781)				LD							
<i>*sibiricus</i> Gyllenhal, 1835		DR	DE	LD							
<i>vulpes</i> (Olivier, 1807)				LD							
<i>Larinus</i> subgen. <i>Larinomesius</i> Reitter, 1924											
<i>canescens</i> Gyllenhal, 1835		DR	DE	LD							
<i>curtus</i> Hochhuth, 1851			DE								
<i>minutus</i> Gyllenhal, 1836	UD	DR	DE	LD	P	MV	SS	E	SL		
<i>nubeculosus</i> Gyllenhal, 1835				LD							
<i>obtusus</i> Gyllenhal, 1835	UD	DR	DE	LD	P	MV					
<i>ruber</i> Motschulsky, 1845		DR									
<i>serratulae</i> Becker, 1864				LD						BL	
<i>Larinus</i> subgen. <i>Phyllonomeus</i> Gistel, 1856											
<i>adspersus</i> Hochhuth, 1847				LD							
<i>centaurii</i> (Olivier, 1807)	UD	DR	DE	LD	P	MV	SS	E	SL		
<i>iaceae</i> (Fabricius, 1775)	UD	DR	DE	LD	P	MV	SS	E	SL	BL	
<i>rusticanus</i> Gyllenhal, 1835	UD	DR	DE	LD	P	MV					
<i>sturnus</i> (Gyllenhal, 1783)			DE	LD		MV	SS				
<i>turbinatus</i> Gyllenhal, 1835	UD	DR	DE	LD	P	MV	SS				
<i>Lixus</i> subgen. <i>Broconius</i> Desbrochers des Loges, 1904											
<i>rubicundus</i> Zoubkoff, 1833		DR	DE	LD	P	MV	SS	E	SL	BL	
<i>Lixus</i> subgen. <i>Callistolixus</i> Reitter, 1916											

<i>cylindrus</i> (Fabricius, 1781)			DE							
<i>Lixus</i> subgen. <i>Compsolixus</i> Reitter, 1916										
<i>albomarginatus</i> Boheman, 1842	UD	DR	DE	LD	P	MV	SS	E	SL	BL
<i>linnei</i> Faust, 1888								E		BL
<i>Lixus</i> subgen. <i>Dilixellus</i> Reitter, 1916										
<i>bardanae</i> (Fabricius, 1787)	UD	DR	DE	LD	P	MV	SS	E	SL	BL
<i>fasciculatus</i> Boheman, 1835		DR	DE	LD	P					
<i>pulverulentus</i> (Scopoli, 1763)		DR	DE	LD				E		BL
<i>punctiventris</i> Boheman, 1835			DE	LD	P		SS	E		
<i>Lixus</i> subgen. <i>Epimeces</i> Billberg, 1820										
<i>cardui</i> Olivier, 1807	UD	DR	DE	LD	P	MV	SS	E	SL	BL
<i>filiformis</i> (Fabricius, 1781)	UD	DR	DE	LD	P	MV	SS	E	SL	BL
<i>scolopax</i> Boheman, 1835										BL
<i>Lixus</i> subgen. <i>Eulixus</i> Reitter, 1916										
<i>iridis</i> Olivier, 1807	UD	DR	DE	LD	P	MV	SS			
<i>kraatzi</i> Capiomont, 1875				LD	P	MV	SS	E	SL	BL
<i>myagri</i> Olivier, 1807			DE	LD			SS	E		BL
<i>punctirostris</i> Boheman, 1842		DR		LD			SS			
<i>Lixus</i> subgen. <i>Lixus</i> Fabricius, 1801										
<i>paraplecticus</i> (Linnaeus, 1758)				LD						
<i>Lixus</i> subgen. <i>Ortholixus</i> Reitter, 1916										
<i>angustus</i> (Herbst, 1795)	UD		DE	LD		MV				
<i>cinerascens</i> Schoenherr, 1832			DE	LD		MV	SS			
<i>Lixus</i> subgen. <i>Phillixus</i> Petri, 1904										
<i>brevipes</i> C. N. F. Brisout de Barnevile, 1866				LD						
<i>incanescens</i> Boheman, 1835			DE	LD	P	MV	SS	E	SL	BL
<i>subtilis</i> Boheman, 1835	UD	DR	DE	LD	P	MV	SS	E	SL	BL
<i>subulatus</i> Faust, 1891						MV	SS	E	SL	BL
<i>Rhinocyllus</i> Germar, 1817										
<i>conicus</i> (Frölich, 1792)			DE	LD	P	MV	SS	E	SL	BL
MESOPTILIINAE Lacordaire, 1863										
<i>Magdalalis</i> subgen. <i>Edo</i> Germar, 1819										
<i>ruficornis</i> (Linnaeus, 1758)	UD	DR	DE	LD						
<i>Magdalalis</i> subgen. <i>Laemosaccidius</i> Smreczyński, 1972										
<i>exarata</i> C. N. F. Brisout de Barnevile, 1862	UD									
<i>Magdalalis</i> subgen. <i>Magdalalis</i> Germar, 1817										
<i>frontalis</i> (Gyllenhal, 1827)			DE							
* <i>phlegmatica</i> (Herbst, 1797)		DR	DE							
<i>violacea</i> (Linnaeus, 1758)		DR								
<i>Magdalalis</i> subgen. <i>Odontomagdalalis</i> Barrios, 1984										
<i>armigera</i> (Geoffroy, 1785)	UD	DR	DE	LD						
<i>Magdalalis</i> subgen. <i>Panus</i> Schoenherr, 1823										
<i>barbicornis</i> (Latreille, 1804)				LD						
<i>Magdalalis</i> subgen. <i>Porrothus</i> Dejean, 1821										

<i>cerasi</i> (Linnaeus, 1758)		DR							
MOLYTINAE Schoenherr, 1823									
Cryptorhynchini Schoenherr, 1825									
<i>Cryptorhynchus</i> Illiger, 1807									
* <i>lopathi</i> (Linnaeus, 1758)			LD						
Gasterocercini Zherikhin, 1991									
<i>Gastrocercus</i> Laporte & Brullé, 1828									
<i>depressirostris</i> (Fabricius, 1792)		DE	LD						
Lepyrini W. Kirby, 1837									
<i>Lepyrus</i> Germar, 1817									
<i>capucinus</i> (Schaller, 1783)	UD		DE	LD	P				
<i>palustris flavidulus</i> Zumpt, 1936		DR	DE	LD	P				
Hylobiini W. Kirbi, 1837									
<i>Hylobius</i> subgen. <i>Callirus</i> Dejean, 1821									
<i>abietis</i> (Linnaeus, 1758)		DR	DE						
<i>transversovittatus</i> (Goeze, 1777)	UD	DR	DE	LD	P	MV			
Molytini Schoenherr, 1983									
<i>Liparus</i> subgen. <i>Liparus</i> Olivier, 1807									
<i>coronatus</i> (Goeze, 1777)	UD	DR	DE	LD					
<i>dirus</i> (Herbst, 1795)		DR		LD					
<i>Liparus</i> subgen. <i>Trysibius</i> Schoenherr, 1842									
<i>tenebrioides</i> (Pallas, 1781)		HD	LD	P	M V	SS			
<i>Minyops</i> Schoenherr, 1823									
<i>minutus redis</i> (Ménétriés, 1849)			DE	LD		MV	SS	E	
Pissodini Gistel, 1848									
<i>Pissodes</i> Germar, 1817									
<i>castaneus</i> (DeGeer, 1775)		DR	DE						
* <i>pini pini</i> (Linnaeus, 1758)				LD					
OROBITIDINAE C. G. Thomson, 1859									
<i>Orobitis</i> Germar, 1817									
* <i>cyaneus</i> (Linnaeus, 1758)		DR		LD					

Acknowledgements

I owe a depth of gratitude to my late teacher and tutor in entomology Margarita E. Ter-Minasian (St. Petersburg, Russia). Boris A. Korotyaev and Genrik E. Davidian (both St. Petersburg, Russia) critically checked identification of some specimens. Pavel P. Ivliev, Denis G. Kasatkin, Eduard A. Khachikov, Maxim V. Nabozhenko, and Igor V. Shokhin (all Rostov-on-Don, Russia) for over two decades collected and supplied weevil specimens. Vasily V. Grebennikov (Ottawa, Canada) critically read an early draft of this paper.

References

- Alonso-Zarazaga M. A. & Lyal C. H. C. 1999.** *A World Catalogue of Families and Genera of Curculionoidea (Insecta: Coleoptera) (Excepting Scolytidae and Platypodidae).* Entomopraxis, Barcelona, 315 pp.
- Alonso-Zarazaga M. A. & Lyal C. H. C. 2002.** Addenda and corrigenda to “A world catalogue of families and genera of Curculionoidea (Insecta: Coleoptera)”. *Zootaxa* 63: 1–37.
- Arzanov Yu. G. 1988.** K faune zhukov-dolgonosikov podsem. Cleoninae (Coleoptera, Curculionidae) severnogo Kavkaza [Contribution to the fauna of Cleoninae (Coleoptera, Curculionidae) of the northern Caucasus]. *Entomologicheskoe Obozrenie* 67(3): 514–522.
- Arzanov Yu. G. 1990.** Obzor fauny zhukov-dolgonosikov (Coleoptera, Curculionidae) Rostovskoi oblasti I Kalmikykskoi ASSR [Review of the fauna of weevils (Coleoptera, Curculionidae) of Rostov province and Kalymik ASSR]. *Entomologicheskoe Obozrenie* 69(2): 313–331.
- Dieckmann L. 1980.** Revision der *Lixus ascanii*-Grupp (Coleoptera, Curculionidae). *Reichenbachia* 18: 203–212.
- Dobrovolsky B. V. 1951.** *Vrednie zhuki* [Noxious beetles]. Rostizdat, Rostov-on-Don, 456 pp.
- Kiseritzky V. A. 1912.** K faune zhukov oblasti voyska Donskogo [On the fauna of beetles of the Don Cossack Host]. *Russkoe Entomologicheskoe Obozrenie* 2: 82–94.
- Korotyeav B. A. & Fridman A. L. L. 2011.** A new species of the weevils genus *Melanobaris* Alonso-Zarazaga & Lyal, 1999 (Coleoptera: Curculionidae: Baridinae) from Mt. Hermon in Israel and commentaries on the composition of the genera *Melanobaris* and *Aulacobaris* Desbrochers, 1892. *Caucasian Entomological Bulletin* 7(2): 169–172.
- Löbl I. & Smetana A. (eds.) 2011.** *Catalogue of Palaearctic Coleoptera. Volume 7. Curculionoidea I.* Apollo Books, Stenstrup, 373 pp.
- Löbl I. & Smetana A. (eds.) 2013.** *Catalogue of Palaearctic Coleoptera. Volume 8. Curculionoidea II.* Brill, Leiden, 700 pp.
- Lukjanovitsh F. K. 1926.** K faune dolgonosikov (Curculionidae) Donskogo okruga [On the fauna of weevils (Curculionidae) Don District]. *Izvestie Stavropolskogo Entomologicheskogo Obshestva* 2: 20–22.
- Lyal C. H. C. 2014.** 3.7.7 Molytinae Schoenherr, 1823. In: Leschen, R.A.B. & Beutel, R.G. (Eds.), *Handbook of Zoology, Arthropoda: Insecta: Coleoptera. Volume 3: Morphology and Systematics (Phytophaga)*. Walter de Gruyter, Berlin, pp. 529–570.
- Osella G. & Bellò C. 2010.** Revisione di *Minyops* Schoenherr e *Paraminyops* nov. gen. (Coleoptera, Curculionidae, Molytinae). *Memorie del Museo Civico di Storia Naturale di Verona, 2. serie, Sezione Scienze della Vita* 19: 3–133.

Correspondence: Yuri G. Arzanov, e-mail: arz99@mail.ru

Received: 27.05.2015 Accepted: 11.07.2015 Published: 01.08.2015

Cite paper: Arzanov Yu. G. 2015. A revised checklist species of the Curculionoidea (Coleoptera, excluding Scolytinae) of Rostov Oblast and Kalmykia, the southern part of European Russia. *Journal of Insect Biodiversity* 3(12): 1–32.

<http://dx.doi.org/10.12976/jib/2015.3.12>

<http://www.insectbiodiversity.org>