



A New Species of *Pleurostachys* (Cyperaceae) from Atlantic coastal Brazil

W. WAYT THOMAS¹, MARCUS V. ALVES² & RAFAEL TREVISAN³

¹The New York Botanical Garden, Bronx, NY 10458-5126, USA. Email: wthomas@nybg.org

²Departamento de Botânica, CCB, Universidade Federal de Pernambuco, Av. Moraes Rego s/n, 50940-000 Recife, PE, Brazil.
Email: alves.marcus@gmail.com

³Departamento de Botânica, CCB, Universidade Federal de Santa Catarina, Av. R. Eng. Agrônomo Andrey Cristian Ferreira, 88040-970 Florianópolis, SC, Brazil. Email: rftrevisan@yahoo.com.br

Abstract

Pleurostachys arcuata is described as a new species endemic to the Atlantic forest of Brazil. It is illustrated and is compared to other species of *Pleurostachys*. It has the large achenes of *P. macrantha* but has small, axillary synflorescences similar to those of *P. gaudichaudii*.

Resumo

Pleurostachys arcuata é descrita como uma espécie nova da Mata Atlântica do Brasil. Está ilustrada e comparada com outras espécies de *Pleurostachys*. Tem os aquêniros grandes de *P. macrantha* mas as sinflorescências pequenas e axilares semelhantes às de *P. gaudichaudii*.

Key words: *Pleurostachys*, Cyperaceae, Brazil, Atlantic forest, taxonomy

Introduction

Pleurostachys Brongn. is a small genus found exclusively in neotropical forests. Kükenthal's monograph of 1952 treated 30 species. Thomas and Alves (2008) recognized 23 species, 21 of which are endemic to Brazil's Atlantic coastal forest. Modern collections have greatly increased the number of specimens available for study, revealing several undescribed species, one of which is described here.

Most authors have maintained *Pleurostachys* as a separate genus closely allied to *Rhynchospora* Vahl (Boeckler, 1875; Clarke, 1908; Pfeiffer, 1925; Koyama, 1961; Kükenthal, 1952; Eiten, 1976; Bruhl, 1995; Goetghebeur, 1986, 1998; Soros & Bruhl, 2000). Recent molecular studies (Thomas *et al.* 2009a) confirm this relationship and raise the question as to whether or not *Rhynchospora* is paraphyletic with respect to *Pleurostachys*. Because the phylogenetic relationship of *Pleurostachys* and *Rhynchospora* is still unclear, and because they are morphologically distinct, we are describing this new species as a member of the genus *Pleurostachys*.

Features that unite *Rhynchospora* and *Pleurostachys* in the tribe Rhynchosporae include simple spikelets, bisexual flowers, lenticular achenes subtended by perianth bristles, and a persistent style base. Characters that distinguish *Pleurostachys* from *Rhynchospora* include an inflorescence comprising a series of axillary synflorescences, distichously (or subdistichously) arranged spikelet scales, distally plumose perianth bristles, and decurrent style bases (these are discussed at length in Thomas & Alves, 2008).

Taxonomy

Pleurostachys arcuata W. W. Thomas, M. Alves & R. Trevis., spec. nov. (Fig. 1)

Pleurostachys arcuata is unique in having basal leaves which are long and wide, small synflorescences of clustered spikelets, and large achenes. It has the large achenes of *P. macrantha* but has the small, axillary synflorescences of *P. gaudichaudii*.

Type:—Brazil. Bahia: Mun. Santa Luzia, Serra da Onça, 10.8 km NE of Santa Luzia (30 km SW of Una) on Una-Santa Luzia road, then 4.2 km N on road to Serra da Onça. 39°30'S, 15°20'W. Southern Bahian Wet Forest. Reddish clay and many rocks. Rare in forest, leaves shiny dark green, inflorescence arching to scandent. 21 Nov 1996. *W. W. Thomas, A. Carvalho, A. Amorim, S. Sant'Ana & J. L. Paixão* 11367 (holotype CEPEC, isotypes K, NY).

Description

Perennial, cespitose or with short woody rhizomes. Culms arching, (30.5–) 58–212 cm long, (1–) 1.8–3 mm wide at lowest synflorescence, tapering to 0.8–1 mm below distal-most synflorescence, glabrous, triangular in cross-section. **Leaves** basal and caudine; basal leaves ca. 5–7 with open sheaths; sheaths gradually widening into the blade, paler than blade, plicate, 12–15 × 1.4–1.8 cm; blade lanceolate to linear, 15–90 × 2–2.4 cm, dark green, usually shiny, the apex acuminate, the margins slightly antrorsely scabrous; caudine leaves (bracts) each subtending an axillary synflorescence and diminishing in size distally along the culm; sheaths closed, the basal ones 2–4 cm long. **Synflorescences** 4–8, each a condensed raceme of 3–6 clusters of 2–3 spikelets each, mature basal one 1.4–3.5 × 1.5–3 cm, the peduncles decreasing in length at each more distal node, the basal one 6.5–8 cm long. **Spikelets** 10–20 per synflorescence; 3.7 × 2.9 mm at anthesis, lenticular, ovate in outline, pale brown; scales ca. 8 visible, distichously or subdistichously arranged, carinate, shiny, cartilaginous, the lowest 2–3 sterile, the lower ones mucronate, the intermediate ones largest, shallowly triangular, 2.8 × 3.5 mm. **Perianth** bristles persistent, 5, flattened basally, antrorsely scabrous to short-plumose, unequal, (1–) 1.5–2.4 mm long, usually one half to two thirds the length of the achene. **Achenes** 2.4–3.2 × (1.7–) 2.2–2.9 mm, deeply biconvex, broadly obovate to circular in outline, shiny, lightly wrinkled to obscurely rugulose; style base inserted in a shallow depressed area at apex of achene and decurrent along lateral margins, the depressed area transversely rhombic, 0.9–1.2 × (1.7–) 2–2.5 mm, the raised central portion 0.6–1.2 × 0.8–1.0 mm, deltate.

Distribution and ecology:—Known from the Atlantic forest region of Brazil, from the state of Bahia south to the state of Santa Catarina (Fig. 2). The known collections are from tropical submontane moist forest (Thomas & Barbosa 2008) above 300 m in Bahia and Espírito Santo and below 300 m in Rio de Janeiro and Santa Catarina, usually among rocks. The flora of one locality, the Serra do Teimoso (Fig. 3), has been well-studied (Amorim *et al.* 2005) and a study of the species composition shows that the forest near the top of the mountain, where *Pleurostachys arcuata* was collected, is distinct from that of the lower forests (Thomas *et al.* 2009b).

Etymology:—The species is named for its characteristic arched culms.

Conservation Status:—*Pleurostachys arcuata* is restricted to intact, rocky, submontane moist forests, and is never common. Furthermore, the Atlantic forest of Brazil has been identified as one of the World's biodiversity "hotspots" (Myers *et al.* 2000), in part, because of continuing deforestation. Overall, the Atlantic forest has been reduced to less than 11 percent of its original extent (Ribeiro *et al.* 2009).

Calculating "area of occupancy" (AOO; IUCN 2012) using grid cells of 3 × 3 km (9 km²) following the methodology developed by Callmander *et al.* (2007) and Miller & Porter Morgan (2011) results in an AOO for *P. arcuata* of 126 km². Calculating "extent of occurrence" (EOO; IUCN 2012) results in an EOO of ca. 160,000 km² (the 1600 km between the northernmost and southernmost known localities multiplied by the average width of the Atlantic forest, ca. 100 km). The AOO estimate is probably too restrictive and does not take into account areas where the species probably occurs but has never been collected. The EOO estimate, on the other hand, is too broad and does not factor in range discontinuity, habitat specificity, or deforestation. We believe the most accurate reflection of its conservation status would be an IUCN Red List category of Vulnerable (VU). More collections would probably increase the AOO of *P. arcuata* significantly, but to less than 2000 km². Furthermore, it occurs in a severely fragmented biome, one that continues to suffer deforestation that will diminish the number and location of its subpopulations.



FIGURE 1. A–G. *Pleurostachys arcuata*. A. Plant habit. B. Synflorescence in fruit. C. Detail of one branch of synflorescence. D. Synflorescence in flower. E. Spikelet prior to anthesis. F. Spikelet scale. G. Achene, side view, top view, and detail of bristle. A–C, F, G from W. W. Thomas et al. 11367 (NY); D and E from W. W. Thomas et al. 14182 (NY). Drawn by Bobbi Angell.

Paratypes:—BRAZIL. Bahia: Mun. Almadina, Serra Corcovado, 9.9 km SW of Coarci on road to Almadina, then N into Fazenda São José, 14°42'21"S, 39°36'12"W, 650–750 m, submontane/montane tropical moist forest with disturbed areas, herb in forest, culms arching, 19 Sep 2004, *W. W. Thomas et al.* 14812 (CEPEC, NY); Serra do Corcovado, acesso pela Estação da EMBASA, 14°42'09"S, 39°36'14"W, 4 Nov 2011, *M. M. Coelho et al.* 486 (CEPEC); Serra do Sete-Paus, Rodovia de Almadina para Ibitupã, 14°44'06"S, 39°41'46"W, *P. Fiaschi et al.* 2730 (CEPEC); Mun. Arataca, Caminho das Pedras, 15°10'25"S, 39°20'30"W, 1000 m, 12 Oct 2005, *A. M. Amorim et al.* 5266 (CEPEC), 15 Jun 2006, *A. M. Amorim et al.* 6065 (CEPEC); Serra das Lontras, 15°12'10"S, 39°24'29"W, 29 Apr 2006, *A. M. Amorim et al.* 5949 (CEPEC); Mun. Jussari, Rod. Jussari/Palmeira, 7.5 km, Fazenda Teimoso, Reserva Serra do Teimoso, 15°09.37S, 39°31.74"W, 300–640 m, 15 Sep 2001, *J. Jardim et al.* 3922 (CEPEC, NY); 21 Aug 2003, *P. Fiaschi, S. C. Sant'Ana & J. L. Paixão* 1579 (CEPEC, NY); near top of mountain, 15°05'44"S, 39°32'33"W, *W. W. Thomas et al.* 13370 (CEPEC); Mun. Una [Santa Luzia], Serra da Onça, 23 Mar 2005, *J. Paixão, S. Sant'Ana & L. Carlos* 378 (NY-2 dupls). Espírito Santo: Mun. Santa Teresa, Estação Biológica da Caixa D'Agua, 4 Nov 1986, *W. Boone* 1081 (MBML, MO); Mun. Cariacica: Reserva Biológica de Duas Bocas, 1 Jun 2000, *M. Alves et al.* 1945 (MBML, SP). Paraná: Mun. Guarqueçaba, Rio de Costa. Mata pluvial encosta de morro, 50 m. 4 Feb 1971. *G. Hatschbach* 26270 (NY). Rio de Janeiro: Mun. Mangaratiba, RPPN Rio das Pedras, *A. M. Amorim et al.* 3378 (CEPEC); Mun. [probably Rio Novo], Serra do Henrique, prope Rio Novo. Sep 1894. *Schwacke* 10973 (RB). Santa Catarina: Mun. Blumenau, Parque Natural Municipal São Francisco de Assis, em floresta Atlântica, 26°55'19.82"S, 49°04'44.87"W, 87 m, 18 Feb 2011, *M. Verdi* 5889 (FLOR, FURB).




FIGURE 2. Distribution map of *Pleurostachys arcuata*.

Observations:—In Kükenthal's (1952) classification, *Pleurostachys arcuata* would belong to section *Millegranae* (Thomas & Alves 2008) because of the presence of blades on the basal leaves and the congested synflorescences. In their phylogeny of *Pleurostachys* based on morphological characters, however, Thomas and Alves (2008) found that section *Millegranae* was probably polyphyletic, and that *P. arcuata* (designated as “sp. nov. 1”) was sister to all other species of the genus.

The only other species with achenes similar in size to those of *Pleurostachys arcuata* is *P. macrantha* Kunth with achenes $2.8\text{--}3.4 \times 2.7\text{--}3.2$ mm. *Pleurostachys macrantha*, however, has large, open panicles with solitary spikelets at the end of each branch. The remaining species of *Pleurostachys* have achenes approximately 1 mm long or less.



FIGURE 3. Forested slope of the Serra do Teimoso Private Reserve (RPPN), Jussari, Bahia.

Acknowledgements

Financial support for field work resulting in collections of this species was provided by the John D. and Catherine T. MacArthur Foundation, the National Science Foundation (DEB 9972116 and DEB 0516233), and the Beneficia Foundation. We thank the curators of the herbaria we consulted, especially those of CEPEC, MBM, NY, R, and RB.

References

- Amorim, A.M., Fiaschi, P., Jardim, J., Thomas, W.W., Clifton, B. & Carvalho, A.M. (2005) The vascular plants of a forest fragment in southern Bahia, Brazil. *Sida* 21(3): 1727–1757.
- Boeckeler, O. (1875) Die Cyperaceen des königlichen Herbariums zu Berlin. *Linnaea* 9: 1–152.
- Bruhl, J.J. (1995) Sedge genera of the world: relationships and a new classification of the Cyperaceae. *Australian Journal of Systematic Botany* 8: 125–305.
<http://dx.doi.org/10.1071/SB9950125>
- Callmander, M.W., Schatz, G.E., Lowry II, P.P., Laivao, M.O., Raharimampoinona, J., Andriambololonera, S. et al. (2007) Identification of priority areas for plant conservation in Madagascar using Red List Criteria: rare and threatened Pandanaceae indicate sites in need of protection. *Oryx* 41: 168–176.
<http://dx.doi.org/10.1017/S0030605307001731>
- Clarke, C.B. (1908) New Genera and Species of Cyperaceae. *Kew Bulletin*, Add. Ser. 8: 1–196.
- Eiten, L. (1976) Inflorescence units in the Cyperaceae. *Annals of the Missouri Botanical Garden* 63: 81–112.
<http://dx.doi.org/10.2307/2395224>
- Goetghebeur, P. (1986) *Genera Cyperacearum. Een bijdrage tot de kennis van de morfologie, systematiek en fylogeneese van de Cyperaceae—genera*. Doctoral Thesis, Rijkuniversiteit Gent.
- Goetghebeur, P. (1998) Cyperaceae. In: K. Kubitzki, H. Huber, P. Rudall, P. Stevens & T. Stützel, *Vascular flowering plants*. Vol. 4. Springer, Berlin., pp. 141–189
- IUCN (2012) The IUCN Red List of Threatened Species. Version 2012.2. At www.iucnredlist.org. Viewed on 13 December 2012.
- Koyama, T. (1961) Classification of the family Cyperaceae 1. *Journal of the Faculty of Science, University of Tokyo, sect. 3, Botany* 8: 37–148.
- Kükenthal, G. (1952) Vorarbeiten zu einer Monographie der Rhynchosporoideae. *Pleurostachys. Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie* 75: 451–497.
- Miller, J.S. & Porter Morgan, H.A. (2011) Assessing the effectiveness of Madagascar's changing protected areas system: a case study of threatened Boraginales. *Oryx* 45: 201–209.
<http://dx.doi.org/10.1017/S0030605310000803>
- Myers, N., Mittermeier, R.A., Mittermeier, C.G., da Fonseca, G.A.B. & Kent, J. (2000) Biodiversity hotspots for conservation priorities. *Nature* 403: 853–858.
<http://dx.doi.org/10.1038/35002501>
- Pfeiffer, H. (1925) Monographia Pleurostachydearum. I. Pars specialis descriptioque specierum. *Botanisches Archiv* 9: 225–242.
- Ribeiro, M.C., Metzger, J.P., Martensen, A.C., Ponzoni, F.J. & Hirota, M.M. (2009) The Brazilian Atlantic Forest: How much is left, and how is the remaining forest distributed? Implications for Conservation. *Biological Conservation* 142: 1141–1153.
<http://dx.doi.org/10.1016/j.biocon.2009.02.021>
- Soros, C. & Bruhl, J.J. (2000) Multiple evolutionary origins of C₄ photosynthesis in the Cyperaceae. Pp. 629–636 in K. L. Wilson & D. A. Morrison, *Monocots: Systematics and Evolution*. CSIRO, Melbourne.
- Thomas, W.W. & Alves, M. (2008) Towards a Revision of the Genus *Pleurostachys* (Cyperaceae): Preliminary Results, In: R. Naczi and B. Ford (eds.) *Sedges: Uses, Diversity, and Systematics of the Cyperaceae*. *Monographs in Systematic Botany from the Missouri Botanical Garden* 108: 269–278.
- Thomas, W.W., Araújo, A.C. & Alves, M. (2009a) (2008 online) A Preliminary Molecular Phylogeny of the Rhynchosporoideae (Cyperaceae). In: W. W. Thomas, D. A. Simpson, A. A. Reznicek and J. R. Starr (eds.), *Cyperaceae – Special Issue. Botanical Review* 75: 22–29.
<http://dx.doi.org/10.1007/s12229-008-9023-7>
- Thomas, W.W. & Barbosa, M.R.V. (2008) Natural Vegetation Types in the Brazilian Atlantic Coastal Forest North of the Rio Doce. In: W. Thomas (ed.), *The Atlantic Coastal Forest of Northeastern Brazil. Memoirs of the New York Botanical Garden* 100: 6–20.
- Thomas, W.W., Jardim, J.G., Fiaschi, P., Mariano-Neto, M. & Amorim, A.M. (2009b) Composição Florística e Estrutura do Componente Arbóreo de uma Área Transicional de Floresta Atlântica no Sul da Bahia, Brasil. *Revista Brasileira de Botânica* 32 (1): 41–54.
<http://dx.doi.org/10.1590/S0100-84042009000100007>