



The male of *Mesoveloidea williamsi* Hungerford, 1929 revisited

FELIPE FERRAZ FIGUEIREDO MOREIRA^{1,3}, JOSÉ RICARDO INACIO RIBEIRO^{2,4} & JORGE LUIZ NESSIMIAN^{1,5}

1. Departamento de Zoologia, Instituto de Biologia, Universidade Federal do Rio de Janeiro, Caixa Postal 68044, Cidade Universitária, 21944-970, Rio de Janeiro, RJ, Brasil.

2. Centro de Ciências Rurais de São Gabriel (CCRSG), Universidade Federal de Santa Maria, UNIPAMPA, Avenida Antônio Mercado 1357, Centro, 97300-000, São Gabriel, RS.

3. felipento@hotmail.com

4. belostom@acd.ufrj.br

5. nessimia@acd.ufrj.br

The present authors recently published in *Zootaxa* (Moreira et al. 2006) a description of the male of *Mesoveloidea williamsi* Hungerford, 1929 and new records for southeastern Brazil. The records therein presented are still valid; however, we overlooked the description of the male form published by Jaczewski (1931). Therefore, Moreira et al. (2006) actually presented a redescription of the male of *Mesoveloidea williamsi*.

Nevertheless, the redescription is useful because it is much more detailed than the one presented by Jaczewski (1931). We also presented several measurements and drawings together with the textual description, while the original description only contained body-length range and drawings of the forewing and paramere.

References

- Jaczewski, T. (1931) The male of *Mesoveloidea williamsi* Hungerford (Hemiptera : Heteroptera). *Proceedings of the Entomological Society of Washington*, 33, 64–66.
- Moreira, F.F.F., Ribeiro, J.R.I. & Nessimian, J.L. (2006) Description of the male and the southernmost record of *Mesoveloidea williamsi* Hungerford, 1929 (Hemiptera: Heteroptera: Mesoveliidae). *Zootaxa*, 1269, 51–56.