



MIRAB-BALOU, M., HU, Q.L., FENG, J.N. & CHEN, X.X. (2011) A new species of Sericothripinae from China (Thysanoptera: Thripidae), with two new synonyms and one new record. *Zootaxa*, 3009, 55–61.

Abstract

One new species of subfamily Sericothripinae (Thysanoptera: Thripidae) is described and illustrated: *Hydatothrips longjingensis* **sp. n.**, and *H. abdominalis* (Kurosawa) is newly recorded from China. The following taxonomic changes are established: *Sericothrips houji* (Chou & Feng) **comb. n.** (from *Hydatothrips*).

Should read as:

Abstract

One new species of subfamily Sericothripinae (Thysanoptera: Thripidae) is described and illustrated: *Hydatothrips longjingensis* **sp. n.**, and *H. abdominalis* (Kurosawa) is newly recorded from China. The following taxonomic changes are established: *Sericothrips houji* (Chou & Feng) **comb. n.** (from *Hydatothrips*), *Sussericothrips melilotus* Han synonymized with *Sericothrips houji* (Chou & Feng), *H. jiawensis* Chou & Feng synonymized with *H. proximus* Bhatti.

Acknowledgements

We are grateful to her critical comments on the draft. We also thank Prof. Ge-Xia Qiao of the Institute of Zoology, Chinese Academy of Sciences, Beijing, for her help in checking type specimens of Han's collection; Prof. Hong Lu and Dr. Shu-Jun Wei from Beijing Academy of Agriculture & Forestry Sciences, for their help in field sampling. The first author wishes to thank Mrs. Shaojie Cao and Mr. Qiu-Lei Men for their help during a visit to NWFU; and Li-Hong Dang for her help during a visit to Institute of Zoology, Chinese Academy of Sciences, Beijing.

Should read as:

Acknowledgements

We are grateful to Dr. Laurence Mound of CSIRO Entomology, Canberra, Australia and Dr. Masami Masumoto of Yokohama Plant Protection Station, Yokohama, Japan for their critical comments on the draft. We also thank Prof. Ge-Xia Qiao of the Institute of Zoology, Chinese Academy of Sciences, Beijing, for help in checking type specimens of Han's collection; Prof. Hong Lu and Dr. Shu-Jun Wei from Beijing Academy of Agriculture & Forestry Sciences, for their help in field sampling. The first author wishes to thank Mrs. Shaojie Cao and Mr. Qiu-Lei Men for their help during a visit to NWFU; and Li-Hong Dang for her help during a visit to Institute of Zoology, Chinese Academy of Sciences, Beijing.