

Description of immatures of the genus *Tropisternus* Solier, subgenus *Pristoternus* Orchymont (Coleoptera: Hydrophilidae), with emphasis on the chaetotaxy and morphometry of larval forms

PATRICIA L. M. TORRES¹, MARIANO C. MICHAT¹, MIGUEL ARCHANGELSKY²
& LILIANA A. FERNÁNDEZ³

¹IBBEA-CONICET, Laboratorio de Entomología, DBBE, FCEyN, Universidad de Buenos Aires, Argentina.
E-mail: patricia.l.m.torres@yahoo.com.ar

²CONICET- LIESA, Laboratorio de Investigaciones en Ecología y Sistemática Animal, Universidad Nacional de La Patagonia, San Juan Bosco, Sarmiento 849, 9200 Esquel, Chubut, Argentina. E-mail: hydrophiloidea@yahoo.com.ar

³CONICET - División Entomología, Museo de La Plata, Paseo del Bosque s/n, 1900 La Plata, Argentina.
E-mail: liliafer@scrym.unlp.edu.ar

Abstract

The egg cases, larvae, and pupae of *Tropisternus (Pristoternus) latus* (Brullé, 1837), *T. (P.) ovalis* Castelnau, 1840, and *T. (P.) laevis* (Sturm, 1826) are illustrated and described for the first time. The egg cases are built with or without a mast, which can be ribbon-like or globular; all larval instars have seven anterior denticles in the anterior margin of nasale, and the antennal seta AN8 is present; the pupae bear 6 or 8 styli on the abdominal segments II–VI, the pleural styli can be either short or very long, and horn-like projections may be present on the abdominal segments II–VI. A comparison of the morphology of the egg cases, pupae and larval chaetotactic and morphometric characters within the genus *Tropisternus* is presented.

Key words: Water scavenger beetles, Hydrophilinae, larva, pupa, egg case

Introduction

The genus *Tropisternus* Solier includes 63 species of aquatic beetles belonging to the family Hydrophilidae (Hansen, 1999; Short & Fikáček, 2011). Species of *Tropisternus* are very frequently collected and often ubiquitous, inhabiting many types of aquatic habitats, from rain puddles and small limnotopes to large lakes and ponds. Adults are detritivorous and can be easily recognized on the basis of their navicular shape, body length less than 15 mm, and the presence of an acute sternal keel. Larvae are carnivorous and typically ambush predators, feeding on a large diversity of prey (Wilson, 1923; Formanowicz & Brodie, 1988). In recent years, several works dealing with the morphology of the preimaginal stages of Neotropical *Tropisternus* have been published (Fernández *et al.*, 2000; Torres *et al.*, 2008, 2011). The genus is presently subdivided into five subgenera: *Tropisternus s. str.*, *Pleurhomus* Sharp, *Homostethus* Orchymont, *Pristoternus* Orchymont, and *Strepitornus* Hansen. *Pristoternus* Orchymont contains 30 species (Hansen, 1999; Spangler & Short, 2008) and is restricted to the Neotropics and the extreme southern Nearctic region. Within *Pristoternus sensu* Hansen (1999), *T. laevis* (Sturm, 1826) and *T. ovalis* Castelnau, 1840 are widely distributed, ranging from Central America to southern South America. On the contrary, *T. latus* (Brullé, 1837) has a more restricted distribution, being known only from specific locations in Brazil (Rio de Janeiro), Uruguay (Montevideo) and Argentina (Buenos Aires and Entre Ríos Provinces).

This paper is meant to be a step towards a better knowledge of the genus *Tropisternus*, and has the following objectives: (i) to describe and illustrate the egg cases, the three larval instars and the pupae of *T. latus*, *T. ovalis* and *T. laevis*; (ii) to perform detailed morphometric and chaetotactic studies on selected larval structures; and (iii) to compare the immatures described here with those of other members of the genus.

thank A. G. Pfaar for helping with the rearing of *T. latus*. This work was supported by projects PICT-2010-0526 (ANPCyT), PIP-5924 (CONICET) and UBACyT-20020100200199 (UBA). This is Scientific Contribution n° 102 from LIESA.

References

- Angus, R.B. (1973) The habitats, life histories and immature stages of *Helophorus* F. (Coleoptera: Hydrophilidae). *Transactions of the Royal Entomological Society of London*, 125, 1–26.
<http://dx.doi.org/10.1111/j.1365-2311.1973.tb00535.x>
- Archangelsky, M., Torres, P.L.M. & Michat, M.C. (2004) Immature stages of *Hydrobiomorpha spinosa* (Orchymont, 1928) (Coleoptera: Hydrophilidae). *Aquatic Insects*, 26, 253–264.
<http://dx.doi.org/10.1080/0165-0420400000593>
- Brullé, A. (1837) Famille des Hydrophiliens. In: Orbigny, A. (Ed.), *Voyage dans l'Amérique méridionale. Vol. 6. Part 2. Insectes (1836–1837)*. P. Bertrand, Paris, Strasbourg, pp. 52–56.
- Byttebier, B. & Torres, P.L.M. (2009) Description of the preimaginal stages of *Enochrus (Hugoscottia) variegatus* (Steinheil, 1869) and *E. (Methydrus) vulgaris* (Steinheil, 1869) (Coleoptera: Hydrophilidae), with emphasis on larval morphometry and chaetotaxy. *Zootaxa*, 2139, 1–22.
- Castelnau, F.L. (1840) *Histoire naturelle des Animaux articulés Histoire naturelle des Insectes Coléoptères. Vol. 2. Nécrophages-Trimères*. P. Duménil, Paris, 565 pp., 38 pls.
- Fernández, L.A., Bachmann, A.O. & Archangelsky, M. (2000) Nota sobre Hydrophilidae neotropicales (Coleoptera) II. Nuevos taxones de *Tropisternus*. *Revista de la Sociedad Entomológica Argentina*, 59, 185–197.
- Fernández, L.A., Torres, P.L.M., Michat, M.C., Fischer, S., Oliva, A. & Bachmann, A.O. (2010) Coleópteros acuáticos y semiacuáticos del Parque Provincial Ernesto Tornquist (Provincia de Buenos Aires, Argentina). *Revista de la Sociedad Entomológica Argentina*, 69, 189–199.
- Fikáček, M., Archangelsky, M. & Torres, P.L.M. (2008) Primary chaetotaxy of the larval head capsule and head appendages of the Hydrophilidae (Coleoptera) based on larva of *Hydrobius fuscipes* (Linnaeus, 1758). *Zootaxa*, 1874, 16–34.
- Formanowicz, D.R. & Brodie, E.D. (1988) Prey density and ambush site changes in *Tropisternus lateralis* larvae (Coleoptera: Hydrophilidae). *Journal of the Kansas Entomological Society*, 61, 420–427.
- Hansen, M. (1999) *World Catalogue of Insects. Hydrophiloidea (s. str.) (Coleoptera)*. Vol. 2. Apollo Books, Stenstrup, Denmark, 416 pp.
- Laker, A.G. (1881) The cocoons of the *Hydrophilus piceus* and *Hydrobius fuscipes*. *Newman's Entomologist*, 14, 82–84.
- Lykins, M.D. (1972) Genetic and environmental factors influencing egg case mast lenght in *Tropisternus collaris* (Coleoptera: Hydrophilidae). *Dissertation Abstracts International B*, 33, 251–252.
- Lyonet, P. (1832) *Recherches sur l'anatomie et les métamorphoses de différentes espèces d'insectes*. J.-B. Baillière, Librairie de l'Académie royale de Médecine, 699 pp.
- Minoshima, Y. & Hayashi, M. (2011a) Larval morphology of the genus *Hydrocassis* Fairmaire (Coleoptera: Hydrophilidae). *Journal of Natural History*, 45, 2757–2784.
<http://dx.doi.org/10.1080/00222933.2011.602805>
- Minoshima, Y. & Hayashi, M. (2011b) Larval morphology of the Japanese species of the tribes Acidocerini, Hydrobiusini and Hydrophilini (Coleoptera: Hydrophilidae). *Acta Entomologica Musei Nationalis Pragae*, 51, 1–118.
- Minoshima, Y. & Hayashi, M. (2012a) The first instar larva of *Hydrobius pauper* Sharp (Coleoptera, Hydrophilidae). *Elytra*, 2, 279–284.
- Minoshima, Y. & Hayashi, M. (2012b) Larval morphology of *Amphiops mater* Sharp (Coleoptera: Hydrophilidae: Chaetarthriini). *Zootaxa*, 3351, 47–59.
- Minoshima, Y., Hayashi, M., Kobayashi, N. & Yoshitomi, H. (2013) Larval morphology and phylogenetic position of *Horelophopsis hansenii* Satô et Yoshitomi (Coleoptera, Hydrophilidae, Horelophopsinae). *Systematic Entomology*, 38, 708–722.
- Minoshima, Y., Iwata, Y. & Hayashi, M. (2012) Morphology of the immature stages of *Hydrochara libera* (Sharp) (Coleoptera, Hydrophilidae). *Elytra*, 2, 285–302.
- Oliva, A., Fernández, L.A. & Bachmann, A.O. (2002) Sinopsis de los Hydrophiloidea acuáticos de la Argentina (Insecta, Coleoptera). *Monografías del Museo Argentino de Ciencias Naturales*, 2, 1–67.
- Portier, P. (1911) Recherches physiologiques sur les insectes aquatiques. *Archives de zoologie expérimentale et générale*, 5, 89–379.
- Short, A.E.Z. & Fikáček, M. (2011) World catalogue of the Hydrophiloidea (Coleoptera): additions and corrections II (2006–2010). *Acta Entomologica Musei Nationalis Pragae*, 51, 83–122.
- Spangler, P.J. & Short, A.E.Z. (2008) Three new species of Neotropical *Tropisternus* Solier (Coleoptera: Hydrophilidae). *Zootaxa*, 1917, 65–68.
- Sturm, J. (1826) *Catalog meiner Insecten-Sammlung. Erster Theil. Käfer*. J. Sturm, Nurnberg, 8 + 207 pp.
- Torres, P.L.M., Mazzucconi, S.A. & Michat, M.C. (2007) Los coleópteros y heterópteros acuáticos del Parque Nacional El

- Palmar (Provincia de Entre Ríos, Argentina): lista faunística, diversidad y distribución. *Revista de la Sociedad Entomológica Argentina*, 66, 127–153.
- Torres, P.L.M., Michat, M.C. & Archangelsky, M. (2008) Description of the preimaginal stages of three species of the genus *Tropisternus* Solier, subgenus *Strepitornus* Hansen (Coleoptera: Hydrophilidae), with emphasis on morphometry and chaetotaxy. *Zootaxa*, 1702, 1–25.
- Torres, P.L.M., Michat, M.C. & Archangelsky, M. (2011) The immature stages of *Tropisternus (Pleurhomus) sahlbergi* (Sharp) (Coleoptera: Hydrophilidae), with considerations on the primary chaetotaxy of *Tropisternus* Solier. *Zootaxa*, 2777, 25–40.
- Torres, P.L.M., Michat, M.C., Libonatti, M.L., Fernández, L.A., Oliva, A. & Bachmann, A.O. (2012) Aquatic Coleoptera from Mburucuyá National Park (Corrientes Province, Argentina). *Revista de la Sociedad Entomológica Argentina*, 71, 57–71.
- Vlasblom, A.G. & Wolvekamp, H.P. (1957) On the function of the funnel on the nest of the water beetle, *Hydrous piceus* L. *Physiologia comparata et oecologia*, 4, 240–246.
- Wiley, E.O. (1981) *Phylogenetics. The theory and practice of phylogenetic systematics*. John Wiley & Sons, New York, 439 pp.
- Wilson, C.B. (1923) Water beetles in relation to pondfish culture with life histories of those found in fishponds at Fairport, Iowa. *Bulletin of the Bureau of Fisheries*, 39, 231–345.