



## Incidence and seasonal occurrence of mites (Acari) on coconut in West Bengal, India\*

DEBJANI BANERJEE & SALIL K. GUPTA

Medicinal Plant Research Unit, Ramakrishna Mission, Narendrapur, Kolkata - 700103; India;

E-mails: debjaniganguly71@yahoo.com; salil\_zsidumdum@yahoo.com, respectively

\* In: Moraes, G.J. de & Proctor, H. (eds) Acarology XIII: Proceedings of the International Congress. Zoosymposia, 6, 1–304.

### Abstract

Coconut (*Cocos nucifera* L.) is an important crop in West Bengal, India. The coconut mite, *Aceria guerreronis* Keifer (Eriophyidae), is becoming a serious threat to its cultivation in that state. This mite lives underneath the perianth, attacking the epidermis and causing damage that leads to considerable yield reduction. The same habitat was found to be shared by several other mites during a one year study in a coconut orchard in South 24 Parganas District of West Bengal, in 2007–2008. In total, 11 species of nine families were found in this study, including mites of different feeding habits, but mainly predatory and phytophagous mites. *Neoseiulus paspalivorus* (De Leon) was the predominant predator, while *A. guerreronis* was the predominant phytophagous species. Both were found throughout the year. The highest abundance of those species occurred in April and June, respectively. The association of *N. paspalivorus* and *A. guerreronis* has been observed in different countries and in other parts of India.

**Key words:** *Aceria guerreronis*, *Neoseiulus paspalivorus*, biological control.

### Introduction

Coconut (*Cocos nucifera* L.) is one of the most important commercial crops in Indian coastal states, covering an area of 1.79 million hectares and producing nearly 14 trillion nuts per year (Nair, 2000). India produces about 23% of world coconut production of 54 billion nuts. The state of West Bengal produces 324 million nuts annually, over an area of 24,200 hectares. Coconut is used mainly for the production of a soft drink, oil, vinegar, copra, chips and coir. India earns a substantial amount of foreign exchange from the coconut based industries.

Unfortunately, coconut is severely affected in this country by the coconut mite, *Aceria guerreronis* Keifer (Eriophyidae), which causes necrosis and malformation of nuts, as well as premature nut fall. These damage reduce yield and quality of copra, leading to substantial economic losses (Haq & Sobha, 2010). This mite was first described in 1965 from the state of Guerrero, Mexico. It was later found on the Caribbean islands and other parts of the American continent and Africa. In Asia, the mite was first found in India (Sathiamma *et al.*, 1998; Haq, 1999a) and Sri Lanka (Fernando *et al.*, 2002), at the end of the 1990's.

Few studies have been conducted on the coconut mite in West Bengal (Karmakar, 2003). Thus, the objective of the present study was to evaluate the mite species from coconut plants, with special attention to those underneath the perianth, in that state in the course of one year.

### Materials and Methods

This study was conducted in a coconut plantation located near Ramakrishna Mission, Ashram Narendrapur, about 25 km south of Kolkata (21.95°N; 88.89°E). Plants were about 7 m tall and about 15 years old. Samples were collected monthly, between February 2007 and January 2008. Altogether, 20 plants were selected in the plantation, from each of which three nuts (3–4 months old) were taken at each sampling date.

The nuts were taken to a laboratory for examination under a stereomicroscope. The perianth was removed and the mites were collected with a moistened brush from three (2.5 x 2.5 cm) areas

randomly determined over the meristematic region. The mites were then mounted in Heinze's medium, identified and counted. Sporadic samples of other coconut parts as well as fallen nuts were also collected, to determine mite diversity.

## Results

Mites of 11 species belonging to nine families were found in this study (Table 1). These belong to families composed predominantly of predatory (Phytoseiidae, Blattisociidae, Melicharidae and Cheyletidae), phytophagous (Eriophyidae, Tarsonemidae and Tetranychidae), fungivorous (Acaridae) and pollinivorous (Ameroseiidae) species.

*Neoseiulus paspalivorus* (DeLeon) was the dominant predator; its population varied from 7.5 to 24.7 individuals per sampling area (total of 6.25 cm<sup>2</sup>). The highest abundance of this species was found in April 2007. Other predatory mites found were *Amblyseius largoensis* (Muma), *Cheyletus malaccensis* Oudemans, *Lasioseius* sp. and *Proctolaelaps* sp.. The former three species were occasionally found, but the latter was uncommon.

Among the four species of phytophagous mites found under the perianth, *A. guerreronis* was dominant, varying from 56.2 to 80.6 individuals per sampling area. It reached its peak abundance in June 2007. The second dominant phytophagous species was an undetermined tarsonemid mite, whose population varied from 0.9 to 30.2 per sampling area. It occurred throughout the year but was found at highest level in October 2007. *Steneotarsonemus furcatus* DeLeon was very infrequently found under the perianth.

The mould mite, *Tyrophagus putrescentiae* (Schrank) (Acaridae), was very occasionally found on the surface of fallen nuts, whereas the pollen feeding *Neocypholaelaps* sp. (Ameroseiidae) and the phytophagous *Oligonychus indicus* (Hirst) (Tetranychidae) were occasionally found on coconut inflorescences and leaves, respectively.

**TABLE 1.** Densities of mites under the perianth of coconuts (mites/ 6.25 cm<sup>2</sup>) determined in South 24 Parganas District of West Bengal, India.

Family / Species	Sampling dates				Nature of association <sup>1</sup>
	Summer Feb–May 2007	Monsoon Jun–Sep 2007	Post monsoon Oct–Nov Jan 2008	Winter Dec 2007– Jan 2008	
<b>Phytoseiidae</b>					
<i>Amblyseius largoensis</i>	2.1	0.4	-	1.6	A
<i>Neoseiulus paspalivorus</i>	24.7	10.2	7.5	10.0	A
<b>Blattisociidae</b>					
<i>Lasioseius</i> sp.	-	0.1	1.1	1.2	A
<b>Melicharidae</b>					
<i>Proctolaelaps</i> sp.	-	-	0.5	0.4	A
<b>Cheyletidae</b>					
<i>Cheyletus malaccensis</i>	0.2	-	2.7	0.8	A
<b>Eriophyidae</b>					
<i>Aceria guerreronis</i>	57.7	80.6	39.5	56.2	B
<b>Tarsonemidae</b>					
<i>Steneotarsonemus furcatus</i>	1.6	0.4	-	-	B
<i>Undetermined tarsonemid</i>	3.7	0.9	30.2	19.7	B
<b>Tetranychidae</b>					
<i>Oligonychus indicus</i>	7.8	6.7	6.0	-	B
<b>Acaridae</b>					
<i>Tyrophagus putrescentiae</i>	0.6	0.2	5.4	0.8	C
<b>Ameroseiidae</b>					
<i>Neocypholaelaps</i> sp.	0.8	0.4	4.7	6.2	D

<sup>1</sup>A= Predatory, B= Phytophagous, C= Fungivorous and D= Pollinivorous

## Discussion

Some studies have been conducted in southern India to evaluate the fluctuation of the population of *A. guerreronis*. Haq (1999b) reported that the highest population levels of *A. guerreronis* in Kerala, Tamil Nadu and Karnataka occurred in July–August. Pushpa & Nandihalli (2009) reported the highest population level of this mite in Karnataka in April–May. As indicated in the present study, in West Bengal that the maximum population level of *A. guerreronis* occurs in a different period, intermediate to those reported in southern India.

The predatory mites known worldwide in association with *A. guerreronis* until the end of the 1990's were reported by Moraes & Zacarias (2002). Shobha (2004) reported some predatory species associated with *A. guerreronis* in Kerala, including species reported as *Typhlodromus pyri* Scheuten, *Cheyletus cocos*, *Agistemus industani* Gonzalez, *Amblyseius* sp. and *Bdella* sp.. Haq (2001) also reported many predatory mites including *A. largoensis*, *Bdella indicata* and *N. paspalivorus*. Mallik & Puttaswami (2000) reported predators belonging to Bdellidae, Cheyletidae, Cunaxidae, Phytoseiidae, Stigmaeidae, Tarsonemidae and Tydeidae in Karnataka. According to those authors, only Cheyletidae and Phytoseiidae were sufficiently dominant and efficient in suppressing *A. guerreronis*. Ramaraju *et al.* (2002) reported the predator *N. paspalivorus* in association with *A. guerreronis* on the surface of infested coconuts in southern India. They also found a large number of the pollen feeder *Neocypholaelaps stridulans* (Evans) on coconut inflorescences.

## Acknowledgements

To the Secretary and Assistant Secretary of the Ramakrishna Mission Ashram, Narendrapur, for providing infrastructure facilities and encouragements.

## References

- Fernando, L.C.P., Wickramananda, I.R. & Aratchige, N.S. (2002) Status of coconut mite, *Aceria guerreronis* in Sri Lanka. In: Fernando, L.C.P., Moraes, G.J. de & Wickramananda, I.R. (eds) *Proceedings of the International Workshop on Coconut Mite (Aceria guerreronis)*. Coconut Research Institute, Lunuwila, pp. 1–8.
- Haq, M.A. (1999a) Coconut mite threat in Kerala. *Journal of Acarology*, 14, 58–63
- Haq, M.A. (1999b) Distribution of the coconut mite, *Aceria guerreronis* in Peninsular India and adjacent islands. *Entomon*, 24, 371–379.
- Haq, M.A. (2001) Culture and rearing of *Aceria guerreronis* and its predators. *Entomon*, 26, 297–302.
- Haq, M.A. & Sobha, T.R. (2010) Weight loss of copra due to infestation of *Aceria guerreronis*. In: Sabelis, M.W. & Bruin, J. (eds) *Proceedings of the 12<sup>th</sup> International Congress of Acarology*, Amsterdam, pp. 509–510.
- Karmakar, K. (2003) Severity of infestation of coconut mite, *Aceria guerreronis* Keifer (Acari: Eriophyidae) in southern districts of West Bengal. *Annals of Entomology*, 21, 29–31.
- Mallik, B. & Puttaswamy (2000) *Biological control of eriophyid mite on coconut*. Group Meeting on Recent Advances in the Management of Coconut Pests. Central Plantation Crops Research Institute, Kayangulam, pp. 12–13.
- Moraes, G.J. de & Zacarias, M.S. (2002) Use of predatory mites for the control of eriophyid mites. In: Fernando, L.C.P., Moraes, G.J. de & Wickramananda, I.R. (eds) *Proceedings of the International Workshop on Coconut Mite (Aceria guerreronis)*. Coconut Research Institute, Lunuwila, pp. 78–88.
- Nair, C.P.R. (2000) Status of coconut eriophyid mite *Aceria guerreronis* Keifer in India. In: Fernando, L.C.P., Moraes, G.J. de & Wickramananda, I.R. (eds) *Proceedings of the International Workshop on Coconut Mite (Aceria guerreronis)*. Coconut Research Institute, Lunuwila, pp. 9–12.
- Pushpa, V. & Nandihalli, B.S. (2009) Seasonal incidence of coconut perianth mite, *Aceria guerreronis* Keifer around Dharwad. *Karnataka Journal of Agricultural Science*, 22, 203–205.
- Ramaraju, K., Natarajan, K., Sundara Babu, P.C., Palaniswamy, S. & Rabindra, R.J. (2002) Studies on coconut

- eriophyid mite, *Aceria guerreronis* Keifer in Tamil Nadu, India. In: Fernando, L.C.P., Moraes, G.J. de & Wickramananda, I.R. (eds) *Proceedings of the International Workshop on Coconut Mite (Aceria guerreronis)*. Coconut Research Institute, Lunuwila, pp. 13–31.
- Sathiamma, B., Radhakrishnan Nair, C.P. & Koshy, P.K. (1998) Outbreak of a nut infesting eriophyid mite, *Eriophyes guerreronis* (K.) in coconut plantations in India. *Indian Coconut Journal*, 29, 1–3.
- Shobha, T.R. (2004) *Acarine Pests of Crop Plants*. PhD Thesis, Calicut University, India.