

A history of systematic studies of the bees of Cuba (Insecta: Hymenoptera, Anthophila)

JULIO A. GENARO

*York University, Department Biology, 4700 Keele Street, Toronto, ON, M3J 1P3, Canada.
E-mail: jagenaro@yorku.ca*

Abstract

A meticulous search was made for papers related to Cuban bees to define the historical context and periods in which studies of their taxonomy and natural history have occurred. Systematic studies of this group of insects began 230 years ago. Such studies are divided into several stages: European (1763–1896), North American (1896–1944), consolidation and current (1944–present). I present the current state of knowledge of natural history of Cuban bees, the status of the main entomological collections in the country, and experience in the use of bees as pollinators in Cuba.

Key words: bees, Hymenoptera, history of taxonomic studies, natural history, collections, Cuba

Resumen

Se efectuó una búsqueda minuciosa sobre los trabajos relacionados con las abejas de Cuba, en cada contexto histórico, para definir las etapas por la que han pasado los estudios de historia natural. Los estudios sistemáticos de este grupo de insectos comenzaron 230 años atrás. Estos, están divididos en varias etapas: europea (1763–1896), norteamericana (1896–1944), de consolidación y actual (1944–presente). Se presenta el estado actual del conocimiento de la historia natural, las características de las principales colecciones entomológicas y experiencia en la utilización de las abejas como polinizadores.

Palabras claves: abejas, Hymenoptera, historia estudios taxonómicos, historia natural, colecciones, Cuba.

Introduction

Bees, ants and wasps are included in the order Hymenoptera and with the Coleoptera, Lepidoptera and Diptera, form the four most numerous and diverse groups of insects. The

world bee fauna contains around 20 000 species, in seven families (Finnamore and Michener 1993; Michener, 2000), while that of Cuba is less than 100 species in four families (Alayo 1976); the Stenotritidae, Andrenidae and Melittidae not being found there.

Bees are among the most popular insects, and are well known for their economic importance. Noteworthy are their production of honey, nutritious products, wax, propolis and drugs; pollination of many plants; and the advanced forms of social behavior which has evolved several times (Asis 1989, Werthein 1993, O'Toole and Raw 1999, Michener 1974, 2000, Stubbs and Drummond 2001).

The Cuban fauna, which has a high endemic component of 47.3% (Genaro and Tejuca 2001), consists of bees that range in length from less than 4 mm (e.g. *Hylaeus* spp.) to the robust, widely distributed and common carpenter bee, *Xylocopa cubaecola* Lucas which is 25 mm. In coloration bees range from black or reddish-brown, to metallic blue or green as among the Cuban halictids, *Augochlora*, *Agapostemon*, some *Lasioglossum* and *Temnosoma*.

At present, no melittologists live in the West Indies, and few entomologist there have a deep knowledge of these important insects. Faunas as diverse and localized as that of Hispaniola remain almost unknown. Contrary to many other Latin American countries or Antillean islands, Cuba has a long entomological tradition, beginning around 1840 (Alvarez 1958, Pruna 2001). Two Cuban entomologists who had a great impact on the studies of their time were Felipe Poey (1799–1891), the first entomologist of Latin America, and Carlos J. Finlay (1833–1915) whose ideas led to the recognition of the role of mosquitoes in transmission of yellow fever (Hogue 1993, Pruna 2001).

In this paper I present a historical overview of the stages of the systematic studies of Cuban bees, a survey of the main collections in the country and summarise what is known about bee natural history in Cuba.

Bibliographic information was obtained during my visit to different United States natural history museums libraries, from 1993 to 2000, as well as Pastor Alayo's personal library, Cuban libraries of the Institute of Ecology and Systematic (IES), Cuban National Museum of Natural History (MNHNCu) and Cuban National Museum of History of Sciences Carlos J. Finlay. The main Cuban collections of native bees were studied in order to evaluate their state of preservation and number of specimens: IES; MNHNCu; Charles Ramsden Museum, University of Oriente (MCRUO); Faculty of Agricultural Sciences, Central University of Las Villas (FAUCV), Oriental Centre of Ecosystems and Biodiversity (BIOECO), and the private collection of Fernando de Zayas (FZ).

Results

The first studies (1763–1896)

The history of the taxonomy of Cuban bees extends over 230 years. Most studies have been performed by taxonomists outside of Cuba. These, from their respective institutions,


compared the material they were receiving and described the new species that appeared, in order to give them names, as is standard practice in taxonomy. Other species, whose distribution includes the island of Cuba, were studied first in other areas and later found to occur in Cuba.

The first Cuban bee species were described or studied by several European entomologists before 1860: M. Spinola (1840, 1851), I. A. Scopoli (1763), Johann C. Fabricius (1793, 1798, 1804), F. E. Guérin-Méneville (1835, 1844), A. Lepeletier (1841), R. Lucas (in Sagra, 1856) and C. E. Blanchard (in Cuvier, 1849). Due to the lack of detail on localities, the material was generally identified as coming from "Americae Meridionalis Insulis", that is, Cuba or another Caribbean island. Thus one cannot determine which was the first described Cuban bee. Genaro (1998a), however, demonstrated that the species *Coelioxys tridentata* described as *Apis tridentata* by Fabricius, in 1775, has Cuba as its type locality.

The English entomologist F. Smith (1853) catalogued the Hymenoptera deposited in the British Museum, including bees from Cuba, and he later described new species from the West Indies (Smith, 1879). The first author that summarized Cuban Hymenoptera was Guérin-Méneville (in Sagra, 1856). This author received specimens from Poey and Sagra to supplement the work of the latter on the natural history of the Island of Cuba, although bees were not abundant in the material. The only species described was *Xylocopa cubaecola*, by Lucas (Sagra, 1856).

After Sagra's work, and stimulated by two great Cuban zoologists, Felipe Poey and Johann C. Gundlach (Fig. 1), Cresson produced thorough taxonomic revision of Cuban Hymenoptera, including the bees (Cresson 1865, 1869a and b). This included the description of many new species, since Cresson had at his disposal the Hymenoptera collection that Poey sold to the Entomological Society of Philadelphia, and also Gundlach's collection borrowed from him. This gave Gundlach's collection, now at the IES (Figs. 2 and 3), great historical and taxonomic value, since it contains many of Cresson's primary types.

Felipe Poey preferred to study butterflies, from among all the insects. However, he carried out a detailed study (Poey 1851) on the nest structure and nesting behavior of the stingless bee *Melipona beecheii* Bennett (as *M. fulvipes* Guérin-Méneville). Later he wrote other works on the bee *Apis mellifera* L., as a producer of honey (Poey 1856–1858, 1888a and b). Gundlach (1886), continued observations on Cuban entomology and provided the most complete account of the bees available at the time, with taxonomic data, diagnoses and distributions in Cuba. In this paper he also incorporated personal observations from field work, accounts of previously well-known Cuban species, and translated Cresson's (1865, 1869b) descriptions. Dalla Torre (1896) catalogued the Hymenoptera of the world. This work, with many West Indian species, included records of species usually not clearly assigned to individual islands from the Caribbean region.



FIGURES 1–3. Gundlach and his collection. 1. J. C. Gundlach (1810–1896) (*Canvas at Cuban National Museum of History of Sciences*). He published the most complete account of Cuban bees until Alayo's (1973) catalog. 2 and 3. Gundlach's collection is housed at the Institute of Ecology and Systematic (IES), Havana City. This important historical collection holds the largest number of primary insect types in Cuba.



Second stage (1896–1944)

The next stage can be considered North American, because of the prevalence in the relevant literature of specialists from the United States. It was begun by the work of Ashmead (1896, 1900) and Friese (1899, 1900, 1902, 1908, 1911, 1921). Friese, a German author, provided lists of the Antillean bee fauna (Friese 1902, 1908). Both authors described new species from West Indian islands. Crawford (1915) described a Jamaican species (*Melissodes foxi*) and commented on species from Cuba. Cresson (1916) listed the holotypes of the new species described during his studies.

In an important contribution at the beginning of the 20th century, Charles F. Baker (Fig. 4), a North American who directed and worked in the Agronomy Experimental Station (EEA) at Santiago de las Vegas, 12 km SW of Havana, presented keys to separate species of halictines, and listed the bee species known from Cuba (Baker, 1906). He also described new species of *Lasioglossum* (as *Chloralictus*), which are small forms, not

studied previously in Cuba. Unfortunately, this author did not designate holotypes and deposited his material mainly in his private collection, now at the United States National Museum of Natural History, Smithsonian Institution.

Cockerell (1898, 1904, 1905*a, b* and *c*, 1906, 1910*a, b* and *c*, 1919) examined many holotypes in the British Museum and Smithsonian Institution, offered keys to separate some West Indian species of genera *Megachile* and *Coelioxys*, wrote a species list and described new species. Other North American authors (Robertson 1902, Smith 1907, Sandhouse 1923), when studying the fauna of their continent, included descriptions and keys for West Indian species. Mitchell (1927) published new species of *Megachile* from the Antilles, but the two species he described from Cuba were synonyms of previously described species (Genaro 1998*a*). He also (Mitchell 1943) presented keys to separate the Neotropical subgenera of *Megachile*, including Cuban species. Schwarz (1932, 1944) revised the genus *Melipona* and commented about the introduction of *M. beecheii* (as *M. beecheii fulvipes*) into Cuba.



FIGURES 4–5. Entomologists in Cuba, working with bees. 4. Charles Fuller Baker (1872–1927). (From Essig 1931). His paper of 1906 is still useful and the only one available for the proper identification of *Lasioglossum* species, which are important pollinators of angiosperms. 5. Pastor Alayo (1915–2001). This Cuban entomologist contributed most to the study of bees. He never described a new species but collected and sent material to specialists. This allowed him to make a fine reference collection and to write an important introductory paper and catalog.

Consolidation stage and current (1944– 2005)

Charles D. Michener is considered to be the world authority on bees. The publication of Michener (1944), greatly furthered the taxonomic studies of bees. This classic paper defined and established morphological terms that improved the classification and created a

favourable basis for the development of later studies, including phylogenetic analyses. At the University of Kansas, he created the largest collection of references about bees. His work has stimulated the development of specialists all over the world. Pastor Alayo (Fig. 5), a Cuban entomologist, had a continuing relationship with him, and sent material for identification, resulting in several papers published (Michener 1966, 1988*a* and *b*). For example, Michener (1988*b*) recorded the presence of the subfamily Lithurginae in the Antilles for the first time and described *Lithurgus antilleorum*, from the xeric coasts of the Greater Antilles.

His subsequent studies on bee natural history, with an emphasis on identification and phylogeny, are necessary references for any specialist of the group. Particularly important are Michener (1953, 1974, 1986, 2000); Michener and Fraser (1978); Michener *et al.* (1994) and Sakagami and Michener (1987).

In the 1950's, the study of the bees of the Western Hemisphere was continued by North American entomologists. Cross (1958) revised the bees of the genus *Nomia*, subgenus *Epinomia*, including the Cuban species. LaBerge (1956, 1961) revised the genus *Melissodes* in North and Central America, redescribing species, describing new species and presenting keys. Mitchell (1960, 1962) studied the bees of the Eastern United States, including species whose distributions embrace the West Indies. Eickwort (1969) in his comparative morphological study and generic revision of the augochlorine bees deal with major works concerning the generic classification of Latin America Augochlorini. Urban (1970), in Brazil, published on the genus *Florilegus*, contributing keys and much historical data. Roberts (1972) revised the genus *Agapostemon* and offered a key to the species. Mitchell (1973, 1980) contributed novel results and new changes in the subgeneric classification of *Megachile* and *Coelioxys*. Timberlake (1980) revised the American and West Indian species of *Exomalopsis*. The studies of Centridini and some parasitic Apidae by Snelling (1984, 1985, 1986) also included Cuban species and in 1990 he revised the subgenus *Chelostomoides*, of the genus *Megachile* (as *Chalicodoma*), and presented keys and illustrations. Alexander and Schwarz (1994) published a catalog of the species of *Nomada* of the world, placing them into the species groups resulting from Alexander's (1994) phylogenetic studies. Brooks (1999) revised the genus *Anthophora* in the West Indies, and recorded *A. hilaris* Smith from Cuba. Brooks (1983, 1988) had previously published on this genus, including the Cuban species *A. atrata* Cresson. Moure (1960), an outstanding Brazilian taxonomist, had access to Fabricius's type material, deposited in various museums in Europe and clarified numerous nomenclatural problems of Western Hemisphere species.

P. Alayo promoted the systematic knowledge of Cuban bees. In the beginning (1931–1951) he was associated with S. C. Bruner, head of the Agronomic Experimental Station of Cuba. Later, he worked at the Institute of Biology of the Academy of Sciences in Havana. He collected bees and formed a very important collection, mainly from the eastern areas of the island. He sent much material to the most active specialists of his time. Since he feared

to create synonyms and because he could not obtain permission to travel outside of Cuba, the impossibility of studying the holotypes of Cuban species deposited in North American museums prevented him from describing new species. He wrote a series of introductory articles about the orders of Cuban insects that stimulated and facilitated their study. Alayo (1973) classified the Hymenoptera of Cuba, in the most complete compilatory work to date. He reviewed the Apoidea and presented keys to separate the families, subfamilies, tribes and genera of bees, besides listing the known species (Alayo 1976).

Krombein *et al.* (1979) included species whose distributions embrace Cuba in their catalog of North American bees. Hurd (1978) classified the species of *Xylocopa* of the Western Hemisphere. Zayas (1981), as a part of his valuable overview of the Cuban Entomofauna, wrote about hymenopterans, offering pleasing illustrations, and comments in clear language on the Cuban bees. Unfortunately, his book was published on poor quality paper. Moure and Hurd (1987) presented much information in their catalog on the halictids of the Western Hemisphere. Carman and Packer (1997), in an electrophoretic study on *Halictus*, found differences among some studied populations and recognized two species existing under the name *H. ligatus* Say. These authors used the name *H. poeyi* Lepeletier, for the Southern form (Southeast of United States, Central America and Caribbean), which also occurs in Cuba, the type locality.

Ayala (1992) revised the stingless bees of Mexico (genus *Melipona*). This study included *M. beecheii* present in Cuba and Jamaica. Ayala commented on the wide use of this species in beekeeping (in this case, meliponiculture), and noted that domestication should have influenced its distribution.

The present author (JAG) has contributed to the study of the taxonomy of West Indian bees, describing new species, making taxonomic changes and expanding the known range of some species (Genaro 1994a, 1997, 1998a, b and c; 1999, 2001a, b and c, 2002, 2003, 2004).

Urban and Moure (2001) published a catalog of the subfamily Diphaglossinae, raising (without giving a rationale) *Alayoapis* from subgenus to genus. Moure and Urban (2002), and Urban and Moure (2002) continued with the publication of catalogue that deal with Neotropical species of Colletidae. Nevertheless, they did not include *Colletes granpiedresis* Genaro, in spite of its being published in a Central American journal (Genaro 2001c).

Raw (2002, 2004) has worked on the taxonomy of *Megachile* of the Americas, including Cuban species but overlooked Genaro's (1998a) paper on distribution and synonymy of some Caribbean bees of this genus. Janjic and Packer (2003) performed a phylogenetic analysis of all *Agapostemon* species and made some biogeographic speculations on the history of these bees. Engel (2004) determined the proper identity of the species known as *Halictus poeyi* Spinola. M. Engel (in press, a and b) will soon publish on Cuban bee taxonomy, describing two new species of *Sphecodes* and a new genus closely related to it.

Cuban collections

Scientific collections provide the basis for systematic and biogeographic studies. They contain the primary evidence for the existence of species, document their presence in local areas, and allow comparison and identification of taxa (Danks 1991, McGinley, 1992). Historically, the importance of the creation and preservation of natural history collections has not been widely appreciated in Cuba. The existence of collections has been due fundamentally to the personal efforts of a few dedicated zoologists performing taxonomic research. Among them, the names J. Gundlach, F. Poey, C. Ramsden, S. Bruner, C. de la Torre, C. G. Aguayo, P. Alayo, F. de Zayas, S. de la Torre, H. Grillo, J. de la Cruz, J. Fernández Milera, G. Silva, L. F. de Armas and G. Alayón appear most commonly. The climatic conditions of Cuba (high humidity and temperatures), and easy proliferation of pest organisms, quickly damage collections and elevate the cost of their maintenance.

Table 1 shows the characteristics of the main Cuban entomological collections that contain bees.

Taxonomic keys for identification

Keys allow identification in a quick and accurate way. Many taxonomic studies lack keys to differentiate species, thereby making the identifications that are indispensable as a basis for ecological, behavioral or biogeographical studies impossible. Table 2 lists the authors which presented keys to differentiate Cuban bees.

Other papers that are useful for the recognition of Neotropical bees, through allowing the identification of families, subfamilies, tribes and genera include Alayo (1976), Goulet and Huber (1993), Michener *et al.* (1994) and Michener (2000).

Natural history

Knowledge about the natural history of Cuban bees is limited. Their behavior, in an environment as diverse and with interspecific interactions as complex as in the tropics and subtropics, remains almost unknown. There are just a few papers dealing with these topics from Cuba. Scaramuzza (1938) discussed two parasites of the bee *Megachile* sp; Alayo (1982) observed the characteristics of the nests of *Anthophora atrata* Cresson; Alayón (1984) contributed data on the biology of *Melipona beecheii* (as *M. fulvipes*); Genaro (1994*b*, 1996*a*, *b*, *c* and *d*, 1997, 1998*a*), and Genaro and Sánchez (1992) published articles on interspecific relationships such as predation, inquilines, phoresy and parasitism, and also nesting behavior.

One of the most interesting behaviors appears in bees of the genus *Megachile*, some of which cut leaves of plants to build their nest cells. In a study on the behavior of the genus, Genaro (1996*a*) demonstrated that they have a preference for leguminous leaves, possibly because of the morphoanatomic characteristic of the leaves. Species of *Megachile* used plants of 25 families, of which 24.1% were introduced. The use of exotic plants demonstrated behavioral plasticity.

TABLE 1. Main Cuban collections of native bees with comments on their holdings.

Institution or collection	Aproximate number of specimens	Number of holotypes	Comment
IES	3 208	12 (belonging to Gundlach's collection, Figs. 2 and 3)	Contains the Gundlach historic collection (with many primary types), the P. Alayo collection, and remains from the Santiago de Las Vegas Agronomic Experimental Station (with voucher specimens and complementary data about natural history). Also with recent collections of other entomologists.
MNHNCu	1 885	10	Collection with some primary types, and more accurate and reliable identifications than the other Cuban collections. The only one with material from Hispaniola, Puerto Rico and other West Indian islands.
MCRUO	200	0	Very important because it holds specimens collected at the beginning of 20th century, in natural areas of the East, many of which have disappeared or been degraded. Many specimens were transferred to other collections, mainly those at IES.
FAUCV	543	0	Contains material mainly from central areas of Cuba (especially Guamuaya Massif). Many specimens need identification.
BIOECO	296	0	Contains material from Eastern mountain areas, and around Santiago de Cuba.
FZ	390	0	Private collection of difficult access because owner limits entrance. Specimens with reliable identifications. Its conservation is endangered because of conditions of storage.

Sotolongo *et al.* (1997) studied pollen in brood cells of *Xylocopa cubaecola* and, León and Sánchez (1998) identified the pollen collected by this bee, demonstrating its generalist foraging habits. Díaz and Vale (2001) studied the behaviour of this carpenter bee as a pollinator of the orchid *Encyclia phoenicea* (Lindl.) Neuman.

It is necessary to deepen studies on the natural history of Cuba's endemic taxa, especially those that may illustrate phenomena of broad interest. For example, phylogenetic studies of several groups are important for biogeographical analyses. *Habralictellus* group of the genus *Lasioglossum*, could be of importance in understanding the origin of sociality in insects and species of this genus present different degrees of social organization (Wcislo 1997, Michener 1990). *Habralictellus* is endemic to the West

Indies (Eickwort 1988), and was behaviorally unknown until I (Genaro 2001a) found up to six females of *L. sierramaestrensis* Genaro sharing the parental nest. The amount of cooperation and reproductive status of each female are questions to investigate.

TABLE 2. Taxonomic keys for the identification of species of Cuban bees, in chronological order.

Author and year	Taxa	Geographic area embraced by the key
Cockerell (1905a)	<i>Megachile</i> , <i>Coelioxys</i>	Some Neotropical <i>Megachile</i> species including West Indian ones, and some <i>Coelioxys</i> spp. from all over the world including <i>C. rufipes</i> Guérin from Cuba.
Baker (1906)	<i>Lasioglossum</i> (<i>Dialictus</i>) <i>Agapostemon</i> , <i>Augochlora</i>	Cuba
Smith (1907)	<i>Ceratina</i>	North and Central America, including Cuba
Cockerell (1910a)	<i>Augochlora</i>	Cuba
Schwarz (1932)	<i>Melipona</i>	South and Central America, including Cuba
LaBerge (1956)	<i>Melissodes</i> (<i>Melissodes</i>)	North and Central America, including Cuba
LaBerge (1961)	<i>Melissodes</i> (<i>Eumelissodes</i>)	North and Central America, including Cuba
Michener (1966)	<i>Caupolicana</i> (<i>Alayoapis</i>)	North America and West Indies
Roberts (1972)	<i>Agapostemon</i>	Throughout the Americas, including West Indies
Timberlake (1980)	<i>Exomalopsis</i>	North America, including West Indies
Snelling (1990)	<i>Megachile</i> (as <i>Chalicodoma</i>) (<i>Chelostomoides</i>)	North America, including Cuba
Genaro (1997)	<i>Megachile</i> [<i>Chalicodoma</i> group <i>sensu</i> Michener (2000)]	Cuba
Genaro (1998b)	<i>Ceratina</i>	Cuba and Hispaniola
Genaro (1999)	<i>Triepeolus</i>	Cuba
Brooks (1999)	<i>Anthophora</i>	Wets Indies
Genaro (2003)	<i>Colletes</i>	Cuba

The honey bee, *A. mellifera*, is an introduced bee in the New World (Michener 2000). It has probably deeply influenced the natural populations of other species by competing with native bees for food sources (flowers). Studies have demonstrated that the honey bee influences the patterns of foraging of native bees by competing (Eickwort and Ginsberg 1980, Buchmann 1996, Engel 2001b, Goulson 2003, Buchmann and Ascher 2005). This is because honey bees are social and generalists, so that in a single field hundreds of individuals may be collecting nectar from a wide variety of flowers, and throughout the whole year. This effect has not been investigated in Cuba, where bee hives have been

introduced for the production of honey and for pollination. Honey bee introduction has been done in many places, including protected areas (e.g. Caguanes National Park, Sancti Spiritus; Zapata Swamp National Park and Alejandro de Humbolt National Park, Guantánamo). The impact of beehives in natural areas on the populations of native Cuban bees remains unknown.

We lack quantitative data to evaluate the current status of the populations of Cuban insects and bees are no exception (and this should be a major concern). An endemic bee, *Megachile armaticeps* Cresson (Fig. 6) is on the list of the threatened species of the International Union for the Conservation of Nature (IUCN). It was proposed (Genaro 1998c) as vulnerable due to the fragmentation and loss of its habitat and possible competition with the introduced *Megachile lanata* (Fabr.), which has led to it having a small population restricted to a very few places.



FIGURE 6. *Megachile armaticeps* (female) showing facial projections. This charismatic species is the only Cuban bee listed in the International Union for the Conservation of Nature “Red Book” of endangered and threatened species. There is need for more study to clarify the threatened status of this bee.

Biogeography of West Indian bees

Many workers have interpreted the biogeographic composition of the Neotropical fauna of bees, and consequently that of Cuba. Michener (1979) examined the distribution of the main groups of bees of the world (at the level of genera and subgenera) and speculated on the historical and ecological explanations of such distributions. Michener (2000), with more current data, interpreted the world patterns of distribution of the bees. Eickwort (1988) analyzed the origin, dispersal and biology of the sweat bees (Halictidae) in the Antilles. This work is important because it comments on the group *Habralictellus* (as a genus), endemic to the Antilles, and proposed routes for its arrival in Cuba. Other authors, as parts of their taxonomic studies (e.g. LaBerge 1956, Camargo *et al.* 1988) have tried to explain the distribution of their groups. For Roberts (1972), the dispersion and

evolution of the genus *Agapostemon* in the West Indies was rapid, since none of the species occurs in the neighbouring continents. Janjic and Packer (2003) cited that *Agapostemon* originated in Central America, from there colonized North and South America and penetrated the West Indies at different times, by way of emerged lands or continuous land masses (Iturralde-Vinent and MacPhee 1999).

In Hispaniola, amber (fossil resin) contains insects that have been trapped and among them are bees. This permits comparison of the present fauna with that of the Miocene (Iturralde-Vinent 2001). The Miocene amber fossils bear the character of the modern faunas (Engel 1995, Engel 1999, Grimaldi and Engel 2005). Keeping in mind current paleogeographic models (Iturralde-Vinent and MacPhee 1999) we can assume that much of that fauna was also present in Cuba.

The members of the tribe Euglossini, well-known as orchid bees, inhabit the tropics of America and have developed a remarkable behavior associated with the pollination of orchids (Roubik and Ackerman 1987). In Cuba, representatives of this tribe are absent, in spite of the presence of 305 species of orchids (Llamacho and Larramendi 2005). Two species of orchid bee occur in Jamaica: *Euglossa jamaicensis* Moure and *E. ignita* F. Smith (Moure 1967, Kimsey and Dressler 1986, Engel 1999). This reaffirms the different paleogeographic history of Jamaica, which was associated for a longer time with Central America (Iturralde-Vinent and MacPhee 1999), and thus was populated with species that have not dispersed to Cuba, in spite of its being relatively near but separated by deep sea. Engel (1999) considered that the genus *Euglossa* was, in the past, more widely distributed in the Caribbean than it is today with subsequent extinction occurring, after the Proto Antilles's land mass became fragmented, islands submerged and reconstituted, and effects of climate changes took place.

Local faunal lists

Due to the lack of multidisciplinary studies that inventory the biota of Cuban areas, the difficulty in identifying the material and the absence of taxonomists, very few local listings of bees exist. Invertebrates have always remained behind in such inventories and despite their importance bees are not exception. Among the better known local lists are those of Rohwer and Holland (1917) for the Isle of Youth (as the Isle of Pines); García *et al.* (1973) for the Jibacoa-Cayajabos area of Havana province; and Portuondo and Fernández (2004) for the Eastern Mountains. Fernández *et al.* (2002) offered preliminary data about the Hymenoptera of Granma province, including bees. Genaro (2004) presented a list of the bees from the Isle of Youth and found no endemics, presumably due to its paleogeographic history and land connections to Cuba during low sea levels in the Pleistocene.

Until now, the Africanized honey bee or "killer bee" has not arrived in Cuba, due to the islands's oceanic isolation and infrequent commerce with neighbouring regions, where it has already arrived. This bee, *Apis mellifera scutellata* Lepeletier, introduced to Brazil in

1956, has dispersed widely in a short time. Cuba has strict quarantine control to avoid its entrance. By 1995 it had appeared in the southwest of the United States, extending northward as far as mild winter temperatures permit its survival (Michener 1982, Taylor 1985, Flakus 1993). Its fame resides in its aggressiveness and propensity to attack en masse and, on occasion causing death to mammals, including humans.

Studies on bees which are producers of honey

The tendency exists, even in Cuba, to believe that the term "bee" includes only or mainly the honey bee (*Apis mellifera* L.) that produces honey and wax. However, this species was introduced to Cuba by the middle 1700's (Romay 1796, Díaz 1985, Bande 1996) whereas there are approximately 90 species of autochthonous bees (Alayo 1973, 1976, Genaro 2002).

The Beekeeping Experimental Station (Estación Experimental Apícola) in Havana, belongs to the Ministry of Agriculture. It has promoted and guided studies of *Apis mellifera* and to a lesser extent, *Melipona beecheii*. The two bees produce honey and consequently are of economic importance. There are numerous studies dedicated to *A. mellifera*: organized meetings (e.g. symposia on propolis and apitherapy), journals dedicated to the topic (e.g. Ciencia y Técnica Agrícola, series Apicultura); translated and published information on beehives, diseases, and breeding techniques. Studies carried out in Cuba range from morphometry to biomedicine (composition, antimicrobial properties and medical therapeutic properties of propolis); and components, properties and uses of honey, pollen, royal jelly and bee venom in the treatment of various illnesses (Díaz and Domínguez 1985, Asis 1989, Werthein 1993, Rodríguez *et al.* 1995, Frias *et al.* 2000).

The keeping of stingless bees in Cuba is growing. Colonies are being domesticated and placed near homes by isolated farmers, without any training. They bring feral colonies to their homes and put them inside an empty tree-trunk (usually a palm-trunk) in order to obtain the honey to sweeten foods. For such reasons wild colonies of *M. beecheii* are disappearing from natural areas. In the Yucatán peninsula of Mexico, *M. beecheii* is threatened by environmental changes and by inappropriate management and conservation efforts (Villanueva-G *et al.* 2005). The situation in Jamaica is similar, where this stingless bee species might become extinct (A. Raw, pers. comm., 2005).

Use of bees in the pollination of crops

Historically, beekeeping has been mainly concerned with the production of honey, wax and propolis, with the use of two introduced species: *A. mellifera* and *M. beecheii*. In Cuba, there is no tradition of using bees in pollination. Farmers do not generally understand the advantages of pollination by bees for the improvement of the yield of crops and for obtaining better seeds (Kremen *et al.* 2004, Ricketts *et al.* 2004).

The promotion of the use of bees to pollinate crops began in 1997 (León 1997). Until now, it has been used for watermelon (Hernández and Lemus 1996a, León and Rivero

1999) and pumpkin (Hernández and Lemus 1996b). The species used in pollination have been: *A. mellifera* on a wide scale; *Melipona beecheii* in a few cases; and *Xylocopa cubaecola* occasionally at the Beekeeping Experimental Station for pollinating pumpkins (A. León, pers. comm., 1999). Much of the use of bees as pollinators has been in greenhouses (León and Rivero 1999).

Acknowledgements

The following specialists offered information on their work: Alejandro León, pollinators; Yoandri Suarez, the Charles Ramsden collection; Elba Reyes (IES) and Horacio Grillo, collection of Universidad Central de Las Villas. Laurence Packer offered me facilities of his laboratory and the use of an extensive library and collections. The following entomologists facilitated my work with literature: Charles Michener, Roy Snelling, Giraldo Alayón, Alejandro León, John Ascher and Michael Engel. I thank M. Engel for the reading the first versions of the manuscript. C. Michener, Stewart B. Peck and overall L. Packer carefully reviewed and improved the English language and offered many helpful comments. To Rolando Fernández de Arcila for sending me information about C. Baker. M. Engel and J. Ascher were always willing to share information about West Indian bees. This research was supported by NSERC postdoctoral fellowship (PDF 314411-2005). Vice-President Research and Innovation of York University Stan M. Shapson and Discovery Grants from the NSERC of Canada to S. B. Peck also provided funding for which I am grateful.

References

- Alayo, P. (1973) *Catálogo de los himenópteros de Cuba*. Editorial Pueblo y Educación, Havana city, 218 pp.
- Alayo, P. (1976) Introducción al estudio de los himenópteros de Cuba. IX- Superfamilia Apoidea. *Serie Biológica, Instituto de Zoología*, 68, 1–35.
- Alayo, R. (1982) Observaciones en himenópteros cubanos. I. Apoidea y Sphecoidea. *Poeyana*, 241, 1–7.
- Alayón, G. (1984) Colmena de *Melipona fulvipes* Guérin (Hymenoptera: Apidae) en un comején de bola, *Nasutitermes rippertii* (Rambur) (Isoptera: Termitidae). *Miscelania Zoológica, Instituto de Zoología*, 19, 3–4.
- Alexander, B.A. (1994) Species-groups and cladistic analysis of the cleptoparasitic bee genus *Nomada* (Hymenoptera: Apoidea). *University of Kansas Science Bulletin*, 55, 175–238.
- Alexander, B.A. & Schwarz, M. (1994) A catalog of the species of *Nomada* (Hymenoptera: Apoidea) of the world. *University of Kansas Science Bulletin*, 55, 239–270.
- Alvarez, J. (1958). *Historia de la Zoología en Cuba*. Publicaciones de la Junta Nacional de Arqueología y Etnología. Editorial Lex, Havana city, 369 pp.
- Ashmead, W.H. (1896) Notes on the Hymenoptera collected by the Bahamas Expedition from the State University of Iowa. *Bulletin Laboratory of Natural Sciences, St. University of Iowa*,

- 4,28–32.
- Ashmead, W.H. (1900) Report upon the Aculeate Hymenoptera of the Islands of St. Vincent and Grenada, with additions to the parasitic Hymenoptera of the West Indies. *Transactions of the Entomological Society of London*, 1900, 207–367.
- Asis, M. (1989) *Propóleo, el oro púrpura de las abejas*. Centro de Información y Documentación Agropecuaria, Ministerio de la Agricultura, Havana city, 255 pp.
- Ayala, R. (1992) *Revisión de las abejas sin agujón de México (Hymenoptera: Apidae: Meliponinae)*. Universidad Nacional Autónoma de México, Master Thesis, 67 pp.
- Baker, C.F. (1906) Halictinae de Cuba. *Primer Informe de la Estación Central Agronómica de Cuba*, pp. 253–274.
- Baker, H.G. & Hurd, Jr., P.D. (1968) Intrafloral ecology. *Annual Review of Entomology*, 13, 385–414.
- Bande, J.M. (1996). Contribución al estudio de la historia de la apicultura en Cuba, introducción de la abeja en la Isla. In: *Simposio de Flora Melífera y Polinización, Resúmenes. Facultad Farmacia y Alimentos*. Habana city.
- Brooks, R.W. (1983) Systematics and bionomics of *Anthophora*: the *bomboides* group and species groups of the New World (Hymenoptera: Apoidea, Anthophoridae). *University of California Publications Entomology*, 98, 1–86.
- Brooks, R.W. (1988) Systematics and phylogeny of the Anthophorine bees (Hymenoptera: Anthophoridae; Anthophorini). *University of Kansas Science Bulletin*, 53, 436–575.
- Brooks, R.W. (1999) Bees of the genus *Anthophora* Latreille 1803 (Hymenoptera, Apidae, Anthophorini) of the West Indies. *Tropical Zoology* 12, 105–124.
- Buchmann, S.L. (1996) *Competition between honey bees and native bees in the Sonoran desert and global bee conservation issues*. In Matheson, A. et al. (Eds), *The Conservation of bees*. Linnean Society Symposium Series #18, Academic Press Inc., pp: 125–142.
- Buchmann, S.L. & Ascher, J. S. (2005) The plight of pollinating bees. *Bee World* 86,71–74.
- Carman, G. M. & Packer, L. (1997) A cryptic species allied to *Halictus ligatus* Say (Hymenoptera: Halictidae) detected by allozyme electrophoresis. *Journal of the Kansas Entomological Society*, 69, 168–176.
- Camargo, J.M.F., Moure, J.S. & Roubik, D.W. (1988) *Melipona yucatanica* new species (Hymenoptera: Meliponinae); stingless bee dispersal across the Caribbean arc and post-Eocene vicariance. *Pan-Pacific Entomologist*, 64, 147–157.
- Cockerell, T.D.A. (1898) On some Panurgine and other bees. *Transactions of the American Entomological Society*, 25,185–198.
- Cockerell, T.D.A. (1904) New and little-known bees in the collection of the British Museum. *Annals and Magazine of Natural History*, 14, 203–208.
- Cockerell, T.D.A. (1905a) Tables for the separation of some bees of the genera *Coelioxys* and *Colletes*. *Psyche*, 12, 85–90.
- Cockerell, T.D.A. (1905b) Notes on some bees in the British Museum. *Transactions of the American Entomological Society*, 31, 309–372.
- Cockerell, T.D.A. (1905c) Notes on some bees in the British Museum. *Transactions of the American Entomological Society*, 31,309–364.
- Cockerell, T.D.A. (1906) The North American bees of the family Anthophoridae. *Transactions of the American Entomological Society*, 32, 63–116.
- Cockerell, T.D.A. (1910a) Some bees of the genus *Augochlora* from the West Indies. *Proceedings of the United States National Museum*, 37, 489–494.
- Cockerell, T.D.A. (1910b) The North American bees of the genus *Nomia*. *Proceedings of the United States National Museum*, 38, 289–298.
- Cockerell, T.D.A. (1910c) Some Neotropical bees. *Psyche*, 17, 142–144.
- Cockerell, T.D.A. (1919) Bees in the collection of the United States National Museum, 3. *Proceed-*

- ings of the United States National Museum, 55,167–221.
- Crawford, J.G. (1915) Descriptions of new Hymenoptera, No 9. *Proceedings of the United States National Museum*, 48, 577–586.
- Cresson, E.T. (1865) On the Hymenoptera of Cuba. *Proceedings of the Entomological Society of Philadelphia*, 4, 1–200.
- Cresson, E.T. (1869a) Descriptions of North American bees. I. *Proceedings of Boston Society of Natural History*, 12, 165–171.
- Cresson, E.T. (1869b) Notes on Cuban Hymenoptera, with descriptions of new species. *Transactions of the American Entomological Society*, 2, 293–298.
- Cresson, E.T. (1916) The Cresson types on Hymenoptera. *Memoirs of the American Entomological Society*, 1, 1–141.
- Cross, E.A. (1958) A revision of the bees of the subgenus *Epinomia* in the New World (Hymenoptera, Halictidae). *University of Kansas Science Bulletin*, 38, 1261–130.
- Cuvier, G.L. C.F.D. (1849) *Le règne animal distribué d'après son organisation*. Disciples ed., Paris, Masson. Insectes 2, 1–443.
- Dalla Torre, C.G. de (1896) *Catalogus Hymenopterorum* Vol. X, Apidae (Anthophilia). Leipzig, Engelmann, 643 pp.
- Danks, H.V. (1991) Museums collections: fundamental values and modern problems. *Collection Forum* 7, 95–111.
- Díaz, I. & Vale, A. (2001) Actividad polinizadora y aspectos conductuales de la abeja *Xylocopa cubaecola* Lucas (Hymenoptera: Apoidea) en condiciones naturales. *Fitosanidad*, 5, 25–30.
- Díaz, M.E. (1985) Contribución al estudio de la historia de la apicultura en Cuba. *Ciencia y Técnica Agrícola, Serie Apicultura*, 1, 109–116.
- Díaz, M.E. & Domínguez, D.A. (1985) Características morfológicas de la abeja (*Apis mellifica*) en Cuba. *Ciencia y Técnica Agrícola, Serie Apicultura*, 1, 31–105.
- Eickwort, G.C. (1969) A comparative morphological study and generic revision of the Augochlorine bees (Hymenoptera: Halictidae). *The University of Kansas Science Bulletin* 48, 325–524.
- Eickwort, G.C. (1988) Distribution patterns and biology of West Indian sweat bees (Hymenoptera: Halictidae). In: Liebherr, J.K. (Editor), *Zoogeography of Caribbean insects*. Cornell University Press, New York, pp. 231–253.
- Eickwort, G.C. & Ginsberg, H. S. (1980) Foraging and mating behavior in Apoidea. *Annual Review of Entomology*, 25, 421–446.
- Engel, M.S. (1995) *Neocorynura electra*, a new fossil bee species from Dominican amber (Hymenoptera: Halictidae). *Journal of New York Entomological Society*, 103, 317–323.
- Engel, M.S. (1999) The first fossil *Euglossa* and phylogeny of the orchid bees (Hymenoptera: Apidae; Euglossini). *American Museum Novitates*, 3272, 1–14.
- Engel, M.S. (2001b) Monophyly and extensive extinction of advanced eusocial bees: insights from an unexpected Eocene diversity. *Proceedings of the National Academy of Sciences of the United States of America*, 98, 1661–1664.
- Engel, M.S. (2004). On the identity of *Halictus cubensis* Spinola, 1851 (Hymenoptera: Halictidae). *Entomological News*, 115, 169–170.
- Engel, M. S. (In press a) The *Sphecodes* of Cuba (Hymenoptera: Halictidae). *Acta Zoologica Cracoviensis*.
- Engel, M. S. (In press b) A new genus of cleptoparasitic bees from the West Indies (Hymenoptera: Halictidae). *Acta Zoologica Cracoviensis*.
- Essig, E.O. (1931) *A history of entomology*. The MacMillan Company, New York, 1029 pp.
- Fabricius, J.C. (1793) *Entomologia Systematica Emendata et Aucta*. Vol. 2. Hafniae: Proft, 519 pp.
- Fabricius, J.C. (1798) *Entomologia Systematica. Emendata et Aucta*. Supplement (p. 272).
- Fabricius, J.C. (1804) *Systema piezatorum*. Brunsvigae: Reichard, 440 pp.

- Fernández, J.L., Sariol, H., Vega, M. A., Ricardo, S., González, M., Portuondo F., E. (2002) Datos preliminares sobre la biodiversidad del orden Hymenoptera en la provincia Granma, Cuba. *Boletín de la Sociedad Entomológica Aragonesa, España*, 31, 43–48.
- Finnamore, A.T. & Michener, C.D. (1993) Superfamily Apoidea. In: Goulet, H. and Huber, J. T. (Eds.), *Hymenoptera of the world: an identification guide to families*. Canada Communication Group Publishing, Ottawa, pp 279–325.
- Flakus, G.T. (1993) *Living with killer bees. The story of the africanized bee invasion*. Quick Trading Comp., California, 144 pp.
- Frias, A.I., Rivera, F., Ferrer, E.M. & Vidal, A. (2000) Toxicidad aguda del veneno de *Apis mellifera* en roedores. *Revista de Biología, University of Havana*, 14, 46–50.
- Friese, H. (1899) Monographie des bienengattungen *Exomalopsis*, *Ptilothrix*, *Melitoma* und *Tetrapedia*. *Annalen des K. K. Naturhistorischen Hofmuseums, Bd*, 14, 247–304.
- Friese, H. (1900) Monographie der bienengattung *Centris* (s. lat.). *Annalen des KK Naturhistorischen Hofmuseums [Wien]*, 15, 237–350.
- Friese, H. (1902) Beitrag zur Apidenfauna der Grossen Antillen. *Zeitschrift für Systematische Hymenopterologie und Dipterologie*, 2, 196–201.
- Friese, H. (1908) Beitrag zur bienenfauna der Kleinen Antillen und der Bermudas. *Zoologische Jahrbücher Suppl.*, 11, 33–40.
- Friese, H. (1911) *Apidae I. Megachilinae*. Das Tierreich 28. R. Friedlander und Sohn, Berlin, 440 pp.
- Friese, H. (1921) Über die kegelbienen (*Coelioxys*) brasiliens. *Zoologische Jahrbücher*, 44, 421–485.
- García, I. et al. (1973) Informe del trabajo faunístico realizado en el Plan Jibacoa-Cayajabos. *Serie Biológica, Instituto de Zoología*, 43, 1–25.
- Genaro, J.A. (1994a) Adiciones a la lista de himenópteros cubanos (Pompilidae, Megachilidae). *Cocuyo*, 1, 6.
- Genaro, J.A. (1994b) Inquilinos de *Sceliphrom assimile* con énfasis en *Podium fulvipes* (Hymenoptera: Vespidae, Sphecidae, Megachilidae). *Caribbean Journal of Science*, 30, 268–270.
- Genaro, J.A. (1996a) Plantas usadas por abejas del género *Megachile* para construir las celdillas de sus nidos (Hymenoptera: Megachilidae). *Caribbean Journal of Science*, 32, 365–368.
- Genaro, J.A. (1996b) Nest parasites (Coleoptera, Diptera, Hymenoptera) of some wasps and bees (Vespidae, Sphecidae, Colletidae, Megachilidae, Anthophoridae) in Cuba. *Caribbean Journal of Science*, 32, 239–240.
- Genaro, J.A. (1996c) Resumen del conocimiento sobre los meloidos de Cuba (Insecta: Coleoptera). *Caribbean Journal of Science*, 32, 382–386.
- Genaro, J.A. (1996d) Foresis de *Caenocara* sp. (Coleoptera: Anobiidae) en *Anthophora atrata* (Hymenoptera: Anthophoridae). *Melissa*, 9, 9.
- Genaro, J.A. (1997) Key to the genus *Megachile*, *Chalicodoma* group (Hymenoptera: Megachilidae). *Revista de Biología Tropical*, 44/45, 193–198.
- Genaro, J.A. (1998a) Distribution and synonymy of some Caribbean bees of the genera *Megachile* and *Coelioxys* (Hymenoptera: Megachilidae). *Caribbean Journal of Science*, 34, 151–152.
- Genaro, J.A. (1998b) El género *Ceratina* en Cuba y La Española (Hymenoptera: Apidae). *Caribbean Journal of Science*, 34, 95–98.
- Genaro, J.A. (1998c) Hoja de datos de taxon para *Megachile armaticeps*. In Pérez, E., Osa, E., Matamoros, Y., Shillcox, J. & Seal, U. S. (Editors), *Conservation breeding specialist group (SSC/UICN). Report of conservation assessment and management plan workshop for selected Cuban species*. CBSG, Apple Valley, Minnesota, pp: 181–189.
- Genaro, J.A. (1999) Revisión del género *Triepeolus* en Cuba (Hymenoptera: Apidae), con descripción de dos especies nuevas. *Caribbean Journal of Science*, 35, 215–220.

- Genaro, J.A. (2001a) Tres especies nuevas del género *Lasioglossum* (*Dialictus*), grupo *Habralictellus* para Cuba (Hymenoptera: Halictidae). *Solenodon*, 1, 38–44.
- Genaro, J.A. (2001b) Especie nueva de *Melissoptila* para Cuba y La Española (Hymenoptera: Apidae). *Avicennia*, 14, 61–64.
- Genaro, J.A. (2001c) Especies nuevas de abejas de Cuba y la Española (Hymenoptera: Colletidae, Megachilidae, Apidae). *Revista de Biología Tropical*, 49, 951–959.
- Genaro, J. (2002) Taxonomía, diversidad y distribución de las abejas de Cuba (Insecta: Hymenoptera) Ph.D. Thesis, University of Havana, Havana city, Cuba.
- Genaro, J.A. (2003) El género *Colletes* en Cuba (Hymenoptera: Apiformes: Colletidae). *Solenodon*, 3, 53–56.
- Genaro, J.A. (2004) Las abejas de la Isla de la Juventud, Cuba (Hymenoptera: Apoidea). *Boletín de la Sociedad Entomológica Aragonesa, Spain*, 34, 177–179.
- Genaro, J.A. & Sánchez, C. S. (1992) Notas sobre la conducta de nidificación de *Philanthus banana-bacoa* Alayo (Hymenoptera: Sphecidae). *Caribbean Journal of Science*, 28, 44–46.
- Genaro, J.A. & Tejuca, A. E. (2001) Patterns of endemism and biogeography of Cuban insects. In: Woods, C. A. and Sergile, F. E. (Editors.), *Biogeography of the West Indies. Patterns and perspectives*. CRC Press, Boca Ratón, Florida, pp: 77–83.
- Goulet, H. & Huber, J.T. (Editors) (1993) *Hymenoptera of the world: An identification guide to families*. Centre for Land and Biological Resources Research Ottawa, Ontario, 668 pp.
- Goulson, D. (2003) Effects of introduced bees on native ecosystems. *Annual Review of Ecology, Evolution and Systematics*, 34, 1–26.
- Grimaldi, D.A. & Engel, M.S. (2005) *Evolution of the insects*. Cambridge University Press, New York. 755 pp.
- Guérin-Méneville, F.E. (1835) *Iconographie du regne animal de G. Cuvier, on représentation d'après nature de l'une des especes les plus remarquables et souvent non encore figurées de chaque genre d'animaux, pouvant servir d'atlas a tous les traités de zoologie*. Vol. 2. Plates. Fain and Thunot, Paris.
- Guérin-Méneville, F.E. (1844) *Iconographie du regne animal de G. Cuvier, on représentation d'après nature de l'une des especes les plus remarquables et souvent non encore figurées de chaque genre d'animaux, pouvant servir d'atlas a tous les traités de zoologie*. Vol. 3 (text). Insects. Fain and Thunot, Paris, 576 pp.
- Gundlach, J. (1886) *Contribución a la entomología cubana*. Tomo II. Hymenópteros. Imprenta La Antilla, Havana, 187 pp.
- Hernández, J.C. & Lemus, Y. (1996a) Comportamiento de la abeja (*Apis mellifera*) como polinizador en el cultivo del melón. In: *Simposio de Flora Melífera y Polinización, Resumen. Facultad Farmacia y Alimentos*. Havana city.
- Hernández, J.C. & Lemus, Y. (1996b) Comportamiento del rendimiento en calabaza utilizando a la abeja (*Apis mellifera*) como insecto polinizador. In: *Simposio de Flora Melífera y Polinización, Resúmenes. Facultad Farmacia y Alimentos*. Havana city.
- Hogue, C.L. (1993) *Latin American insects and Entomology*. University of California Press, Berkeley, 536 pp.
- Hurd, Jr., P.D. (1978) *An annotated catalog of the carpenter bees (genus Xylocopa Latreille) of the Western Hemisphere (Hymenoptera: Anthophoridae)*. Smithsonian Institution Press, Washington, D. C., 106 pp.
- Iturralde-Vinent, M.A. (2001) Geology of the amber-bearing deposits of the Greater Antilles. *Caribbean Journal of Science*, 37, 141–167.
- Iturralde-Vinent, M.A. & MacPhee, R.D.E. (1999) Paleogeography of the Caribbean Region: implications for Cenozoic biogeography. *Bulletin of the American Museum of Natural History*, 238, 1–95.
- Janjic, J. & Packer, L. (2003) Phylogeny of the bee genus *Agapostemon* (Hymenoptera: Halictidae).

- Systematic Entomology*, 28, 101–123.
- Kinsey, L.S. & Dressler, R.L. (1986) Synonymic species list of Euglossini. *Pan-Pacific Entomologist*, 62, 229–236.
- Kremen, C., Williams, N.M., Bugg, R.L., Fay J.P. & Thorp, R.W. (2004) The area requirements of an ecosystem service: crop pollination by native bee communities in California. *Ecology Letters*, 7, 1109–1119.
- Krombein, K.V., Hurd, Jr., P.D., Smith, D.R. & Burks, B.D. (eds). (1979) *Catalog of Hymenoptera in America North of Mexico*. 2, 1199–2209. Smithsonian Institution Press, Washington, D.C.
- LaBerge, W.E. (1956) A revision of the bees of the genus *Melissodes* in North and Central America. Part I. (Hymenoptera, Apidae). *University of Kansas Science Bulletin*, 37, 911–1194.
- LaBerge, W.E. (1961) A revision of the bees of the genus *Melissodes* in North and Central America. Part III (Hymenoptera, Apidae). *University of Kansas Science Bulletin*, 42, 283–663.
- León, A.A. (1997) Polinización con abejas. *Se puede*, 10, 16–19.
- León, A.A. & Rivero, L. (1999) Polinización de melón (*Cucumis melo* L.) en casas de cultivo, con *Apis mellifera*. *Apiciencias* (electronic journal, Havana city) 1, 10–15.
- León, A.A. & Sánchez, U. (1998) Feeding and nesting plants of *Xylocopa cubaecola* (Hymenoptera: Apidae). *Caribbean Journal of Science*, 34, 152–155.
- Lepeletier, A. (1841) *Histoire naturelle des insectes. Hymenopteres*. Tome 2. Pp. 262–289.
- Llamacho, J.A. & Larramendi, J.A. (2005). *The orchids of Cuba*. Greta editors, Sevilla, Spain, 287 pp.
- McGinley, R.J. (1992) Where's the management in collections management? Planning for improved care, greater use, and growth of collections. In: Palacios, F., Martínez, C. & Thomas, B. (Editors.) *The International Symposium and First World Congress on the Preservation and Conservation of Natural History Collections, held in Madrid, 10–15 May 1992. Congress Book. Volume 3. Current Issues, Initiatives, and Future Directions for the Preservation and Conservation of Natural History Collections*. pp. 309–338.
- Michener, C.D. (1944) Comparative external morphology, phylogeny, and a classification of the bees (Hymenoptera). *Bulletin of the American Museum of Natural History*, 82, 151–326.
- Michener, C.D. (1953) Comparative morphological and systematic studies of bee larvae with a key to the families of hymenopterous larvae. *University of Kansas Science Bulletin*, 35, 987–1102.
- Michener, C.D. (1966) The classification of the Diphaglossinae and North American species of the genus *Caupolicana* (Hymenoptera, Colletidae). *University of Kansas Science Bulletin*, 46, 717–751.
- Michener, C.D. (1974) The social behavior of the bees. A comparative study. Harvard Univ. Press, Cambridge, 404 pp.
- Michener, C.D. (1979) Biogeography of the bees. *Annals of the Missouri Botanical Garden*, 66, 277–347.
- Michener, C.D. (1982) The africanized honey bee. In: Jaison, P. (Editor). *Social insects in the tropics*. University Paris-Nord, pp: 205–207.
- Michener, C.D. (1986) Family-group names among bees. *Journal Kansas of the Entomological Society*, 59, 219–234.
- Michener, C.D. (1988a) The parasitic anthophorid genus *Xeromelecta* in Cuba (Hymenoptera: Apoidea). *Annals of the Entomological Society of America*, 81, 377–379.
- Michener, C.D. (1988b) The genus *Lithurge* in the Antilles (Hymenoptera: Megachilidae). *Folia Entomológica Mexicana*, 76, 159–164.
- Michener, C.D. (1990) Reproduction and castes in social halictine bees. In: Engels, W. (Editor). *Social insects. An evolutionary approach to castes and reproduction*. Berlin: Springer-Verlag, pp. 77–121.
- Michener, C.D. (2000) *The bees of the world*. Johns Hopkins Univ. Press, Baltimore, 913 pp.
- Michener, C.D. & A. Fraser (1978) A comparative study of mandibular structure in bees. *University*

- of *Kansas Science Bulletin*, 51, 463–482.
- Michener, C.D., McGinley R.J. & Danforth, B.N. (1994) *The bee genera of North and Central America (Hymenoptera: Apoidea)*. Smithsonian Institution Press, Washington, D. C, 209 pp.
- Mitchell, T.B. (1927) New West Indian *Megachile*. *Psyche*, 34, 47–57.
- Mitchell, T.B. (1943) On the classification of Neotropical *Megachile* (Hymenoptera: Megachilidae). *Annals of the Entomological Society of America*, 35, 656–671.
- Mitchell, T.B. (1960) Bees of the Eastern United States. I. *North Carolina Agricultural Experimental Station Technical Bulletin*, 141: 1–538.
- Mitchell, T.B. (1962) Bees of the Eastern United States. II. *North Carolina Agricultural Experimental Station Technical Bulletin*, 152, 1–557.
- Mitchell, T.B. (1973) A subgeneric revision of the bees of the genus *Coelioxys* of the Western Hemisphere. *Contributions of the Department of Entomology of North Carolina State University*, 124 pp.
- Mitchell, T.B. (1980) A generic revision of the Megachiline bees of the Western Hemisphere. *Contributions of the Department of Entomology of North Carolina State University*, 95 pp.
- Moure, J. (1960) Notes on the types of Neotropical bees described by Fabricius (Hymenoptera: Apoidea). *Studia Entomologica, Peotrópolis*, 3, 97–160.
- Moure, J. (1967) A check-list of the known Euglossine bees (Hymenoptera, Apidae). *Atas Simpósio sobre a Biota Amazônica*, 5, 395–415.
- Moure, J.S. & Hurd, P.D. (1987) *An annotated catalog of the halictid bees of the Western Hemisphere (Hymenoptera:Halictidae)*. Smithsonian Institution Press, Washington, D.C, 405 pp.
- Moure J.S. & Urban, D. (2002) Catálogo de Apoidea de Región Neotropical (Hymenoptera, Colletidae). III. Colletini. *Revista Brasileira de Zoología*, 19, 1–30.
- O'Toole, C. & Raw, A. (1999) *Bees of the World*. Bandford, London, 192 pp.
- Poey, F. (1851) Historia de la abeja de la tierra. *Trigona fulvipes*. *Memorias sobre la Historia Natural de la Isla de Cuba*, 1, 122–176.
- Poey, F. (1856–1858) Apéndice sobre la abeja de la tierra. *Memorias sobre la Historia Natural de la Isla de Cuba*, 2, 72.
- Poey, F. (1888a) La abeja (*Apis mellifica*, Linn.). *Memorias sobre la Historia Natural de la Isla de Cuba*, 3, 71–72.
- Poey, F. (1888b) Historia de la abejas en la formación de una colonia. *Memorias sobre la Historia Natural de la Isla de Cuba*, 3, 72–74.
- Portuondo, E., & Fernández, J.L. (2004) Biodiversidad del orden Hymenoptera en los macizos montañosos de Cuba Oriental. *Boletín de la Sociedad Entomológica Aragonesa, Spain*, 35:121–136.
- Proctor, M. & Yeo, P. (1973) *The pollination of flowers*. Collins, Londres, 418 pp.
- Pruna, P.M. (2001) *Ciencia y científicos en Cuba Colonial. La Real Academia de Ciencias de La Habana. 1861–1898*. Editorial Academia, Havana city, 243 pp.
- Raw, A. (2002) New combinations and synonymies of leafcutter and mason bees of the Americas (*Megachile*, Hymenoptera, Megachilidae). *Zootaxa*, 71, 1–43.
- Raw, A. (2004) Nomenclatural changes in leafcutter bees of the Americas: *Megachile* Latreille 1802 (Hymenoptera; Megachilidae). *Zootaxa*, 766, 1–4.
- Ricketts, T.H., Daily, G.C., Ehrlich, P.R. & Michener C.D. (2004) Economic value of tropical forest to coffee production. *Proceedings of the National Academy of Sciences of the United States of America* 101, 12579–12582.
- Robertson, C. (1902) Synopsis of Halictinae. *The Canadian Entomologist*, 34, 243–250.
- Roberts, R.B. (1972) Revision of the bee genus *Agapostemon* (Hymenoptera: Halictidae). *University of Kansas Science Bulletin*, 49, 437–590.
- Rodríguez, Y., Rivera, A. I., Frasi, F., Vidal, A. & Bellón, Y. (1995) Evaluación analgésica y toxicológica del veneno de abeja *Apis mellifera*. *Revista Cubana de Farmacia*, 30, 145–152.

- Rohwer, S.A. & Holland, W.J. (1917) List of the Hymenoptera collected on the Isle of Pines by G. A. Link, Sr., 1912–1913, and contained in the Carnegie Museum. *Annals of the Carnegie Museum*, 11, 291–296.
- Romay, T. (1796) *Discurso sobre los obstáculos que han impedido progresen las colmenas en la Isla de Cuba*. Imprenta de la Capitanía General, Havana, 34 pp.
- Roubik, D.W. (1989) *Ecology and natural history of tropical bees*. Cambridge University Press, Cambridge, 514 pp.
- Roubik, D.W. & Ackerman, D. (1987) Long-term ecology of euglossine orchid-bees (Apidae: Euglossini) in Panama. *Oecologia*, 73, 321–333.
- Sagra, R. de la. (1856) (Compiler) *Historia física, política y natural de la isla de Cuba. Tomo VII. Crustáceos, arácnidos e insectos*. Librería A. Bertrand, París, 371 pp.
- Sakagami, S.F. & Michener, C.D. (1987) Tribes of Xylocopinae and origin of the Apidae (Hymenoptera: Apoidea). *Annals of the Entomological Society of America*, 80, 439–450.
- Sandhouse, G.A. (1923) The bee-genus *Dialictus*. *The Canadian Entomologist*, 55, 193–195.
- Scaramuzza, L.C. (1938) Breve nota acerca de dos parásitos de "*Megachile* sp." (Hymenoptera, Apoidea, Megachilidae). *Memorias de la Sociedad Cubana de Historia Natural Felipe Poey*, 12, 87–88.
- Schwarz, H.F. (1932) The genus *Melipona*. The type genus of the Meliponidae or stingless bees. *Bulletin of the American Museum of Natural History*, 63, 231–460.
- Schwarz, H.F. (1944) Stingless bees of the Western Hemisphere. *Bulletin of the American Museum of Natural History*, 90, 1–546.
- Scopoli, I.A. (1763) *Entomologia Carniolica exhibens Insecta Carnioliae indigena et distributa in ordines, genera, species, varietates methodo Linnaeana*. Vindobonae; Trattner, 420 pp.
- Smith, F. (1853). *Catalogue of hymenopterous insects in the collection of the British Museum. Part. I. Andrenidae and Apidae*. London, 197 pp.
- Smith, F. (1879) *Descriptions of new species of Hymenoptera in the collection of the British Museum*. London, 240 pp.
- Smith, H.S. (1907) Contribution toward a monograph of the Ceratinidae of North and Middle America. *Transactions of the American Entomological Society*, 33, 115–124.
- Snelling, R.R. (1984) Studies on the taxonomy and distribution of American Centridine bees (Hymenoptera: Anthophoridae). *Contributions in Science, Natural History Museum of Los Angeles County*, 347, 1–69.
- Snelling, R.R. (1985) A review of the genera of cleptoparasitic bees of the tribe Ericrocini (Hymenoptera: Anthophoridae). *Contributions in Science, Natural History Museum of Los Angeles County*, 369, 1–34.
- Snelling, R.R. (1986) Contributions toward a revision of the New World Nomadine bees. A partitioning of the genus *Nomada* (Hymenoptera: Anthophoridae). *Contributions in Science, Natural History Museum of Los Angeles County*, 376, 1–32.
- Snelling, R.R. (1990) A review of the native North American bees of the genus *Chalicodoma* (Hymenoptera: Megachilidae). *Contributions in Science, Natural History Museum of Los Angeles County*, 421, 1–39.
- Sotolongo, L.O., Machado S.T. & León, A.A. (1997) Diversidad de polen en masas de aprovisionamiento de *Xylocopa cubaecola* (Hymenoptera: Apoidea). *Cocuyo*, 6, 11–12.
- Spinola, M. (1840) Observations sur les apiaires méliponides. *Annales des Sciences Naturelle, série 2, Zoologie* 13: 116–140, pl. 2.
- Spinola, M. (1851) *Hymenópteros*. Pp. 153–569. In C. Gay (Ed.) *Historia física y política de Chile, Zoología* [volumen 6]. Maulde et Renou. Paris, France, 596+[2] pp.
- Stubbs, C.S. & Drummond, F.A. (eds). (2001) Bees and crop pollination-crisis, crossroads, conservation. *Proceedings Thomas Say Publications in Entomology*, 156 pp.
- Taylor, O.R., Jr. (1985) African bees: potential impact in the United States. *Bulletin of the Entomo-*

- logical Society of America* 31, 15–24.
- Timberlake, P.H. (1980) Review of North American *Exomalopsis* (Hymenoptera, Anthophoridae). *University of California Publications in Entomology*, 86, 1–158.
- Urban, D. (1970) As espécies do género *Florilegus* Robertson, 1900 (Hymenoptera, Apoidea). *Boletim Universidade Federal do Parana, Zoologia* 3, 245–280.
- Urban D. & Moure, J.S. (2001) Catálogo de Apoidea de Região Neotropical (Hymenoptera, Colletidae). II. Diphaglossinae. *Revista Brasileira de Zoologia*, 18, 1–34.
- Urban D. & Moure, J.S. (2002) Catálogo de Apoidea de Região Neotropical (Hymenoptera, Colletidae). IV. Hylaeinae. *Revista Brasileira de Zoologia*, 19, 31–56.
- Villanueva-G, R, Roubik, D.W. & Colli-Ucán, W. (2005) Extinction of *Melipona beecheii* and traditional beekeeping in the Yucatán peninsula. *Bee World*, 86, 35–41.
- Wcislo, W.T. (1997) Behavioral environments of sweat bees (Halictinae) in relation to variability in social organization. In: Choe, J.C. & Crespi, B.J. (Editors) *Social behavior in insects and arachnids*. Cambridge University Press, United Kingdom, pp: 316–332.
- Werthein, I. (1993) *Cuba, paraíso de las abejas*. Editorial Oriente, Santiago de Cuba, 174 pp.
- Zayas, F. de (1981) *Entomofauna cubana. Sección Oligoneoptera. Ordenes Hymenoptera y Strepsiptera*. Tomo VIII. Editorial Científico-Técnica, Havana city, 111 pp.