



***Intellagama lesueurii* (Gray, 1831), the correct binomial combination for the Australian Eastern Water Dragon (Sauria, Agamidae)**

ANDREW P. AMEY^{1,3}, PATRICK J. COUPER¹ & GLENN M. SHEA²

¹Biodiversity, Queensland Museum, PO Box 3300, South Brisbane 4101, Australia

²Faculty of Veterinary Science B01, University of Sydney 2006 NSW, Australia

³Corresponding author. E-mail: andrew.amey@qm.qld.gov.au

The Eastern Water Dragon is a large, conspicuous agamid, well known to many inhabitants of eastern Australia. It was first described in the scientific literature as *Lophura lesueurii* by Gray (1831). Gray's allocation of this taxon to his earlier genus *Lophura* Gray, 1827, created for the species *Lacerta lophura* Shaw, 1802, does not mention that Cuvier (1829) had erected *Istiurus* for *amboinensis* Schlosser, 1768, which Cuvier treated as a senior synonym of *lophura*. Cuvier considered the generic name *Lophura* to be too similar to *Lophyrus* Latreille, 1802, a genus of conifer sawflies belonging to the family Diprionidae, hence the need for a new genus. Both Gray and Cuvier were evidently unaware that *Lophura* was unavailable as this name had already been assigned to a genus of phasianid birds (Fleming 1822). The Eastern Water Dragon has subsequently appeared in the taxonomic literature under the following synonyms: *Iguana paramatensis* Fitzinger, 1843; *Amphibolurus maculiferus* Girard, 1857; *Amphibolurus heterurus* Peters, 1866 and *Amphibolurus branchialis* De Vis, 1884. The combination *Physignathus lesueurii* was first used in 1845 by Gray in his *Catalogue of the Specimens of Lizards in the British Museum* and has been in use ever since. The only other generic name proposed for this taxon is *Intellagama* Wells and Wellington, 1985, by which the authors implied its distinctiveness from the other member of *Physignathus*, the Chinese Water Dragon *P. cocincinus* Cuvier, 1829. However, as their description provided no evidence to demonstrate that Australian water dragons are generically distinct from their foreign congener, this name has not been adopted by subsequent authors, and has been informally treated as a synonym of *Physignathus*.

In a recent paper by Townsend *et al.* (2011), evidence was presented showing that the genus *Physignathus* is not monophyletic, a conclusion supported by earlier molecular studies (Hugall *et al.* 2008; Macey *et al.* 2000; Schulte *et al.* 2003). These authors analysed DNA sequence data from 29 nuclear protein-coding genes for 47 iguanian taxa. They concluded that "...the southeast Asian species *P. cocincinus* (Cuvier, 1829) is the sister taxon to the entire Australian agamid radiation, which includes its sole congener, the Australian species *P. lesueurii*...". To rectify this paraphyly, they resurrected the genus *Istiurus*, to which Duméril and Bibron (1837) had assigned *cocincinus*, *lesueurii* and *amboinensis*, stating that the name is "not currently used and therefore is available". Regrettably, these authors overlooked the fact that Duméril and Bibron (1837) did not erect that generic name, which was instead created by Cuvier (1829). In his original description, Cuvier assigned only a single species, *Lacerta amboinensis* to *Istiurus* and thus *amboinensis* is the type species of this genus by monotypy. The resurrection by Poche (1903) of *Hydrosaurus* Kaup, 1828 as the earliest available generic name for *amboinensis* places *Istiurus* in the synonymy of this genus. In light of this, *Intellagama* Wells and Wellington 1985 is the earliest generic name available for *lesueurii* and *Intellagama lesueurii* (Gray, 1831) is the appropriate binomial combination.

Georges and Thomson (2010) have questioned the validity of Wells' more recent taxonomic contributions, (specifically Wells 2007a, b, c, 2009) on the grounds that they violate ICZN Articles 8 and 9 and Recommendation 8D. However, these arguments cannot be extended to the 1985 Wells and Wellington publication in which *Intellagama* was erected. As this was printed and distributed as multiple copies, it meets the minimum requirements of the Code (ICZN, 1999). The ICZN acknowledged this in 1991 when it refused to vote on a proposal to suppress the works of Wells and Wellington published up to that date, despite the "clear rejection by Wells and Wellington of virtually every tenet of the voluntary Code of Ethics which forms Appendix A of the Code" (ICZN 1991). Since this time, many of the generic names proposed by Wells and Wellington (1983, 1985) have gained widespread acceptance and now appear in many scientific and popular publications (for example, Bauer *et al.* 1997; Cogger 1996, 2000; Couper *et al.* 2006; Dolman & Hugall 2008; Donnellan *et al.* 2000; Gardner *et al.* 2008; Greer 1989, 1997; Greer & Cogger 1985; Kluge 1993;

Queensland Museum 2000, 2007; Sadlier *et al.* 1993; Wilson 2005; Wilson & Swan 2003, 2008, 2010). Now that appropriate evidence has been presented to demonstrate that *lesueurii* is generically distinct from *cocincinus* (Townsend *et al.* 2011), *Intelligama* should be regarded as the only available generic name for this taxon.

References

- Bauer, A.M., Good, D.A. & Branch, W.R. (1997) The taxonomy of the southern African leaf-toed geckos (Squamata: Gekkonidae), with a review of Old World "*Phyllodactylus*" and the description of five new genera. *Proceedings of the California Academy of Sciences*, 49, 447–497.
- Cogger, H.G. (1996) *Reptiles and Amphibians of Australia* (5th ed.). Chatswood: Reed Books.
- Cogger, H.G. (2000) *Reptiles and Amphibians of Australia* (6th ed.). Sydney: Reed New Holland.
- Couper, P.J., Covacevich, J.A., Amey, A.P. & Baker, A.M. (2006) The genera of skinks (family Scincidae) of Australia and its island territories: diversity, distribution and identification. In: J. R. Merrick, M. Archer, G. M. Hickey & M. S. Y. Lee (Eds), *Evolution and Biogeography of Australasian Vertebrates*. Oatlands: Auscipub, pp. 367–383.
- Cuvier, G. (1829) *Le Règne Animal distribué d'après son organisation, pour servir de base à l'histoire naturelle des animaux et d'introduction à l'anatomie comparée. Nouvelle édition* (Vol. 2). Paris: Déterville.
- De Vis, C.W. (1884) On a new Australian lizard. *Proceedings of the Royal Society of Queensland*, 1, 53–56.
- Dolman, G. & Hugall, A. (2008) Combined mitochondrial and nuclear data enhance resolution of a rapid radiation of Australian rainbow skinks (Scincidae: *Carlia*). *Molecular Phylogenetics and Evolution*, 49, 782–794.
- Donnellan, S.C., Aplin, K.P. & Dempsey, P.J. (2000) Generic and morphological variation in Australian *Christinus* (Squamata: Gekkonidae): preliminary overview with recognition of a cryptic species on the Nullarbor Plain. *Australian Journal of Zoology*, 48, 289–315.
- Duméril, A.M.C. & Bibron, G. (1837) *Erpétologie Générale, ou Histoire Naturelle Complète des Reptiles* (Vol. 4). Paris: Roret.
- Fitzinger, L. (1843) *Systema Reptilium, Fasciculus Primus Amblyglossae*. Vienna: Braumüller et Seidel.
- Fleming, J. (1822) *The philosophy of zoology; or a general view of the structure, functions, and classification of animals*. Edinburgh: Constable.
- Gardner, M.G., Hugall, A., Donnellan, S.C., Hutchinson, M.N. & Foster, R. (2008) Molecular systematics of social skinks: phylogeny and taxonomy of the *Egernia* group (Reptilia: Scincidae). *Zoological Journal of the Linnean Society*, 154, 781–794.
- Georges, A. & Thomson, S. (2010) Diversity of Australasian freshwater turtles, with an annotated synonymy and keys to species. *Zootaxa*, 2496, 1–37.
- Girard, C. (1857) Descriptions of some new reptiles, collected by the United States Exploring Expedition, under the command of Capt. Charles Wilkes, U.S.N. Fourth Part: including the species of saurians exotic to North America. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 1857, 195–199.
- Gray, J.E. (1827) A synopsis of the genera of Saurian Reptiles, in which some new genera are indicated, and the others reviewed by actual examination. *Annals of Philosophy, new series*, 2, 54–58.
- Gray, J.E. (1831) A synopsis of the species of the class Reptilia. In: E. Griffith & E. Pidgeon (Eds), *The Animal Kingdom Arranged in Conformity with its Organization by the Baron Cuvier, member of the Insititute of France, &c. &c. &c. with additional descriptions of all the species hitherto named, and of many not before noticed. Volume 9. The class Reptiles arranged by the Baron Cuvier, with specific descriptions*. London: Whittaker, Treacher & Co., p. 481+110.
- Gray, J.E. (1845) *Catalogue of the Specimens of Lizards in the Collection of the British Museum* (Vol. 28). London: British Museum.
- Greer, A.E. (1989) *The Biology and Evolution of Australian Lizards*. Sydney: Surrey Beatty & Sons.
- Greer, A.E. (1997) *The Biology and Evolution of Australian Snakes*. Chipping Norton: Surrey Beatty & Sons.
- Greer, A.E. & Cogger, H. (1985) Systematics of the reduce-limbed and limbless skinks currently assigned to the genus *Anomalopus* (Lacertilia: Scincidae). *Records of the Australian Museum*, 37, 11–54.
- Hugall, A., Foster, R., Hutchinson, M. & Lee, M.S.Y. (2008) Phylogeny of Australasian agamid lizards based on nuclear and mitochondrial genes: implications for morphological evolution and evolution and biogeography. *Biological Journal of the Linnean Society*, 93, 343–358.
- ICZN (1991) Three works by Richard W. Wells and C. Ross Wellington: proposed suppression for nomenclatural purposes. *Bulletin of Zoological Nomenclature*, 48, 337–338.
- ICZN (1999) *International Code of Zoological Nomenclature* (4th ed.). London: International Trust for Zoological Nomenclature.
- Kaup, J.J. (1827) Ueber *Hyaena*, *Uromastix*, *Basiliscus*, *Corythacolus*, *Acontias*. *Isis von Oken*, 21, 1144–1150.
- Kluge, A.G. (1993) *Aspidites* and the phylogeny of pythonine snakes. *Records of the Australian Museum, Supplement*, 19, 1–77.

- Latreille, P.A. (1802) *Histoire Naturelle des Fourmis, et Recueil de Mémoires et d'Observations sur les Abeilles, les Araignées, les Faucheurs, et Autres Insectes*. Paris: Barrois.
- Macey, J.R., Schulte, J.A., Larson, A., Ananjeva, N.B., Wang, Y.-Z., Pethiyagoda, R., Rastegar-Pouyani, N. & Papenfuss, T.J. (2000) Evaluating trans-Tethys migration: an example using acrodont lizard phylogenetics. *Systematic Biology*, 49, 233–256.
- Peters, W. (1866) Eine mittheilung über neue Amphibien (*Amphibolurus*, *Lygosoma*, *Cyclodus*, *Masticophis*, *Crotaphopeltis*) und Fische (*Diagramma: Haplagenys*) des Kgl. zoologischen museums. *Monatsberichte der Königlich Akadamie der Wissenschaften zu Berlin*, 1866, 86–96.
- Poche, F. (1903) Einige nothwendige Änderungen in der herpetologischen nomenklatur. *Zoologischer Anzeiger*, 26, 698–703.
- Queensland Museum (2000) *Wildlife of Tropical North Queensland*. South Brisbane: Queensland Museum.
- Queensland Museum (2007) *Wildlife of Greater Brisbane* (2nd ed.). South Brisbane: Queensland Museum.
- Sadlier, R.A., Colgan, D.J. & Shea, G.M. (1993) Taxonomy and distribution of the scincid lizard *Saproscincus challengeri* and related species in southeastern Australia. *Memoirs of the Queensland Museum*, 34, 139–158.
- Schlosser, J.A. (1768) *Brief van Johannes Albertus Schlosser (...) aan den zeer ervaarenen en kundigen heere Ferdinand Dejean, onlangs beroemd stads heelmeester der stad Batavia. Behelzende eene naauwkeurige beschrijving der Amboinsche haagdis. Benevens eene zeer wel-gelijkende afbeelding deezer volstrekt-nieuwe, fraaije en tot heden toe genoegzaam onbekende soort, door bekwaame kunstenaars vervaardigt*. Amsterdam.
- Schulte, J.A., Melville, J. & Larson, A. (2003) Molecular tests of phylogenetic taxonomies: a general procedure and example using four subfamilies of the lizard family Iguanidae. *Molecular Phylogenetics and Evolution*, 10, 367–376.
- Shaw, G. (1802) *General Zoology, or Systematic Natural History. Vol III. Part I. Amphibia*. London: G Kearsley.
- Townsend, T.M., Mulcahy, D.G., Noonan, B.P., Sites, J.W., Kuczynski, C.A., Wiens, J.J. & Reeder, T.W. (2011) Phylogeny of iguanian lizards inferred from 29 nuclear loci, and a comparison of concatenated and species-tree approaches for an ancient, rapid radiation. *Molecular Phylogenetics and Evolution*, 61, 363–380.
- Wells, R.W. (2007a) Some taxonomic and nomenclatural considerations on the Class Reptilia in Australia. A new genus of the family Chelidae from Eastern Australia. *Australian Biodiversity Record*, 3, 1–13.
- Wells, R.W. (2007b) Some taxonomic and nomenclatural considerations on the Class Reptilia in Australia. Notes on the recently described freshwater turtle *Chelodina canni* McCord and Thomson, 2002 and a redescription of *Chelodina rankini* Wells and Wellington, 1985. *Australian Biodiversity Record*, 1, 1–5.
- Wells, R.W. (2007c) Some taxonomic and nomenclatural considerations on the Class Reptilia in Australia. Some comments on the *Elseya dentata* (Gray, 1863) complex with redescriptions of the Johnstone River Snapping Turtle, *Elseya stirlingi* Wells and Wellington, 1985 and the Alligator Rivers Snapping Turtle, *Elseya jukesi* Wells, 2002. *Australian Biodiversity Record*, 2, 1–12.
- Wells, R.W. (2009) Some taxonomic and nomenclatural considerations on the Class Reptilia in Australia. A new species of freshwater turtle in the genus *Wollumbinia* Wells 2007 (Reptilia: Chelidae) from eastern Australia. *Australian Biodiversity Record*, 2009, 1–12.
- Wells, R.W. & Wellington, C.R. (1983) A synopsis of the class Reptilia in Australia. *Australian Journal of Herpetology*, 1, 73–129.
- Wells, R.W. & Wellington, C.R. (1985) A classification of the Amphibia and Reptilia of Australia. *Australian Journal of Herpetology Supplementary Series*, 1, 1–61.
- Wilson, S.K. (2005) *A Field Guide to Reptiles of Queensland*. Sydney: New Holland Press.
- Wilson, S.K. & Swan, G. (2003) *A Complete Guide to Reptiles of Australia*. Sydney: Reed New Holland.
- Wilson, S.K. & Swan, G. (2008) *A Complete Guide to Reptiles of Australia* (2nd ed.). Sydney: New Holland Publishers.
- Wilson, S.K. & Swan, G. (2010) *A Complete Guide to Reptiles of Australia* (3rd ed.). Sydney: New Holland Publishers.