



<http://dx.doi.org/10.11646/zootaxa.3635.5.11>

<http://zoobank.org/urn:lsid:zoobank.org:pub:EFEACFDF-204E-4E22-8880-2E5A5BAFE578>

Designation of a lectotype for *Deuterodon pedri* Eigenmann, 1908 (Characiformes: Characidae)

CARLOS ALBERTO SANTOS DE LUCENA¹ & ZILDA MARGARETE SEIXAS DE LUCENA²

Museu de Ciências e Tecnologia da Pontifícia Universidade Católica do Rio Grande do Sul. Av. Ipiranga 6681, prédio 40. 90619-900. Porto Alegre, RS, Brazil. E-mail: ¹lucena@pucrs.br; ²margarete@pucrs.br

Deuterodon pedri was briefly described in 1908 by Carl Eigenmann on the basis of seven specimens of 73–100 mm standard length (SL), all "in very poor condition" according to the author. In the original description, Eigenmann did not mention the location where the specimens were captured, even though the publication presented results of the Thayer expedition to Brazil (Eigenmann 1908: 93).

Eigenmann (1927:348) later redescribed the species, adding new data, and indicated the seven specimens as Cotypes (=syntypes) (MCZ 21081), all collected during the Thayer Expedition in "Rio San Antonio, Santa Anna de Ferros" (currently Ferros, Minas Gerais, Doce River basin). Both Eschemeyer (2012) and the database of the Museum of Comparative Zoology collection of fishes (<http://mczbase.mcz.harvard.edu/guid/MCZ:Fish:21081>) indicate five syntypes instead of the seven originally noted.

We had the opportunity to examine two (58.7 mm and 78.4 mm SL) of these five specimens and found that they represent distinct species, readily distinguishable by the shape of the cusps on premaxillary teeth and the way in which the dentary teeth decrease in size. These two traits are mentioned in the redescription of the species by Eigenmann (1927) and are present in the 78.4 mm SL specimen: "the denticles [=cusps] [of] the second row of teeth [are] in a slight crescent" and "dentary with about ten teeth graduated from the minute lateral teeth to the large first tooth." However, these traits are different in the smaller specimen (58.7 mm SL), in which the premaxillary inner row teeth are straight and dentary teeth decrease abruptly as of the fifth tooth. Furthermore, this specimen has tricuspid premaxillary teeth on the outer series and quadri- to pentacuspoid teeth on the inner series; among the dentary teeth, the first five are pentacuspoid, the sixth quadricuspoid, the seventh tricuspid and eighth bicuspoid (vs. premaxillary teeth on outer series pentacuspoid, the first tooth of the inner series pentacuspoid, the other four hexa- to heptacuspoid, and the dentary teeth hexa- to heptacuspoid in the specimen of 78.4 mm SL).

We did not examine the other three syntype specimens (also in very poor condition), but to establish the application of the name *Deuterodon pedri* the 78.4 mm SL specimen is designated here as lectotype. Consequently, the other four specimens are paralectotypes (and remain unidentified). A redescription of the species is urgently needed due to the poor condition of the types.

Acknowledgements

Marilyn Weitzman sent data of the two syntypes in the beginning of our work on *Deuterodon* in 1992. Karten Hartel (MCZ) allowed Paulo Buckup (MNRJ) to send us the syntypes. A. Leyva revised the text to improve our writing in English.

References

- Eigenmann, C.H. (1908) Preliminary descriptions of new genera and species of tetragonopterid characins (Zoölogical Results of the Thayer Brazilian expedition). *Bulletin of the Museum of Comparative Zoology*, 52(6), 91–106
- Eigenmann, C.H. (1927) The American Characidae. Cambridge. *Memoirs of the Museum of Comparative Zoology*, 53(4), 311–428.
- Eschemeyer, W. N. (2012) *Catalog of Fishes*. California Academy of Sciences (<http://research.calacademy.org/research/ichthyology/catalog/fishcatmain.asp>). Electronic version accessed 22 November 2012.