



Zootaxa 3697 (1): 001–088
www.mapress.com/zootaxa/

Copyright © 2013 Magnolia Press

Monograph

ISSN 1175-5326 (print edition)

ZOOTAXA

ISSN 1175-5334 (online edition)

<http://dx.doi.org/10.11646/zootaxa.3697.1.1>

<http://zoobank.org/urn:lsid:zoobank.org:pub:492264BB-E919-447D-9D67-C226DE21A0CE>

ZOOTAXA

3697

Taxonomic exploration of Neotropical Microdontinae (Diptera: Syrphidae) mimicking stingless bees

MENNO REEMER

Naturalis Biodiversity Center / EIS, P.O. Box 9517, 2300 RA Leiden, the Netherlands, menno.reemer@naturalis.nl



Magnolia Press
Auckland, New Zealand

Accepted by C. Kehlmaier: 18 Jun. 2013; published: 9 Aug. 2013

Licensed under a Creative Commons Attribution License <http://creativecommons.org/licenses/by/3.0>

MENNO REEMER

Taxonomic exploration of Neotropical Microdontinae (Diptera: Syrphidae) mimicking stingless bees
(*Zootaxa* 3697)

88 pp.; 30 cm.

9 Aug. 2013

ISBN 978-1-77557-238-1 (paperback)

ISBN 978-1-77557-239-8 (Online edition)

FIRST PUBLISHED IN 2013 BY

Magnolia Press

P.O. Box 41-383

Auckland 1346

New Zealand

e-mail: zootaxa@mapress.com

<http://www.mapress.com/zootaxa/>

© 2013 Magnolia Press

ISSN 1175-5326 (Print edition)

ISSN 1175-5334 (Online edition)

TABLE OF CONTENTS

ABSTRACT	4
INTRODUCTION	4
MATERIAL AND METHODS	5
Acronyms for collections	5
Terminology	6
Lectotype designations	6
KEYS	6
Key to genera of Neotropical <i>Microdantinae</i> mimicking stingless bees	7
Key to species of <i>Carreramyia</i>	8
Key to species of <i>Ceratophya</i>	8
Key to species of <i>Hypselosyrphus</i>	8
Key to species of <i>Mermerizon</i>	9
Key to species of <i>Stipomorpha</i>	9
Key to species of <i>Ubristes</i>	10
Species accounts: descriptions, redescrptions and notes	10
<i>Carreramyia flava</i> (Sack, 1941)	10
<i>Carreramyia megacephalus</i> (Shannon, 1925)	12
<i>Carreramyia megacera</i> spec. nov.	12
<i>Carreramyia tigrina</i> spec. nov.	14
<i>Ceratophya argentinensis</i> spec. nov.	15
<i>Ceratophya carinifacies</i> (Curran, 1934)	17
<i>Ceratophya longicornis</i> Wiedemann, 1824 (excluded from <i>Ceratophya</i>)	19
<i>Ceratophya notata</i> Wiedemann, 1824	19
<i>Ceratophya panamensis</i> (Curran, 1930)	20
<i>Ceratophya scolopus</i> (Shannon, 1927)	22
<i>Hypselosyrphus amazonicus</i> Reemer, 2013	23
<i>Hypselosyrphus anax</i> (Thompson, 1976)	25
<i>Hypselosyrphus corbiculipes</i> Papavero, 1962	26
<i>Hypselosyrphus helvus</i> spec. nov.	28
<i>Hypselosyrphus marshalli</i> spec. nov.	28
<i>Hypselosyrphus maurus</i> spec. nov.	29
<i>Hypselosyrphus pingo</i> spec. nov.	32
<i>Hypselosyrphus plaumanni</i> (Curran, 1940)	34
<i>Hypselosyrphus pseudorhoga</i> spec. nov.	35
<i>Hypselosyrphus trigonus</i> Hull, 1937	35
<i>Hypselosyrphus ulopodus</i> (Hull, 1944)	37
<i>Hypselosyrphus vexillipennis</i> spec. nov.	38
<i>Mermerizon inbio</i> Reemer, 2013	40
<i>Mermerizon mellosus</i> spec. nov.	41
<i>Mermerizon mesmerizus</i> spec. nov.	41
<i>Stipomorpha apicula</i> (Curran, 1930)	43
<i>Stipomorpha crematogastris</i> spec. nov.	45
<i>Stipomorpha dichromata</i> spec. nov.	47
<i>Stipomorpha elcopala</i> spec. nov.	47
<i>Stipomorpha fallax</i> spec. nov.	48
<i>Stipomorpha fraudator</i> (Shannon, 1927)	49
<i>Stipomorpha goettei</i> (Shannon, 1927) comb. nov.	51
<i>Stipomorpha guianica</i> (Curran, 1925)	54
<i>Stipomorpha inarmata</i> (Curran, 1925)	55
<i>Stipomorpha lacteipennis</i> (Shannon, 1927)	57
<i>Stipomorpha lanei</i> (Curran, 1936)	58
<i>Stipomorpha litoralis</i> (Papavero, 1964)	59
<i>Stipomorpha mackiei</i> (Curran, 1940)	59
<i>Stipomorpha maculipennis</i> spec. nov.	60
<i>Stipomorpha mendax</i> spec. nov.	62
<i>Stipomorpha micromidas</i> (Shannon, 1925)	63
<i>Stipomorpha mixta</i> (Curran, 1940)	64
<i>Stipomorpha panamana</i> spec. nov.	66
<i>Stipomorpha simillima</i> (Hull, 1950)	67
<i>Stipomorpha spuria</i> spec. nov.	68
<i>Stipomorpha tenuicauda</i> (Curran, 1925)	70
<i>Stipomorpha trigoniformis</i> (Shannon, 1927)	72

<i>Stipomorpha wheeleri</i> (Mann, 1928)	73
<i>Stipomorpha zophera</i> spec. nov.	75
<i>Ubristes flavitibia</i> Walker, 1852	79
<i>Ubristes ictericus</i> spec. nov.	80
<i>Ubristes jaguarinus</i> spec. nov.	82
REST GROUP	83
<i>Microdon</i> (<i>Chymophila</i>) <i>angulatus</i> Hull, 1943	83
<i>Ubristes chrysopygus</i> Giglio-Tos, 1892	85
Discussion	85
ACKNOWLEDGEMENTS	86
REFERENCES	87

ABSTRACT

Several species of Neotropical Microdontinae (Diptera: Syrphidae) are mimics of stingless bees. Most of these species have previously been grouped in *Ubristes* Walker, 1852, with *Carreramyia* Doesburg, 1966, *Hypselosyrphus* Hull, 1937 and *Stipomorpha* Hull, 1945 treated as synonyms in recent literature. The species of the recently described genus *Mermerizon* Reemer, 2013 are also treated in the present paper. Recent evidence (Reemer and Ståhls 2013a, b) supports an independent origin for all of these taxa, which is why they are now treated as different genera. The present paper revises all specific taxa previously associated with these genera. A total number of 52 species is treated, 22 of which are described as new. These are divided among the genera as follows: *Carreramyia* (4 species, 2 new: *C. megacera*, *C. tigrina*), *Ceratophya* (5 species, 1 new: *C. argentinensis*), *Hypselosyrphus* (12 species, 6 new: *H. helyus*, *H. marshalli*, *H. maurus*, *H. pingo*, *H. pseudorhoga*, *H. vexillipennis*), *Mermerizon* (3 species, 2 new: *M. mellosus*, *M. mesmerizus*), *Stipomorpha* (25 species, 9 new: *S. crematogastris*, *S. dichromata*, *S. elcopala*, *S. fallax*, *S. maculipennis*, *S. mendax*, *S. panamana*, *S. spuria*, *S. zophera*), *Ubristes* (3 species, 2 new: *U. ictericus*, *U. jaguarinus*). *Microdon scolopus* Shannon, 1927, previously classified in *Ubristes*, was recently transferred to *Ceratophya* Wiedemann, 1824, which is why this genus is also treated in this paper. *Ceratophya longicornis* Wiedemann, 1824 is excluded from *Ceratophya* and treated as a species *incertae sedis*. Two other species are excluded, because they belong to other groups of Microdontinae not treated in the present paper: *Microdon angulatus* Hull, 1943 (now in *Microdon* subgenus *Chymophila* Macquart, 1834) and *Ubristes chrysopygus* Giglio-Tos, 1892 (now in *Peradon* Reemer, 2013). Three new synonyms are proposed: *Microdon procteri* Curran, 1941 syn. nov. = *Microdon procedens* Curran, 1941 syn. nov. = *Ubristes flavitibia* Walker, 1852; *Microdon triangularis* Curran, 1940 syn. nov. = *Microdon lacteipennis* Shannon, 1927. Two specific taxa previously considered as synonyms are rendered valid status: *Ubristes litoralis* Papavero, 1964 stat. rev. (placed in *Stipomorpha*) and *Ubristes plaumanni* Curran, 1940 stat. rev. (placed in *Hypselosyrphus*). One new name is introduced to replace a junior primary homonym: *Hypselosyrphus amazonicus* nom. nov. for *Microdon scutellaris* Shannon, 1927, preoccupied by Schummel (1842). Lectotypes are designated for *Stipomorpha goettei* (Shannon, 1927) and *Stipomorpha lactipennis* (Shannon, 1927). A key to the genus-groups and to the species is given. The genus *Rhoga* Walker, 1857 is included in the key to the genus-groups, but specific taxonomy is not worked out. The paper concludes with some considerations on mimicry as a possible drive for speciation, and on species of *Stipomorpha* visiting flowers.

Key words: revision, new species, key, mimicry of Meliponini

INTRODUCTION

Mimicry of noxious Hymenoptera commonly occurs in hoverflies (Diptera: Syrphidae). For instance, approximately 22% of all European species are considered to be mimics of bees or aculeate wasps to varying extent (Gilbert 2005). The potential selective advantage of mimicking noxious insects is obvious. It may come as a surprise, therefore, that several species of Syrphidae seem to mimic apparently harmless models of aculeate Hymenoptera: the stingless bees (Apidae: Apinae: Meliponini), which are characterized by their rudimentary sting. Harmless these bees may seem, but certain taxa are known to secrete formic acid from cephalic glands, which can cause an itching or even burning sensation when bitten by such a bee (Roubik *et al.* 1987). Such chemical properties of stingless bees may be an explanation for the noxiousness underlying their use of models for the evolution of mimicry.

Stingless bees are found all over the tropics, but their greatest diversity (about 75% of 500 species) occurs in the Neotropics (Costa *et al.* 2003). Likewise, this seems to be the only region in which many species of Syrphidae have evolved as mimics of stingless bees. Unpublished observations by the author in Surinam indicate that

stingless-bee mimics occur among all three currently recognized subfamilies of Syrphidae: Syrphinae (e.g. *Ocyptamus* Macquart, 1834), Eristalinae (e.g. *Copestylum* Macquart, 1846, *Lepidomyia* Loew, 1864) and Microdontinae. Especially the latter subfamily is rich in mimics of stingless bees.

Most species of Neotropical Microdontinae resembling stingless bees have traditionally been grouped in the genus *Ubristes* Walker, 1852 (Thompson *et al.* 1976). This genus was originally erected for *U. flavitibia* Walker, 1852. Shannon (1927) applied the name *Ubristes* to several other species of Microdontinae with long, brush-like pilosity on the hind-tibia, resembling the corbicula of stingless bees. Subsequent authors, such as Hull (1949) and Van Doesburg (1966), adopted this application of the name, sometimes considering *Ubristes* as a genus, sometimes as a subgenus of *Microdon* Meigen, 1803. In the catalogue of South American Syrphidae, Thompson *et al.* (1976) treated *Ubristes* as a subgenus of *Microdon*, in which they included 31 specific taxon names (two of which as synonyms). They included the following genus-group names as synonyms of *Ubristes*: *Carreramyia* Doesburg, 1966, *Hypselosyrphus* Hull, 1937 and *Stipomorpha* Hull, 1945. Subsequently, Cheng and Thompson (2008) considered *Ubristes* as a genus, *Carreramyia* as a subgenus of *Ubristes*, and *Hypselosyrphus* and *Stipomorpha* as species groups of *Ubristes*. Recent evidence indicates that all of these genus group names represent unrelated taxa, implying that mimicry of stingless bees has evolved several times independently within the Microdontinae (Reemer and Ståhls 2013b). Therefore, the groups have all been given generic status, and the species previously listed under *Ubristes* were divided over these genera by Reemer and Ståhls (2013a). So far, the taxonomy of the species belonging to these genera has not been studied in detail. Keys to species were published by Shannon (1927), Curran (1940) and Van Doesburg (1966), but these are very incomplete.

The aim of the present paper is to revise the species listed under *Ubristes* by Thompson *et al.* (1976), to attribute them to the available supraspecific taxa, and to describe hitherto undescribed species. Three new species are assigned to the recently described genus *Mermerizon* Reemer, 2013 (in Reemer and Ståhls 2013a). Although Reemer and Ståhls (2013a) provide a key to all genera of Microdontinae, an additional key is here presented to distinguish the genera of Neotropical Microdontinae resembling stingless bees (i.e. those taxa with widened and/or brush-like pilose hind tibiae). For each genus keys to species are given. The genus *Ceratophya* Wiedemann, 1824 is also included in this paper, because one of its species (*C. scolopus* (Shannon, 1927)) used to be included in *Ubristes* s.l. Although species of *Rhoga* Walker, 1857 also have ‘corbiculate’ hind tibiae, making them resemble stingless bees, this taxon has always been considered distinct from *Ubristes* auct. It has been well characterized by e.g. Cheng and Thompson (2008) and Hull (1949). Reemer and Ståhls (2013b) recovered *Rhoga* within *Hypselosyrphus*, but refrained from changing the taxonomy of these groups because of limited taxon sampling. To avoid confusion *Rhoga* is also included in the generic key provided in this paper, but no key to the species is provided nor are the species treated separately.

MATERIAL AND METHODS

Acronyms for collections

AMNH	American Museum of Natural History, New York
BMNH	British Museum (Natural History), London
CNC	Canadian National Collection, Ottawa
CSCA	California State Collection of Arthropods, Sacramento
CU	Cornell University, Ithaca
INBIO	Instituto Nacional de Biodiversidad, Santo Domingo de Heredia (Costa Rica)
MCZ	Museum of Comparative Zoology, Harvard
MRSN	Museu Regionale di Scienze Naturali, Torino
MZLU	Museum of Zoology, Lund
MZUSP	Museu de Zoologia da Universidade de São Paulo
NMW	Naturhistorisches Museum Wien, Vienna
PUCE	Pontífica Universidad Católica del Ecuador, Quito
RMNH	National Museum of Natural History, Leiden
SEMC	Snow Entomological Museum, University of Kansas, Lawrence

SNSD	Senckenberg Naturhistorische Sammlungen Dresden
UCD	University of California, Davis
USNM	United States National Museum, Smithsonian Institutions, Washington D.C.
ZMAN	Zoölogisch Museum Amsterdam (recently included in collection RMNH, Leiden)
ZMHB	Museum für Naturkunde der Humboldt Universität, Berlin

Terminology

For morphology the terminology of McAlpine (1981) is used, as specifically applied to Syrphidae by Thompson (1999). For some additional characters terms of Hippa and Ståhls (2005) are used (e.g. antennal fossa, antetergite). A description and discussion of morphology of Microdontinae can be found in Reemer and Ståhls (2013b).

Lectotype designations

Lectotype designations have been carried out in accordance with the 4th edition of the ICZN Code, in order to fix the names involved and ensure a universal and consistent interpretation of the taxa in the future.

KEYS

Keys to species of *Ubristes* auct. have been published by Shannon (1927), Curran (1940) and Van Doesburg (1966), but these only contain a small part of the species treated in this paper. The following key covers all species listed under *Ubristes* by Thompson *et al.* (1976) (including *Carreramyia*, *Hypselosyrphus* and *Stipomorpha*), all species of *Ceratophya*, and all new species described in the present paper, which include the species of the recently described genus *Mermerizon* Reemer, 2013. A summary of diagnostic characters for distinguishing the genera treated here is given in table 1.

Not all Neotropical Microdontinae resembling stingless bees can be identified with the keys below. However, the key should work for all species included in the treated genera and species groups. In general, Neotropical microdontines with widened hind tibiae and / or brushes of long pile on the hind tibiae should get identified using these keys. If a specimen does not belong to one of those groups, the key to the genera will tell you so.

Be aware that probably several undescribed species lurk on the South American continent. Always check identification with the figures, diagnoses and (re)descriptions as given in the species accounts.

TABLE 1. Morphological differences between the genera of Microdontinae treated in the present paper.

	<i>Carreramyia</i>	<i>Ceratophya</i>	<i>Hypselosyrphus</i>	<i>Mermerizon</i>	<i>Rhoga sepulchrasilva</i>	<i>Stipomorpha</i>	<i>Ubristes</i>
Basoflagellomere	male: bifurcate; female: unfurcate	unfurcate	unfurcate	unfurcate	unfurcate	unfurcate	unfurcate
Face	wider than one eye in frontal view	as wide as or wider than one eye in frontal view	narrower than one eye in frontal view	narrower than or as wide as one eye in frontal view	narrower than one eye in frontal view	narrower than or as wide as one eye in frontal view (except wider in <i>S. wheeleri</i>)	wider than one eye in frontal view
Eye	bare	bare	short pilose	bare	short pilose	bare or appearing so under low magnification	bare

..... continued on the next page

TABLE 1. (Continued)

	<i>Carreramyia</i>	<i>Ceratophya</i>	<i>Hypselosyrphus</i>	<i>Mermerizon</i>	<i>Rhoga sepulchrasilva</i>	<i>Stipomorpha</i>	<i>Ubristes</i>
Occiput	dorsally wide, laterally and ventrally narrow	narrow over entire length	narrow at least dorsally	dorsally wide, ventrally narrow	wide over entire length	dorsally wide, ventrally narrow	dorsally wide, ventrally narrow
Scutellum	apicomediaally sulcate	varies between species	triangular or apicomediaally more or less sulcate	semicircular, not sulcate	apicomediaally slightly sulcate	semicircular, not sulcate	semicircular, not sulcate
Anepimeron	bare on ventral half	entirely pilose (pile very short)	entirely pilose	bare on ventral half	entirely pilose	varies between species	entirely pilose
Vein sc	joins costal vein at about same level as crossvein r-m	joins costal veins proximal of crossvein r-m	joins costal vein distal of crossvein r-m	joins costal vein distal of crossvein r-m	joins costal vein distal of crossvein r-m	joins costal vein at same level as or proximal of crossvein r-m	joins costal veins proximal of crossvein r-m
Vein R4+5	without appendix	with appendix	without appendix	with appendix	without appendix	mostly with, sometimes without appendix	with appendix
Tergites 3 and 4	fused	not fused	fused	fused	fused	fused	fused
Sternite 1	bare or pilose	pilose	pilose	bare	bare *	bare	pilose
Membranes between sternites 1–3	normal	normal	normal	normal	normal	with very wide membranous parts in between	normal
Epandrium	without lateral 'fenestrae'	without lateral 'fenestrae'	without lateral 'fenestrae'	without lateral 'fenestrae'	without lateral 'fenestrae'	without lateral 'fenestrae'	with lateral 'fenestrae'
Hypandrium	basally bulged	basally not bulged	basally bulged	basally bulged	basally bulged	basally bulged	basally not bulged
Aedeagus	furcate, furcation point close to apex	furcate, furcation point close to base	furcate, furcation point close to apex	furcate, furcation point close to apex	furcate, furcation point close to apex	unfurcate	furcate, furcation point close to base; long and slender

*: In certain other *Rhoga* species sternite 1 is pilose.

Key to genera of Neotropical Microdontinae mimicking stingless bees

1. Hind tibia without long pile and not conspicuously widened. other groups of Microdontinae
- Hind tibia appearing corbiculate (as in bees): with long, brush-like pile and/ or hind tibia conspicuously widened medially or apically 2
2. Scutellum with calcars other groups of Microdontinae
- Scutellum without calcars. 3
3. Sternites 2 and 3 separated by unusually wide membranous parts, about as wide as tergite 2 medially or wider (fig. 152, 153) (may be hard to see in dry specimens). Aedeagus unfurcate (figs. 230–251). *Stipomorpha*
- Sternites 2 and 3 not separated by very wide membranous part. Aedeagus furcate apically or basally (figs. 55–57, 108–112, 124–126, 265–266). 4
4. Vein R4+5 with posterior appendix in cell r4+5 (e.g. figs. 40, 161, 165). 5
- Vein R4+5 without posterior appendix in cell r4+5 (e.g. figs. 8, 62, 87) 7
5. Tergites 3 and 4 not fused, posterior margin of tergite 3 strongly overlapping with tergite 4 (figs. 31, 37). Tergite 4 in lateral view perpendicular to tergite 2. Face in most species laterally depressed, appearing somewhat carinate medially *Ceratophya*

- Tergites 3 and 4 fused, posterior margin of tergite 3 not overlapping with tergite 4 (figs. 9, 143, 205). Tergite 4 in lateral view not perpendicular to tergite 2. Face laterally not depressed 6
- 6. Tergite 2 with lateral tubercles (fig. 258). Basoflagellomere longer than scape. Face in frontal view wider than an eye (fig. 259).. *Ubristes*
- Tergite 2 without lateral tubercles. Basoflagellomere about as long as scape. Face in frontal view narrower than an eye (fig. 115, 120, 123) *Mermerizon*
- 7. Vertex wider than an eye (fig. 2, 8, 21). Basoflagellomere at least four times as long as scape; bifurcate in male. *Carreramyia*
- Vertex narrower than an eye (fig. 60, 85). Basoflagellomere maximally twice as long as scape, but usually shorter than scape; not furcate 8
- 8. Occiput wide dorsally (also ventrally) (fig. 1), wider than length of ocellar triangle, also wider than length of pedicel *Rhoga* (no key to species included in this paper)
- Occiput narrow dorsally, usually also ventrally (fig. 61, 86), but not always (fig. 103), narrower than length of ocellar triangle, also narrower than length of pedicel *Hypselosyrphus*

Key to species of *Carreramyia*

- 1. Abdomen yellowish with dark vittae medially and laterally (fig. 2, 17) 2
- Abdomen unicolorous black or yellowish brown (fig. 8, 13) 3
- 2. Scutum mostly yellow; only with dark stripe laterally between notopleuron suture and posterior margin. Scutellum yellow. Hind leg: tibia wider than femur. Female basoflagellomere with arista; shape as in fig. 7 *flava*
- Scutum mostly black; with yellow margins and narrow yellow lines. Scutellum black laterally, yellow medially. Hind leg: tibia wider than femur. Female basoflagellomere without arista; shape as in fig. 19. *tigrina*
- 3. Yellowish brown species. *megacephalus*
- Black species. *megacera*

Key to species of *Ceratophya*

- 1. First tarsomeres of all tarsi with strong ventrobasal tooth (fig. 54). Tergites 2 and 3 black. *scolopus*
- First tarsomeres without ventrobasal tooth. Tergites 2 and 3 at least partly yellow 2
- 2. Scutellum (dorsal view) semicircular, without apicomedian sulcus 3
- Scutellum (dorsal view) with (sometimes weak) apicomedian sulcus. 4
- 3. Tergite 2 posteromedially black, laterally with wide, oblique yellow vitta (fig. 30) *carinifacies*
- Tergite 2 posteromedially yellow, laterally black (fig. 24) *argentinensis*
- 4. Scutellum with weak apicomedian sulcus (fig. 41). Male: tergite 3 blackish brown with posterior margin broadly yellow. Female: tergites predominantly brownish black; tergite 2 with two oblique yellow vittae, tergites 3 and 4 with yellow posterior margins *notata*
- Scutellum with deep apicomedian sulcus (fig. 46). Male: tergite 3 blackish brown with posterior margin broadly yellow and with yellow lateral vittae (based on description of Curran 1930; in holotype the abdomen is missing). Female: tergites yellow, only brownish along lateral margins *panamensis*

Key to species of *Hypselosyrphus*

- 1. Scutellum triangular, apex acute. (fig. 65) *amazonicus*
- Scutellum not triangular: semicircular or apicomediaally sulcate. 2
- 2. Alula entirely microtrichose 6
- Alula with small bare area basomedially 3
- 3. Thorax and abdomen entirely yellow (fig. 70, 71) *helvus*
- Thorax and abdomen largely black 4
- 4. Basoflagellomere about as long as scape or longer. Wing either with strongly contrasting yellowish patch posteriad of pterostigma (male) or with middle 1/3 strongly contrasting whitish yellow (female) (figs. 74, 75).. *marshalli*
- Basoflagellomere maximally half as long as scape. Wing without strongly contrasting yellow parts 5
- 5. Tergite 4 with posterior margin widely yellowish; sternite 4 largely yellowish. Scutellum entirely black pilose. Wing tinged yellowish on apical third, otherwise brownish.. *anax*
- Tergite and sternite 4 entirely blackish brown. Scutellum black pilose, except pale pilose posteroventrally. Wing entirely tinged brownish. *plaumanni*
- 6. Wing with contrasting pattern of dark brown and yellow fasciae (fig. 87, 106). 7
- Wing without contrasting colour pattern (may be tinged with brown or yellow). 8
- 7. Mesonotum and scutellum yellow pilose *pingo*
- Mesonotum and scutellum black pilose *vevillipennis*

8.	Hind tibia yellow, yellow pilose. Only female known	<i>corbiculipes</i>
–	Hind tibia at least partly brown, black pilose	9
9.	Abdomen orange to reddish brown	11
–	Abdomen black	10
10.	Occiput narrow over entire length (fig. 79)	<i>maurus</i>
–	Occiput ventrally widened (fig. 103)	<i>ulopodus</i>
11.	Scutellum apicomediaally sulcate. Face dark brown. Thorax dark brown.	<i>trigonus</i>
–	Scutellum more or less semicircular, not sulcate. Face yellow. Thorax yellow, except for blackish brown maculae on mesoscutum.	<i>pseudorhoga</i>

Key to species of *Mermerizon*

1.	Hind tibia with appressed pile, which are shorter than half the width of the tibia. Antenna entirely black. Wing with vaguely defined, greyish transverse fasciae (fig. 122)	<i>mesmerizus</i>
–	Hind tibia with more or less erect brush-like pile, which are at least half as long as the width of the tibia. Antenna with at least basoflagellomere yellow. Wing without greyish fasciae (fig. 117, 118).	2
2.	Mesonotum and scutellum entirely yellow pilose. Hind femur and basal half of hind tibia yellow pilose	<i>mellosus</i>
–	Mesonotum, scutellum and hind leg entirely black pilose.	<i>inbio</i>

Key to species of *Stipomorpha*

1.	Anepisternum pilose posterodorsal margin pilose (often only sparsely)	2
–	Anepisternum with posterodorsal margin bare (only anterior part pilose)	14
2.	Katepisternum dorsally pilose.	3
–	Katepisternum dorsally bare	7
3.	Abdomen with contrasting colour pattern: tergite 2 orange brown, tergites 3 and 4 blackish (fig. 134)	<i>dichromata</i>
–	Abdomen more or less unicolourous.	4
4.	Abdomen yellowish brown	5
–	Abdomen blackish.	6
5.	Front and mid legs yellow.	<i>micromidas</i>
–	Front and mid legs black (except apical tarsomeres yellow).	<i>elcopala</i>
6.	Wing with whitish transverse fascia posterior to pterostigma (view against dark background). Male genitalia as in fig. 237	<i>lacteipennis</i>
–	Wing without whitish fascia. Male genitalia as in fig. 239.	<i>litoralis</i> (only male known)
7.	Alula entirely microtrichose	8
–	Alula partly bare basomedially	9
8.	Head in frontal view clearly wider than high (fig. 130). Basoflagellomere at least four times as long as wide. Frontal ocellus round, not split in two	<i>crematogastris</i> spec. nov. (only female known)
–	Head in frontal view about as wide as high (fig. 222). Basoflagellomere less than three times as long as wide. Frontal ocellus split in two.	<i>wheeleri</i>
9.	Vertex shining black, more or less convex (fig. 211, 214)	10
–	Vertex mostly yellow, irregularly swollen (not convex) (fig. 179, 198).	12
10.	Vein R2+3 joins costal vein at about same level as junction of M1 and R4+5. Wing more or less colourless. Male genitalia as in fig. 247	<i>spuria</i>
–	Vein R2+3 joins costal vein clearly distal of junction of M1 and R4+5. Wing tinged with yellow on anterobasal ½.	11
11.	Male genitalia as in fig. 248: surstylus in lateral view slender with 'hooked' appearance. Alula: band of microtrichia along posterior margin maximally as wide as 1/6 of width of alula (fig. 176).	<i>tenuicauda</i>
–	Male genitalia as in fig. 240: surstylus in lateral view approximately quadrate. Alula: band of microtrichia along posterior margin about as wide as 1/4 of width of alula (fig. 175).	<i>mackiei</i>
	[Two externally very similar species, both quite variable in colouration. In both species the hind leg and the metanotum can be entirely yellow or almost entirely dark. The character of the distribution of microtrichia on the alula should be used with caution, as it could only be verified on a small number of males.]	
12.	Wing with dark brown spot anteromedially. Male genitalia as in fig. 241	<i>maculipennis</i>
–	Wing without dark brown spot anteromedially.	13
13.	Face narrower than one eye in frontal view. Hind tibia as wide as or slightly wider than hind femur. Male genitalia as in fig. 235.	<i>guianica</i>
–	Face wider than one eye in frontal view. Hind tibia about twice as wide as hind femur. Male genitalia as in fig. 245	<i>panamana</i>
14.	Alula entirely microtrichose	15
–	Alula partly or entirely bare	23
15.	Wing uniformly hyaline or slightly infuscated, without whitish cloud or fascia (may be tinged with yellow)	16

- Wing with yellow or white cloud at or posterior to pterostigma, sometimes small, sometimes extending to posterior wing margin, thus forming a transverse fascia (may be inconspicuous, be sure to view against dark background!) 20
- 16. Abdomen black. Face with median black vitta occupying about 3/4 of width of face. *trigoniformis* (only male known)
- Abdomen yellow. Face yellow or with narrow, vague brownish median vitta. 17
- 17. Front- and mid-legs: first 3-4 tarsomeres dark brown. Basoflagellomere slightly shorter than scape. *fraudator*
- Front- and mid-legs: tarsi entirely yellow. Basoflagellomere at least as long as scape. 18
- 18. Abdomen somewhat constricted: narrowest width at tergite 3 (less pronounced in female). Pilosity of hind tibia less than half as long as width of tibia. Female: vertex at least partly yellow, sometimes with vague dark markings. *lanei*
- Abdomen not constricted at tergite 3; tapering from anterior margin of tergite 3 towards apex. Pilosity of hind tibia about as long as half the width of the tibia. Female: vertex black. 19
- 19. Anterior margin of tergite 2 not curled around tergite 1 laterally (fig. 229). Females: greatest width of tergite 2 posteriad of half its length. More robust species *mixta*
- Anterior margin of tergite 2 more or less curled around tergite 1 (fig. 228). Females: greatest width of tergite 2 at or anterior of half its length. Slender species *mendax*
- 20. Scutellum yellow. 21
- Scutellum black. 22
- 21. Vertex black *mendax* (female)
- Vertex yellow, except ocellar triangle darkened. *goettei*
- 22. Basoflagellomere 1.5 times as long as scape. Vein sc joins costal vein proximal of crossvein r-m. Pale mark on wing large, extended to posterior wing margin. Black facial vitta occupying about 1/2 of width of face. *puerilis* (only female known)
- Basoflagellomere about as long as scape. Vein sc joins costal vein at about same level as crossvein r-m. Pale mark on wing small: reaching from pterostigma to halfway wing or less. Black facial vitta occupying about 1/6 of width of face. Male genitalia as in fig. 246. *simillima*
- 23. Scutellum yellowish 24
- Scutellum brown or black. 26
- 24. Vertex yellow. *goettei*
- Vertex black 25
- 25. Scape 4 times as long as pedicel. Genitalia as in fig. 242. *mendax*
- Scape 2 times as long as pedicel. Genitalia as in fig. 232. *fallax* (only male known)
- 26. Basoflagellomere clearly longer than scape. Anterior part of anepisternum black pilose. *zophera*
- Basoflagellomere slightly shorter than or as long as scape. Anterior part of anepisternum yellow pilose. 27
- 27. Abdomen black or dark brown. *inarmata*
- Abdomen pale orange brown. *apicula*

Key to species of *Ubristes*

- 1. Abdomen black (fig. 252). *flavitibia*
- Abdomen partly or entirely yellow (figs. 256, 261). 2
- 2. Abdomen entirely yellow (fig. 256). *ictericus*
- Abdomen brownish with yellow maculae (fig. 261). *jaguarinus*

Species accounts: descriptions, redescriptions and notes

Carreramyia flava (Sack, 1941)

Figs 2–7.

Ceratophya flava Sack, 1941: 117.

Studied type specimens. HOLOTYPE. Female. PERU. Label 1 (green): "Peru – Rosalina / 28.8.03 / Urubambafl."; label 2 (green): "Coll. W. Schnuse / 1911 - 3"; label 3: [female sign]; label 4: "Ceratophya / flava Schnuse"; label 5 (orange): "16a. / ? Ceratophya Wied ? / flava sp. nov."; label 6: "Staatl. Museum für / Tierkunde, Dresden". Coll. SNSD.

Description (based on holotype)

Adult female. Body size: 5 mm.

Head. Face occupying almost 3/5 of head width in frontal view; yellow with small brown spot laterad of antennal fossa; short yellow pilose. Gena yellow; yellow pilose. Frons yellow, bare. Vertex strongly produced medially; yellow with a vague brown transverse fascia; yellow pilose, with pile getting longer posteriorly. Occiput

yellow; yellow pilose. Eye bare. Antennal fossa about as wide as high. Antenna yellow, with basoflagellomere and base of scape darker; ratio of scape:basoflagellomere approximately as 1:4; pedicel very short, only about 1/18 of basoflagellomere; arista about as long as pedicel, yellowish white.



FIGURES 1–10. 1. *Rhoga sepulchrasilva* male, head lateral. 2–7. *Carreramyia flava* female (holotype). 2. habitus dorsal; 3. habitus lateral; 4. head frontal; 5. head lateral; 6. wing; 7. basoflagellomere. 8–10. *Carreramyia megacephalus* male (Costa Rica, coll. M. Hauser). 8. habitus dorsal; 9. habitus lateral; 10. head frontal.

Thorax. Scutum yellow with brown lateral vitta between notopleuron and posterior margin; yellow pilose, except black pilose on part of lateral brown vitta. Postpronotum yellow; bare. Postalar callus pale yellow; yellow pilose. Scutellum yellow; yellow pilose basally, black pilose apically; sulcate posteromedially. Anepisternum a little convex, without sulcus; brown anteriorly, yellow posteriorly; yellow pilose anterodorsally and posterodorsally. Katepisternum yellow, except brown posterodorsally; sparsely yellow pilose dorsally, bare ventrally. Katepimeron pale yellow; convex; bare. Calypter dark brown. Halter yellow.

Wing: hyaline, vaguely infuscated halfway from stigmal crossvein to r-m and along marginal crossveins; with yellowish white transverse fascia from pterostigma almost to posterior margin (view against dark background); microtrichose, except bare on 1st costal cell, basal 1/2 of cell br, posterobasal 1/3 of cell bm, anterobasal 1/5 of cell cup.

Legs: front and hind legs yellow [mid legs missing in type specimen], with hind tibia somewhat darker; short yellow pilose, except hind tibia dorsally longer appressed black pilose. Coxae and trochanters yellowish brown; yellow pilose.

Abdomen. Tergites yellow with brown markings on the following parts: tergite 1 anterolaterally; tergite 2 laterally and vaguely posteromedially; tergite 3-5 laterally and with median vitta. Tergites short dark pilose, except tergite 2 anterolaterally with longer yellow pile. Sternites yellow; yellow pilose, except sternite 1 bare.

Diagnosis. Distinguished from *C. megacephalus* and *C. megacera* by the striped pattern of the abdomen. From *C. tigrina* it differs by the mostly yellow scutum, the entirely yellow scutellum, the hind leg with its femur wider than its tibia, the presence of an arista in the female and the shape of the female basoflagellomere.

Notes. This species was listed by Thompson *et al.* (1976) under the 'Unrecognized species' of Syrphidae, not placed in any genus of Syrphidae, not even in a subfamily. Examination of the type revealed that Sack (1941) was right in placing this species in the Microdontinae. It fully fits the characters described as diagnostic for *Carreramyia*.

Distribution. Only known from Peru.

***Carreramyia megacephalus* (Shannon, 1925)**

Figs 8–12, 23.

Microdon megacephalus Shannon, 1925: 213. Type locality: Panama.

Studied type specimen. HOLOTYPE. Male. PANAMA: Old Panama, 31.I.1911, leg. A. Busck, coll. USNM.

Additionally studied specimens. COSTA RICA: 2 males & 1 female, Guanacaste, 3 km SE R. Naranjo, 3–8.III.1992, leg. F.D. Parker, males coll. M. Hauser, female coll. RMNH; 1 male, same locality & leg as previous, 11–21.IV.1992, coll. RMNH; 1 female, Guanacaste, Parque Nacional Santa Rosa, Sector Santa Rosa, 5–18.VII.2001, leg. A.R. Deans, coll. M. Hauser.

Diagnosis. Body size 6–8 mm. Immediately distinguished from the other three *Carreramyia*-species by its entirely yellowish colouration, without any dark markings.

Distribution. Known from Costa Rica and Panama.

***Carreramyia megacera* spec. nov.**

Figs 13–16.

Studied type specimens. HOLOTYPE. Female. SURINAM: Commewijne, Peperpot, 05°46'08"N–55°07'33"W, 17–24.II.2006 (malaise trap), leg. M. Reemer, coll. RMNH.

Description (based on holotype)

Adult female. Body size: 6 mm.

Head. Face occupying about 2/5 of head width in frontal view; pale yellow with two vague submedian vittae; with short, sparse black pile, getting longer and more bristly around oral margin. Gena pale yellow, with a few black setae. Frons pale yellow, except for black lunula, black macula laterad of antennal fossa and narrow black line along eye margin; bare. Vertex pale yellow and bare on anterior half, black and with black bristly pile on posterior half; strongly elevated. Occiput black; black pilose dorsally, white pilose ventrally. Eye bare. Antennal

fossa about as wide as high. Antenna black; ratio of scape:basoflagellomere approximately as 1:5; pedicel very short; basoflagellomere very long, about 1.5 times as long as face; arista yellow, about 2/3 of length of scape.



FIGURES 11–21. 11–12. *Carreramyia megacephalus*. 11. head male lateral; 12. head female frontal (Costa Rica, coll. RMNH). 13–16. *Carreramyia megacera* female (holotype). 13. habitus dorsal; 14. habitus lateral; 15. head lateral; 16. wing. 17–21. *Carreramyia tigrina* female (holotype). 17. habitus dorsal; 18. habitus lateral; 19. head frontal; 20. head lateral; 21. head dorsal.

Thorax. Scutum black; bristly black pilose. Postpronotum yellow; bare. Postalar callus yellow; bristly black pilose. Scutellum black; bristly black pilose; deeply sulcate posteromedially; in lateral view making angle of about 45° with scutum. Anepisternum a little convex, without sulcus; black anteriorly, yellow posteriorly; bristly black pilose anterodorsally and along posterior margin, with wide bare part in between. Katepisternum black, except for small yellow macula at dorsal margin; dorsally bristly black pilose, bare ventrally. Katepimeron yellow; convex; bare. Calypter black. Halter yellow.

Wing: hyaline, faintly infuscated between base and stigmal crossvein, with faint yellow cloud apically of stigmal crossvein and crossvein r-m; microtrichose, except bare on 1st costal cell, on cell br except along vena spuria, on most of cell bm except apical 1/6 and a narrow median stripe from base to apex, on anterior 1/3 of cell cup.

Legs: missing in holotype, except middle leg: long and slender; femur black, except faintly yellow near apex; tibia black, except yellow at basal and apical 1/10; tarsus yellow; entirely black pilose. Coxae and trochanters black; bristly black pilose.

Abdomen. Black. Second segment wider than thorax, widest point at posterior margin; tergites 3 and 4 strongly narrowing. Tergites 1 and 2 black pilose, except yellow pilose medially along posterior margin of tergite 2. Tergite 3 black pilose, except for two large submedian patches of yellow pile which reach posterior margin. Tergites 4 and 5 mainly black pilose, with limited yellow pile. Sternites black, with wide yellow membrane between sternites 1 and 2. Sternite 1 bare, other sternites bristly black pilose.

Etymology. The specific epithet refers to the very long antennae: *mezas* (Gr., large), *keras* (Gr., antenna, horn).

Diagnosis. Immediately distinguished from the other three *Carreramyia*-species by its black colouration and the deeply sulcate scutellum.

Notes. The male is unknown, so whether it has the furcate basoflagellomere characteristic for this genus or not can only be guessed at. Nevertheless the female is very similar to *C. megacephalus* in the diagnostic and other important characters: basoflagellomere very long, vertex strongly produced, face very wide, near oral margin with bristly pile, scutellum sulcate, vein R4+5 without appendix, crossvein r-m close to base of cell dm. Because of these characters, this new species is placed in *Carreramyia*.

Distribution. Only known from Surinam.

Ecology. The holotype was collected in a malaise trap in secondary forest on moist clay soil in a former coffee- and cocoa plantation.

Carreramyia tigrina spec. nov.

Figs 17–22.

Studied type specimens. HOLOTYPE. Female. PERU, Madre de Dios, Rio Tambopata, Sachavacayoc Centre, 12°51'S-69°22'W, malaise trap, 16-26.X.2008, leg. J.T. Smit, coll. RMNH.

Description (based on holotype)

Adult female. Body size: 5 mm.

Head. Face occupying almost 3/5 of head width in frontal view; yellow with small black spot laterad of antennal fossa; short yellow pilose. Gena yellow; yellow pilose. Frons yellow, short black pilose. Vertex strongly produced medially; yellow with a wide black transverse fascia; black pilose, with pile getting longer posteriorly. Occiput yellow; black pilose dorsally, yellow pilose ventrally. Eye short pale pilose. Antennal fossa about as wide as high. Antenna with scape and pedicel black, basoflagellomere yellow except black on dorsobasal 2/5; ratio of scape:basoflagellomere approximately as 1:5.5; pedicel very short; basoflagellomere very long and wide; arista absent.

Thorax. Scutum black medially with two narrow submedian yellow vittae, widely pale yellow along margins; yellow pilose, except black pilose posterolaterally. Postpronotum pale yellow; bare. Postalar callus pale yellow; bristly black pilose. Scutellum yellow on median 1/3, black on lateral 1/3; yellow pilose, with some long and bristly pile posteriorly; sulcate posteromedially. Anepisternum a little convex, without sulcus; black, except pale yellow along posterior margin; long black pilose dorsally, shorter and partly pale pilose medially, bare on ventral 1/4. Katepisternum black, except for yellow macula at dorsal margin; short yellow pilose dorsally, bare ventrally. Katepimeron pale yellow; convex; bare. Calypter blackish. Halter whitish yellow.

Wing: hyaline, infuscated blackish around veins, with yellow transverse fascia between pterostigma and vein M, also yellow on and around vein CuA; microtrichose, except bare on 1st costal cell, basal 1/4 of cell br, posterobasal 1/3 of cell bm, anterobasal 1/4 of cell cup.

Legs: front and middle legs yellow, except basal 1/3 of femora blackish; hind femur blackish except yellow on subbasal 1/5, on narrow stripe dorsally and on postero-apical 1/3; hind tibia blackish except narrowly yellow at apex; hind tarsus yellow. Coxae and trochanters blackish; black pilose.

Abdomen. Tergite 1 black; other tergites yellow with black median vitta and widely black lateral margins; short black pilose. Tergite 2 wider than thorax, widest point at posterior 1/3. Sternites yellow, except for small dark lateral macula on tergite 5. All sternites short black pilose.

Etymology. The name *tigrina* (L., 'of tigers') is inspired by the pattern of black and buff spots and stripes on head, wings and abdomen.

Diagnosis. Distinguished from *C. megacephalus* and *C. megacera* by the striped pattern of the abdomen. From *C. flava* it differs by the mostly black scutum, the laterally black scutellum, the hind leg with its tibia wider than its femur, the absence of an arista in the female and the shape of the female basoflagellomere.

Notes. The male is unknown, so whether it has the furcate basoflagellomere characteristic for this genus or not can only be guessed at. Nevertheless the female is very similar to *C. megacephala* in the diagnostic and other important characters: basoflagellomere very long, vertex strongly produced, face very wide, scutellum sulcate, vein R4+5 without appendix, crossvein r-m close to base of cell dm. Therefore, this new species is placed in *Carreramyia*.

Unfortunately, after description and taking photographs, the holotype was severely damaged by accident. The head and a large part of the thorax are lost.

Distribution. Only known from Peru.

Ecology. The holotype was collected in a malaise trap at the edge of primary rainforest (varzea forest) in the Amazonian part of Peru.

***Ceratophya argentinensis* spec. nov.**

Figs 24–28.

Studied type specimens. HOLOTYPE. Female. ARGENTINA. Label 1: "Argentina, Tucuman / Rio Potrerillo / S26.80674° / W65.46934°, 969 m", other side of label: "01.XI.2008 / leg. T. Ekrem"; label 2: "309 / Y1008 / DNA voucher". Coll. RMNH.

Description (based on holotype)



Adult female. Body size: 7 mm.

Head. Face occupying 1/3 of head width in frontal view; sides converging ventrad; laterally depressed; pale yellow, with vague, narrow, brown median vitta; black pilose on dorsal half and along eye margins, yellow pilose on ventral half. Face in profile slightly convex, slightly produced downward at anterior oral margin. Gena yellow. Frons anteriorly yellow, black pilose; posteriorly black, yellow pilose. Vertex blackish brown, yellow pilose. Occiput blackish; white pilose. Eye bare. Antennal fossa about as wide as high. Antenna: scape and pedicel blackish, basoflagellomere pale brown; antennal ratio approximately 4:1:5.

Thorax. Scutum blackish, except narrowly yellow along anterior margin; short appressed golden yellow pilose, except short black pilose posteriad of transverse suture on lateral 1/3. Postpronotum yellow; yellow pilose. Postalar callus yellow; black pilose. Scutellum yellow, short yellow pilose; semicircular, without sulcus or calcars. Anepisternum not differentiated by sulcus; anterior half black, posterior half yellow; mixed black and golden pilose anterodorsally. Anepimeron black; black pilose on dorsal 3/4. Katepisternum blackish brown, except yellow along dorsal and ventral margins; black pilose dorsally and ventrally, widely bare in between. Katatergum and anatergum blackish brown; short microtrichose. Calypter grey. Halter yellow.

Wing: hyaline; microtrichose, except bare on 1st costal cell, basal 1/4 of cell r1, most of cell br except microtrichose along vena spuria, entirely on cell bm, basal 1/2 of cell cup, basal 1/10 of cell dm.

Legs: Front and hind legs yellow, slightly infuscated on basal 1/4 of femora and tarsi; pilosity short, mixed yellow and black. Coxae and trochanters blackish brown; yellow pilose.



FIGURES 22–31. 22. *Carreramyia tigrina* (holotype), scutellum. 23. *Carreramyia megacephalus* male (Costa Rica, coll. RMNH), genitalia lateral. 24–28. *Ceratophya argentinensis* female (holotype). 24. habitus dorsal; 25. habitus lateral; 26. head frontal; 27. head lateral; 28. wing. 29–31. *Ceratophya carinifacies* female (holotype). 29. habitus dorsal; 30. abdomen dorsal; 31. abdomen lateral.

Abdomen. Tergite 1 yellow anteriorly, dark brown posteriorly; yellow pilose laterally. Tergite 2 blackish brown with large, triangular marking posteromedially, which extends to anterior margin by narrow vitta; short black pilose, except bare on yellow parts; coarsely punctured over entire surface. Tergites 3 and 4 with colouration as tergite 2; bare, except tergite 4 short black pilose along lateral margin. Tergite 5 blackish, except narrowly yellow along lateral margin; short black pilose. In profile with tergite 4 almost perpendicular to tergite 2, so apex of abdomen curved downward. Sternites blackish anteriorly, yellow posteriorly; bare, except sternite 4 posteriorly and sternite 5 short black pilose.

Male. Unknown.

Diagnosis. Recognizable by the black posteromedian part of tergite 2 in combination with the unsulcate scutellum.

Distribution. Only known from northern Argentina (prov. Tucuman).

***Ceratophya carinifacies* (Curran, 1934)**

Figs 29–35.

Microdon carinifacies Curran, 1934: 376.

Studied type specimens. HOLOTYPE. Female. GUYANA. Label 1 (red): "*Microdon carinifacies* Curran Type"; label 2: "Trop. research station New York Zool. Society, No. 201330"; coll. AMNH. The specimen is in bad condition: ventral parts of the thorax are missing, as well as front legs and all tarsi, and the wings are badly damaged. Attached to the pin is also an empty puparium, from which the female holotype has apparently been reared.

Additionally studied specimens. BRAZIL: 1 female, RO Fazenda, Rancho Grande, 62 km s. Ariquemes, 10°18'S 62°53'W, 8–20.IV.1997, malaise trap, leg. A.C. Rhen & C. Alexander, coll. UCD.

Redescription (based on holotype)

Adult female. Body size: 8 mm.








Head. Face occupying 2/5 of head width in frontal view; parallel-sided; submedially depressed; pale yellow, with brown median vitta from antennal fossa to oral margin, where it is at its narrowest; entirely short, appressed yellow pilose. Face in profile straight, produced downward at anterior oral margin. Gena blackish. Frons brown, short pale pilose; vertex blackish brown, short white pilose. Occiput blackish. Eye bare. Antennal fossa about as wide as high. Antenna with scape basally yellow, gradually getting brown in apical 1/2; pedicel and basoflagellomere blackish brown; antennal ratio approximately 4:1:5.5. Basoflagellomere parallel-sided, with apex directed a little upward. Arista slender, about half the length of the basoflagellomere.

Thorax. Scutum blackish brown, except narrowly yellow along margins; short black pilose, except lateral fasciae of white pile along anterior margin and transverse suture, and a complete white pilose fascia between postalar calli. Postpronotum and postalar callus yellow; short yellow pilose. Scutellum yellow, short yellow pilose; semicircular, without sulcus or calcars. Anterior and posterior part of anepisternum not differentiated; short yellow pilose except on ventral 1/3. Anepimeron entirely white pilose. Katepisternum white pilose, at least dorsally (ventral part not visible in type specimen). Katatergum and anatergum microtrichose. Other pleurae bare. Calypter and halter yellow.

Wing: hyaline; microtrichose, except bare on costal cells, basal 1/2 of cell r1, basal 1/3 of cell r4+5, entirely on cell r-m except for traces of microtrichia around vena spuria, on cell bm, on basal 1/4 of cells DM and CuA1, on anterobasal 1/2 of cell cup.

Legs: Front legs and all tarsi missing in holotype. Mid- and hindfemora and -tibiae yellowish brown, mid-femora paler on apical half. Legs short and appressed pilose, black on femora, yellow on tibiae. Coxae and trochanters brownish, with pale pile.

Abdomen. Tergite 1 brownish. Tergite 2 brownish with large, oblique, lateral yellow markings over entire length, which leave narrow brown lateral margins and a large median brown triangle. Tergite 3 brownish with yellow lateral margins and a widely yellow posterior margin, which vaguely extends forward medially and gradually turns into pale brown. Tergite 4 mostly yellow, except brownish sublaterally. Tergite 5 yellowish brown. In profile with tergite 4 almost perpendicular to tergite 2, so apex of abdomen curved downward. Tergites short pilose, mostly black on brown parts and yellow on yellow parts. Sternites yellowish brown, short pale pilose.



FIGURES 32–41. 32–35. *Ceratophya carinifacies* female (holotype). 32. head frontal; 33. head lateral; 34. puparium lateral; 35. puparium dorsal. 36–41. *Ceratophya notata* male (holotype). 36. habitus dorsal; 37. habitus lateral; 38. head frontal; 39. head lateral; 40. wing; 41. scutellum.

Male. Unknown.

Puparium. Figs 34–35. Length about 8 mm., dorsally more or less flat, ventrally convex. Head skeleton and anterior parts (including anterior spiracles) lost. Posterior spiracle not visible.

Diagnosis. Recognizable by the yellow posteromedian part of tergite 2 in combination with the unsulcate scutellum.

Notes. The female from Brazil is darker in overall colouration: the scutellum has two blackish marks posteroventrally, tergites 3 and 4 are black with yellow posterior margins (tergite 3 also with yellow lateral margins).

Distribution. Known from Brazil and Guyana.

***Ceratophya longicornis* Wiedemann, 1824** (excluded from *Ceratophya*)

Ceratophya longicornis Wiedemann, 1824: 14. Type locality: Brazil. [type lost]

English translation of German description in Wiedemann (1830): "Antennae black, basally brown; third segment four times longer than first; face fawn-coloured ['rehhaarbräunlich']; frons black. Scutum black, with traces of fawn-coloured hairs along anterior margin, transverse suture, posterior margin and lateral margins. Abdomen black, second segment longer than in preceding species [*C. notata*], a little narrower than the other [segments], with 'longish', posteriorly forked ['hinten kurz gabelförmigen oder zweispitzigen'] yellow marking which does not reach the posterior margin. Base of abdomen ventrally and sternal margins widely yellow. Wing yellowish. Femora black, with narrowly yellow apex; tibiae and tarsi yellow.—In my collection, a specimen treated with arsenic-solution, because of which the colours are not well discernable."

Notes. According to Wiedemann (1830) the type was in his personal collection, which usually means that it is conserved in the NMW collection. However, no specimen recognizable as the type of *C. longicornis* is present in the NMW (pers. comm. P. Sehnal). It is not in the ZMHB collection either (pers. comm. J. Ziegler). Wiedemann (1830) wrote that the specimen had been treated with arsenic, a common practice in those days to prevent insect depredation of entomological collections. Sometimes arsenic solutions were applied in a mixture with other ingredients, e.g. soap or mercury (Albrecht 1993). Perhaps the chemical treatment of the specimen has eventually led to its disappearance.

The original description provides some indications that *C. longicornis* is quite different from other *Ceratophya* species:

- basoflagellomere four times longer than scape (in all other *Ceratophya* species the basoflagellomere is less than twice as long as the scape);
- the second tergite is longer than in *C. notata* and narrower than the other tergites (suggesting a constricted abdomen, which is found in no other *Ceratophya* species).

These characters indicate that *C. longicornis* is probably not a true *Ceratophya* as defined by Cheng and Thompson (2008) and Reemer and Ståhls (2013a). At present the taxonomic affinities of *C. longicornis* remain unclear. However, there are few Neotropical species of Microdontiinae with such an antennal ratio combined with a constricted abdomen, so possibly the identity of *C. longicornis* can be clarified later.

***Ceratophya notata* Wiedemann, 1824**

Figs 36–41, 55.

Ceratophya notata Wiedemann, 1824: 14.

Studied type specimens. HOLOTYPE. Male. BRAZIL. Label 1: "Brasilia, coll. Winthem"; label 2: "*notata*, det. Wiedem."; label 3: "*notata* Wied. Brasilia"; label 4 (yellow): "Lectotype *Ceratophya notata* Wied. Desig. Thompson 1977". Coll. NMW.

Additionally studied material. SURINAM: 1 female, Zanderij, 18-21.VII.1964, leg. D.C. Geijskes, coll. RMNH.

Adult male. Body size: 7 mm.

Head. Face parallel-sided, occupying 1/3 of head width in frontal view; laterally depressed; pale yellow, with

brown median vitta from antennal fossa to oral margin, where it is at its narrowest; entirely short yellow pilose. Face in profile straight, produced downward at anterior oral margin. Gena brown. Oral cavity with lateral margins not produced. Frons blackish brown, short white pilose; vertex blackish, short white pilose. Occiput blackish brown. Eye bare. Antennal fossa about as wide as high. Antenna with scape yellow, gradually getting brown in apical 1/3, pedicel and scape brown; antennal ratio approximately 6:1:9. Basoflagellomere curled upward in apical 1/3. Arista slender, a little longer than half the length of the basoflagellomere.

Thorax. Scutum blackish brown; short black pilose, except lateral fasciae of white pile along transverse suture and posteriorly between postalar calli. Postpronotum yellow, short yellow pilose. Postalar callus pale brown, short black pilose. Scutellum yellow, short yellow pilose; slightly sulcate apicomediaally; without calcars. Anterior and posterior part of anepisternum not differentiated; short yellow pilose except on ventral 1/3. Anepimeron entirely white pilose. Katepisternum white pilose. Katatergum and anatergum microtrichose. Other pleurae bare. Calypter and halter yellow.

Wing: hyaline; microtrichose, except bare on costal cell, anteriorly along vein RS, on cell br except for traces of microtrichia around vena spuria, on cell bm, on anterobasal 1/3 of cell cup.

Legs: Brown, with tarsi and apical 1/3 of pro- and mesotibiae yellow. Legs short yellow pilose. Coxae and trochanters brownish, with yellow to white pile.

Abdomen. Blackish brown, with large, oblique, lateral yellow markings over entire length of tergite 2, and yellow posterior margins of tergites 3 and 4. In profile with tergite 4 almost perpendicular to tergite 2, so apex of abdomen curved downward. Tergites 1 and 2 short black pilose, except yellow pilose on yellow markings. Tergites 3 and 4 entirely short yellow pilose. Sternites brown, short pale pilose. Male genitalia as in fig. 55.

Female. Same as male, except the following differences in the Abdomen. about 1.5 times wider than in male, with posterior margins of tergites strongly extending over next tergites, suggesting 'telescopic' capacities. The yellow posterior margins of tergites 3 and 4 are somewhat swollen.

Diagnosis. Recognizable by the following combination of characters: scutellum weakly sulcate, male: tergite 3 blackish brown with posterior margin broadly yellow, female: tergite 2 with two oblique yellow vittae, tergites 3 and 4 with yellow posterior margins.

Notes. The holotype of *C. notata* carries the label "Lectotype *Ceratophya notata* Wied. Design. Thompson 1977". However, assuming that this specimen indeed was used by Wiedemann to describe the species, and considering that there seems to have been only one specimen on which he based his description, it seems that this specimen should be regarded as the holotype.

Distribution. Known from Brazil and Surinam.

***Ceratophya panamensis* (Curran, 1930)**

Figs 42–47, 56.

Microdon panamensis Curran, 1930: 6.

Studied type specimens. HOLOTYPE. PANAMA. Male. Label 1: "France Field, Canal Zone. 18.I.1929"; label 2 (red): "Type *Microdon panamensis* Curran"; label 3: "Collector C.H. Curran". Coll. AMNH.


PARATYPE.—Female. Attached to same pin as male holotype, with which it has been collected 'in coitu' (Curran 1930).

Notes on type specimens. The male holotype and the female paratype are attached to the same pin. There is no question as to which of the specimens should be regarded as holotype, because Curran (1930) clearly designated the male as such. Unfortunately, the abdomen of the male is missing. Nonetheless, its genitalia are conserved in a microvial attached to the pin.

Adult male. Body size: 7–8.5 mm (Curran 1930).

Head. Face parallel-sided, occupying slightly less than 1/3 of head width in frontal view; sublaterally depressed, so medially slightly carinate; pale yellow, with brown median vitta from base of antenna to oral margin, where it is at its narrowest; entirely short yellow pilose. Face in profile straight, produced downward at anterior oral margin. Gena black. Frons black, short yellow pilose; vertex black, short yellow pilose. Occiput black. Eye bare. Antennal fossa about as wide as high. Antenna with scape brown, yellowish basally; pedicel and scape

brown; antennal ratio approximately 5:1:8. Basoflagellomere parallel-sided, curled slightly upward in apical 1/4. Arista slender, about half the length of the basoflagellomere.



FIGURES 42–54. 42–47. *Ceratophya panamensis*. 42. male holotype (above) & female paratype (photo: American Museum of Natural History); 43. female, head frontal; 44. female, head lateral; 45. male, wing; 46. male, scutellum; 47. male, hind tarsus. 48–54. *Ceratophya scolopus* male (holotype). 48. habitus dorsal; 49. habitus lateral; 50. abdomen posterodorsal; 51. head frontal; 52. wing; 53. scutellum; 54. apex of hind tibia with basitarsus, lateral.

Thorax. Scutum black; short black pilose, except fasciae of pale pile, which are badly visible in type specimen. Curran (1930): 'golden hair forming three bands, the anterior one situated on the anterior margin, interrupted, the median one narrowest and entire, posterior band widest, situated on the posterior border, irregularly margined in front'. Postpronotum yellow, short yellow pilose. Postalar callus brown, short black pilose. Scutellum yellow, short yellow pilose, except short black pilose apicoventrally; deeply sulcate apicomediaally; without calcars. Anterior and posterior part of anepisternum not differentiated; short yellow pilose except on ventral 1/3. Anepimeron entirely white pilose. Katepisternum white pilose dorsally and ventrally, these patches widely separated. Katatergum and anatergum microtrichose. Other pleurae bare. Calypter pale brown, halter yellow.

Wing: hyaline with brown veins; microtrichose, except bare on costal cells, on basal 1/4 of cell r1, on cell br except for traces of microtrichia around vena spuria, on most cell bm except antero-apically, on anterobasal 1/3 of cell cup.

Legs: Profemora and -tibiae reddish brown, with apical 1/3 of tibiae paler. Meso- and metafemora and -tibiae blackish brown. Femora very short blackish pilose; tibiae very short appressed golden pilose, most densely so on apical 1/3 of pro- and mesotibiae. Procoxae yellow and white pilose, other coxae brown and white pilose. Trochanters brown, short pale pilose.

Abdomen. Missing in type specimen. Curran (1930): 'Abdomen brownish black, with yellow markings. Second segment on either side with a large pale triangle which is continuous with a broad pale vitta on the third segment, the very broad apex of the third segment, except at the sides, yellowish, the base more or less yellow; fourth segment with the posterior border broadly yellow. Pile very short, golden yellow, on the fourth segment less abundant and more brassy. Second to fourth sternites brownish yellow with pale-yellow apices.' Genitalia as in fig. 56.

Female. Same as male, except the following differences. Face occupying slightly more than 1/3 of total head width in frontal view. Scutum and pleurae brownish. Postalar callus pale brown. Wing: veins yellow. Abdomen about 1.5 times wider than in male; tergites and sternites strongly overlapping and with posterior margins of tergite 3 and 4 appearing swollen; entirely yellow, except narrowly brown along lateral margins of tergites.

Diagnosis. Recognizable by the following combination of characters: scutellum deeply sulcate, male: tergite 3 blackish brown with posterior margin broadly yellow and with yellow lateral vittae (based on description of Curran 1930; in holotype the abdomen is missing), female: tergites yellow, only brownish along lateral margins.

The colouration of the abdomen appears to be strongly sexually dimorphic: mostly blackish in the male, mostly pale orange in the female.

Distribution. Only known from Panama.

***Ceratophya scolopus* (Shannon, 1927)**

Figs 48–54, 57.

Microdon scolopus Shannon, 1927: 20. Type locality: Amazon region.

Microdon (Ubristes) scolopus Shannon in Thompson *et al.* (1976)

Studied type specimens. HOLOTYPE. Male. Label 1 (small, round, red-bordered): "Holotype"; label 2: "Amazon. 66 53"; label 3: "*Microdon scolopus* Snn.". Coll. BMNH.

Adult male. Body size: 7 mm.

Head. Face occupying about 1/3 of the head width in frontal view; parallel-sided; laterally weakly depressed; pale yellow, with blackish median vitta from antennal fossa to oral margin, where it is at its narrowest; entirely short yellow pilose. Face in profile straight, produced downward at anterior oral margin. Gena black. Frons and vertex black, short pilose. Occiput black. Eyes bare. Antennal fossa about as wide as high. Antenna brown, except scape yellowish basally; antennal ratio approximately 5:1:8; arista slender, slightly longer than half the length of the basoflagellomere.

Thorax. Scutum black; short pilose. Postpronotum and postalar callus brown (perhaps yellow in better preserved specimens); short pilose. Scutellum yellow, with basal 1/3 and apical 1/4 blackish; apicomediaally sulcate. Anterior and posterior part of anepisternum not differentiated; short pale pilose except on ventral 1/3. Anepimeron entirely white pilose. Katepisternum white pilose dorsally (ventral part not visible in type specimen). Calypter greyish brown. Halteres missing in type specimen.

Wing: hyaline; microtrichose, except bare on costal cells, on basal 1/3 of cell r1, on cell br except for traces of microtrichia around vena spuria, on cell bm, posterobasal 1/3 of cell r4+5, on basal 1/6 of cells DM and CuA1, on anterobasal 1/2 of cell cup.

Legs: Blackish, with tarsi of pro- and mesolegs yellow. Hind leg with first tarsomere black, second tarsomere yellow (other tarsomeres missing in type specimen). First tarsomeres of all tarsi with strong basoventral tooth. Coxae and trochanters brownish, with pale pile.

Abdomen. Blackish brown, with yellow posterior margin of tergite 4; mostly whitish pilose. In profile with tergite 4 almost perpendicular to tergite 2. Sternites 1-3 brown, sternite 4 yellow; short pale pilose. Genitalia as in fig. 57.

Female. Unknown.

Diagnosis. This species is unique among *Ceratophya*-species in the presence of a strong basoventral tooth on the first tarsomeres of all tarsi. Furthermore, it's the only known *Ceratophya*-species with an almost black abdomen (except for the yellow posterior margin of tergite 4).

Notes. When Shannon (1927) described this species, he mentioned the similarity in general appearance to the stingless-bee mimics which he described in the subgenus *Ubristes*. Nonetheless, he did not place it in *Ubristes*, as Thompson *et al.* (1976) did. Examination of the type revealed that it has all the characters of the genus *Ceratophya*. The specimen is a little dirty and greasy, so colours and pilosity are not always easy to assess.

Distribution. Only known from the holotype from the Amazon region, probably from Brazil.

***Hypselosyrphus amazonicus* Reemer, 2013**

Figs 58–66, 108.

Hypselosyrphus amazonicus Reemer, in Reemer & Ståhls 2013: 132. Replacement name.

Microdon (Ubristes) scutellaris Shannon, 1927: 20. Preoccupied by Schummel (1842).

Studied type specimens. HOLOTYPE. BRAZIL. Female. Label 1 (small, round, red-bordered): "Holotype"; label 2: "Amazon. 6653"; label 3: "*Microdon Ubristes scutellaris* Snn.". Leg.: H.W. Bates (Shannon 1927), coll. BMNH.

Additionally studied specimens: PERU: 1 male, Madre de Dios, Rio Tambopata, Sachavacayoc Centre, 12°51'S–69°22'W, mal. trap 26–28.X.2008, leg. J.T. Smit, coll. RMNH.


Redescription (based on holotype)

Adult female Body size: 8 mm.


Head. Face occupying 1/5 of head width in frontal view; black; entirely with long white pilosity, with lateral 1/4 along eye margins white pollinose. Gena hardly developed. Oral cavity directly bordering eye margins; with lateral margins not produced. Frons black; white pilose. Vertex more or less convex, dull black; white pilose anteriorly, black pilose posteriorly; ocellar triangle not elevated compared to rest of vertex. Occiput black; very narrow, barely visible in lateral view; with anterior row of dorsally orientated black pile on dorsal 1/3; with posterior row of posteriorly orientated pale pile on ventral 2/3. Eye entirely with short, pale pile, a little longer than ommati diameter. Antennal fossa about as wide as high. Antenna yellow; antennal ratio 4:1:? (basoflagellomeres missing in holotype; Shannon 1927 gives an antennal ratio of 1:0.25:1).

Thorax. Dark brown. Scutum densely black pilose, except for a few white pile along transverse suture. Postpronotum and postalar callus black pilose. Scutellum equilaterally triangular, with posterior corner quite blunt; directed upward, making an angle with the scutum of about 40°; paler brown than scutum; long black pilose anteriorly and dorsally, long pale pilose posteroventrally. Anepisternum a little convex, no clear division between anterior and posterior part; anterior part black pilose, posterior part bare. Anepimeron entirely black pilose. Katepisternum black pilose dorsally; with a few pile ventrally. Katatergum long microtrichose. Anatergum short microtrichose. Other pleurae bare. Calypter grey, halter yellowish.


Wing: hyaline, subtly tinged with brown; with vague, dark transverse fascia on anterior half just before middle of wing, and with vague, whitish transverse fascia just after middle of wing. Microtrichose, except for most of cell br (except for traces of microtrichia along vena spuria), posterobasal 1/2 of cell bm, anterobasal 2/5 of cell cup, and on basomedian 1/2 of alula.




55



56



57



58



59



60



61



62

FIGURES 55–62. 55–57. *Ceratophya*, male genitalia. 55. *C. notata* (holotype); 56. *C. panamensis* (holotype); 57. *C. scolopus* (holotype). 58–62. *Hypselosyrphus amazonicus* male (Peru, coll. RMNH). 58. habitus dorsal; 59. habitus lateral; 60. head frontal; 61. head lateral; 62. wing.

Legs: Blackish brown, except for whitish yellow last three tarsomeres of front- and hindlegs (mid-tibia and -tarsus missing in holotype); black pilose, except pale pilose on pale tarsomeres. Hind-tibia with greatest width apically of middle; about as wide as hind femur; with strong excavation at cicatrice (lateral view); pilosity about as long as width of tibia. Hind-basitarsus enlarged; about twice as wide as tibia in dorsal view. Coxae and trochanters brownish black, with black pile.

Abdomen. Oval, 1.5 times as wide as thorax, with largest width at tergite 3 (which only slightly narrows posteriorly). Yellowish brown; with wide dark lateral margins, dark posterior margins of tergites 2, 3 and 4, and a narrow median vitta; tergite 5 yellow with dark median vitta. All sternites yellow and yellow pilose.

Male. As female. The studied male is darker in overall colouration than the female holotype, but possibly the holotype has lost some of its colour over time.

Diagnosis. This is the only known species of *Hypselosyrphus* with a triangular scutellum.

Notes. The name *Microdon scutellaris* Shannon, 1927 is a junior primary homonym of *Microdon scutellaris* Schummel, 1842. In such cases it is a good custom to name the species after the person who described it first. In this case however, the name *shannoni* is preoccupied too by Curran (1940), so another name was chosen: *amazonicus*, referring to the apparent distribution of the species. This name was introduced by Reemer in Reemer and Ståhls (2013a).

Distribution. Known from Brazil and Peru.

Hypselosyrphus anax (Thompson, 1976)

Figs 67, 109.

Microdon analis Curran, 1940: 3. Preoccupied by Macquart, 1842, new name *anax* introduced by Thompson *et al.* (1976).

Studied type specimens. HOLOTYPE. BRAZIL. Male. Label 1: "Brasilien, Nova Teutonia, 27°11' B, 52°23'L, Fritz Plaumann, 10.12.1937"; label 2 (red): "Microdon analis Curran Holotype". Coll. AMNH.

Redescription (based on holotype)

Adult male Body size: 8.5 mm.

Head. Face occupying 1/5 of head width in frontal view; black; entirely with long white pilosity except for black pile ventrolaterad of antennal fossa, with lateral strips of white pollinosity along eye margins. Gena black. Oral cavity with lateral margins not produced. Frons black; black pilose. Vertex black; black pilose; in profile almost vertical directly anterior of anterior ocellus. Occiput black; with anterior row of dorsally orientated black pile on dorsal 1/3; with posterior row of posteriorly orientated pale pile over entire length. Eye entirely with short, pale pile, a little longer than ommati diameter. Antennal fossa about as wide as high. Antenna yellow; antennal ratio 4:1:? (basoflagellomere missing in holotype).

Thorax. Black, a little brownish on pleurae, posterior callus and scutellum. Scutum densely black pilose, except for a few white pile along transverse suture. Postpronotum and postalar callus black pilose. Scutellum directed upward, a little sulcate posteriorly; entirely long black pilose. Anepisternum a little convex, no clear division between anterior and posterior part; anterior part black pilose, posterior part black pilose along posterior margin. Anepimeron entirely black pilose. Katepisternum black pilose dorsally. Katatergum and anatergum pilose and microtrichose, respectively. Other pleurae bare. Calypter dark grey, halter pale brown.

Wing: hyaline, tinged with brown on basal 2/3, tinged yellowish on apical 1/3. Microtrichose, except for posterobasal 1/3 of cell br, anterobasal 1/3 of cell bm, anterobasal 1/6 of cell cup, anterior to vein A2 and on a narrow basal strip on the alula.

Legs: Femora dark brown, black pilose; metafemur gradually paler towards apex. Front- and mid-tibia brown, black pilose, except for short yellow pile on posteroapical 1/3 of front-tibia. Hind-tibia with greatest width apically of middle; about as wide as hind femur; yellow, entirely with long (a little longer than maximal width of tibia) black pile. Tarsi yellow; yellow pilose, except first tarsomere of mid-leg. Coxae and trochanters brownish black, with black pile.

Abdomen. Blackish brown, except tergite 4 with widely yellow posterior and posterolateral margins. First tergite laterally with black pile, sublaterally with white pile. Tergites 2 and 3 black pilose. Tergites 3 and 4 not fused. Tergite 4 black pilose, except yellow pilose laterally and posteriorly. Sternites 1-3 blackish brown; sternite 4 yellow, except blackish brown anteriorly and laterally. Sternite 1 posteriorly with black pile; sternites 2 and 3 black

pilose; sternite 4 yellow pilose on yellow parts, black pilose on dark parts. Pregenital segments yellowish brown. Genitalia as in fig. 109.

Female. Unknown.

Diagnosis. A mostly black species with partly bare alula, apicomediaally sulcate scutellum and basoflagellomere maximally half as long as scape. Very similar to *H. plaumanni*. Differences are: tergite 4 with posterior margin widely yellowish, sternite 4 largely yellowish, scutellum entirely black pilose, wing tinged yellowish on apical third, otherwise brownish.

Notes. Curran (1940) made a mistake in the key in which he separates his *M. analis* from *M. plaumanni*: he states that the posterior femora are tawny pilose on the basal half in *M. plaumanni*, while the posterior femora are black pilose in *M. analis*. This character applies to the posterior tibiae, not to the femora.

Distribution. Only known from Brazil.

***Hypselosyrphus corbiculipes* Papavero, 1962**

Figs 68–69.

Hypselosyrphus corbiculipes Papavero, 1962: 320.

Studied type specimens. HOLOTYPE. BRAZIL. Female. Label 1: "Rio de Janeiro D.F. Sumaré, Werner, Alvarenga, 1.55"; label 2: "28.632"; label 3 (red): "Holotipo"; label 4: "Hypselosyrphus corbiculipes sp. n. N. Papavero det. 1962". Coll. MZUSP.

Redescription (based on holotype)

Adult female Body size: 7 mm.

Head. Face occupying 1/4 of head width in frontal view; shining blackish brown; with long white pilosity and white pollinosity on lateral 1/3. Gena undeveloped, oral cavity directly bordering eye margins, with lateral margins not produced. Frons shining dark brown, pale pilose, with small spot of white pollinosity along eye margin. Vertex convexly produced, shining dark brown; brown pilose; ocellar triangle not elevated compared to rest of vertex. Occiput brownish black; narrow, only slightly widened dorsally; pale pilose; entirely pale pollinose; with anterior row of dorsally orientated short black pile on dorsal 1/2; with posterior row of posteriorly orientated pale pile over entire length; black pollinose on dorsal half, white pollinose on ventral half. Eye entirely with short, pale pile, about as long as ommati diameter. Antennal fossa about as wide as high. Antenna pale brown; antennal ratio 4:1:4. Basoflagellomere with acute apex; with small sensory pit slightly beyond half the segment. Arista pale, about as long as basoflagellomere.

Thorax. Dark brown. Postpronotum, scutum, postalar callus and scutellum black pilose, except for a few pale pile along transverse suture and on postalar callus. Scutellum apicomediaally sulcate, without calcars; directed upward, making an angle with the scutum of about 45°. Anepisternum a little convex, no clear division between anterior and posterior part; anterior and posterior part black pilose, widely bare in between. Anepimeron entirely black pilose. Katepisternum black pilose dorsally; bare ventrally. Katatergum long microtrichose. Anatergum short microtrichose. Other pleurae bare. Calypter greyish brown, halter yellow with dark grey knob.

Wing: hyaline, except blackish on anterobasal 2/3; microtrichose, except on 1st costal cell, anterobasal 1/2 of 2nd costal cell, basal 1/10 of cell r1, basal 1/4 of cell br, posterobasal 2/3 of cell bm, anterobasal 1/4 of cell cup.



Legs: yellowish brown, except tarsi and hind tibia yellow; black pilose, except yellow pilose on hind tibia, hind tarsus and apical three tarsomeres of other tarsi. Metatibiae strongly widened, with greatest width slightly apically of middle, about 1.5 times as wide as posterior femur at largest width; with strong excavation at cicatrice (lateral view); pilosity about half as long as width of tibia. Hind-basitarsus enlarged; almost twice as wide as apex of metatibia in dorsal view. Coxae and trochanters brown, with black pile.

Abdomen. Oval, wider than thorax, with largest width at posterior margin of tergite 2; entirely shining brown; white pilose. All sternites shining brown; pale pilose.

Male. unknown.

Diagnosis. Recognized by the following combination of characters: scutellum sulcate, alula entirely microtrichose, hind tibia yellow and yellow pilose.

Distribution. Only known from Brazil.



FIGURES 63–71. 63–66. *Hypselosyrphus amazonicus* female (holotype). 63. habitus dorsal; 64. habitus lateral; 65. scutellum; 66. wing. 67. *Hypselosyrphus anax* male (holotype), habitus dorsal. Photo: American Museum of Natural History. 68–69. *Hypselosyrphus corbiculipes* female (holotype). 68. habitus dorsal; 69. habitus lateral. 70–71. *Hypselosyrphus helvus* female (holotype). 70. habitus dorsal; 71. habitus lateral.

***Hypselosyrphus helvus* spec. nov.**

Figs 70–73.

Studied type specimens. HOLOTYPE. BRAZIL. Female. Label 1: "Brasil: Roraima / Rio Uraricoera / Ilha de Maraca"; label 2: "19-24-vii-1987 / J.A. Rafael / L.S. Aquino"; label 3: "Armadilha de / malaise"; label 4 (yellow): "INPA / Rafael 1989". Coll. INPA (when studied the specimen was on loan in the USNM-collection).

Redescription (based on holotype)

Adult female Body size: 10 mm.

Head. Face occupying 1/4 of head width in frontal view; yellow; entirely yellow pilose. Gena hardly developed. Oral cavity directly bordering eye margins; with lateral margins not produced. Frons yellow, except black around lunula; yellow pilose, except black pilose around lunula. Vertex convexly produced, shining blackish brown; yellow pilose; ocellar triangle not elevated compared to rest of vertex. Occiput very narrow; black on dorsal half, yellow on ventral half; entirely yellow pilose. Eye entirely with short, white pile, a little longer than ommati diameter. Antennal fossa a little higher than wide. Antenna yellowish; antennal ratio approximately as 4:1:3.

Thorax. Entirely yellow and yellow pilose. Scutellum semicircular; with posterior margin apicomediaally very faintly slightly sulcate; pale. Anepisternum convex, without sulcus; anterior part pilose, posterior part bare. Anepimeron entirely pilose. Katepisternum pilose dorsally; bare ventrally. Katatergum long yellow microtrichose. Anatergum short microtrichose. Other pleurae bare. Calypter and halter yellow.

Wing: hyaline; with veins yellow and anterior half tinged with yellow. Microtrichose, except bare on 1st costal cell, basal 1/10 of 2nd costal cell, basal 1/2 of cell br, posterobasal 1/3 of cell bm, anterobasal 1/6 of cell cup, basomedian 1/4 of alula.

Legs: Yellow; yellow pilose; hind tibia strongly widened and with pile as long as width of tibia.

Abdomen. Widest point at posterior 1/3 of tergite 2. Tergites and sternites (including sternite 1) entirely yellow and yellow pilose.

Male. unknown.

Etymology. The Latin adjective *helvus* (bay, yellow) was chosen because of the entirely yellow thorax and abdomen of this species.

Diagnosis. This is the only *Hypselosyrphus*-species with entirely yellow thorax and abdomen and with hyaline wings.

Distribution. Only known from Brazil (Roraima).

***Hypselosyrphus marshalli* spec. nov.**

Figs 74, 75.

Studied type specimens. HOLOTYPE. ECUADOR. Female. Label 1: "ECU: Napo, Tiputini / Biodiversity Stn., 0°36'50"S. / 76°9'1"W, May 2011, S.A. / Marshall, debu00339701"; label 2: "Voucher code M. Reemer / MR398". Coll. PUCE.

Additionally studied specimens. 1 male was photographed by S.A. Marshall at same locality and date as holotype (fig. 75), but not collected.

Description (based on holotype)

Adult female Body size: 9 mm.

Head. Face occupying between 1/5 and 1/4 of head width in frontal view; shining black; entirely white pilose. Gena hardly developed. Oral cavity directly bordering eye margins; with lateral margins not produced. Frons shining black, short white pilose. Vertex strongly produced, shining black; black pilose. Occiput black; narrow; black pilose on dorsal half, white pilose on ventral half; entirely pollinose. Eye entirely with short, pale pile, about as long as ommati diameter. Antennal fossa about as wide as high. Antenna blackish brown; antennal ratio 5:1:6. Arista blackish brown, about as long as basoflagellomere.

Thorax. Black, except postalar callus brownish. Scutum densely black pilose, except for medially interrupted transverse fasciae of long yellowish pile along suture. Postpronotum and postalar callus black pilose. Scutellum apicomediaally sulcate, without calcars; directed upward, making an angle with the scutum of about 50°; black pilose, except yellowish pilose ventrally. Anepisternum a little convex, no clear division between anterior and

posterior part; anterior part black pilose, posterior part with a few black pile along posterior margin. Anepimeron entirely black pilose. Katepisternum black pilose dorsally; bare ventrally. Katatergum long microtrichose. Anatergum short microtrichose. Other pleurae bare. Calypter blackish, halter pale yellow.

Wing: basal and apical 1/3 somewhat infuscated, middle 1/3 yellowish white; microtrichose, except bare on 1st costal cell, basal 1/20 of cell r1, basal 3/4 of cell br, posterior 1/3 of cell bm, basal 1/6 of cell cup.

Legs: Black, except fore- and middle-tarsi yellow and apical three tarsomeres of hindleg yellow; black pilose, except yellow pilose on apical three tarsomeres. Hind tibia strongly widened, with greatest width at apical 1/3, a little wider than posterior femur at largest width; with strong excavation at cicatrice (lateral view); pilosity about as long as width of tibia. Hind-basitarsus enlarged; about twice as wide as apex of metatibia in dorsal view. Coxae and trochanters black, with black pile.

Abdomen. More or less oval, wider than thorax, with largest width at posterior 1/3 of tergite 2; black, except tergite 4 posteriorly with pair of submedian dark yellowish vittae of half the length of the tergite, and tergite 5 almost entirely dark yellowish (only black along anterior margin). Tergites black bilose, except tergite 2 with pair of narrow yellow pilose transverse fasciae along posterior margin, tergite 4 white pilose posteriorly, tergite 5 white pilose. Sternites black except sternite 5 yellowish; black pilose, except posterior margin of tergite 4 and tergite 5 entirely yellow pilose.

Male (based on photograph). As female, except for usual sexual dimorphism and following differences. Wing largely infuscated, except for yellow pterostigma and yellow patch between pterostigma and vein R4+5.

Etymology. This species is attributed to Stephen A. Marshall, who collected the holotype, after having made excellent photographs of living specimens. A photograph of the living holotype was also included in his book *Flies* (Marshall 2012: 428).

Diagnosis. Recognized by the following combination of characters: scutellum sulcate, alula with small bare patch basomedially, thorax and abdomen largely black, basoflagellomere about as long as scape or longer.

Distribution. Only known from Ecuador.

***Hypselosyrphus maurus* spec. nov.**

Figs 76–82, 110.

Studied type specimens. HOLOTYPE. FRENCH GUYANA. Male. Label 1: "FRENCH GUIANA / Kaw Mountains / 4°32.893'N 52°10.245'W / Leg. V. Soon 29.12.2002. Coll. RMNH.

PARATYPE. FRENCH GUYANA: 1 female, Kaw Road, PK 37, Relais Patawa, N 4°32'42" / W 52°9'9", IX.2008 (malaise trap), leg. O. Morvan, coll. RMNH.

Additionally studied specimens. PERU: 1 female, Madre de Dios, Rio Tambopata, Sachavacayoc Centre, 12°51'S-69°22'W, malaise trap, 4-10.IX.2009, leg. J.T. Smit, coll. RMNH.

Redescription (based on holotype)

Adult male Body size: 7 mm.

Head. Face occupying 1/5 of head width in frontal view; shining black; long black pilose on lateral 1/4, also with some white pile ventrolaterally; grey pollinose on lateral 1/4. Gena hardly developed. Oral cavity directly bordering eye margins; with lateral margins not produced. Frons shining black, dark pilose. Vertex strongly produced, shining black; black pilose. Occiput black; narrow; with anterior row of dorsally orientated short black pile on dorsal 1/2; with posterior row of posteriorly orientated pale pile over entire length; entirely pollinose. Eye entirely with short, pale pile, about as long as ommati diameter. Antennal fossa about as wide as high. Antenna brown; antennal ratio 4:1:4. Basoflagellomere with acute apex; with small sensory pit at half the length of the segment. Arista pale, about as long as basoflagellomere.

Thorax. Black. Scutum densely black pilose, except for medially interrupted transverse fasciae of shorter white pile along suture. Postpronotum and postalar callus black pilose. Scutellum apicomediaally sulcate, without calcars; directed upward, making an angle with the scutum of about 30°; black pilose. Anepisternum a little convex, no clear division between anterior and posterior part; anterior part black pilose, posterior part with a few black pile along posterior margin. Anepimeron entirely black pilose. Katepisternum black pilose dorsally; bare ventrally. Katatergum long microtrichose. Anatergum short microtrichose. Other pleurae bare. Calypter dark grey, halter brown with knob blackish.



FIGURES 72–75. 72–73. *Hypselosyrphus helvus* female (holotype). 72. head frontal; 73. head lateral. 74–75. *Hypselosyrphus marshalli* (photographs by S.A. Marshall). 74. female (holotype), habitus; 75. male, habitus.



FIGURES 76–84. 76–80. *Hypselosyrphus maurus* male (holotype). 76. habitus dorsal; 77. habitus lateral; 78. head frontal; 79. head lateral; 80. wing. 81–82. *Hypselosyrphus maurus* female (paratype). 81. habitus dorsal; 82. habitus lateral. 83–84. *Hypselosyrphus pingo* female (holotype). 83. habitus dorsal; 84. habitus lateral.

Wing: hyaline, veins darkened around stigmal crossvein, veins around pterostigma yellow; microtrichose, except on 1st costal cell, basal 1/2 of 2nd costal cell, basal 1/10 of cell r1, entirely on cell br except microtrichose along vena spuria, posterior 1/2 of cell bm, basal 1/3 of cell cup.

Legs: Black, except fore- and middle-tarsi yellow and apical four tarsomeres of hindleg yellow; black pilose, except yellow pilose on apical two tarsomeres. Hind tibia strongly widened, with greatest width at apical 1/3, about 1.5 times as wide as posterior femur at largest width; with strong excavation at cicatrice (lateral view); pilosity about half as long as width of tibia. Hind-basitarsus enlarged; about 1.5 times as wide as apex of metatibia in dorsal view. Coxae and trochanters blackish, with black pile.

Abdomen. More or less oval, wider than thorax, with largest width at posterior 1/3 of tergite 2; blackish brown. Tergite 1 shining, tergite dull except for shining median 1/3, tergite 3 dull except shining along lateral margins, tergite 4 shining. Tergites black bilose, except tergite 1 and posterior and lateral margins of tergite 4 white pilose. Sternite 1 white pilose, other sternites black pilose. Genitalia as in fig. 110.

Female. As male, except for usual sexual dimorphism. In the paratype, the colouration of the wing veins is entirely uniform (not dark around stigmal crossvein and yellow around pterostigma). In the additionally studied female from Peru, however, the colouration of the wing veins is as in the male holotype. These differences are considered to be intraspecific.

Etymology. The specific epithet *maurus* (Latin for 'dark') refers to the black appearance of this species.

Diagnosis. 7–8 mm. Recognized by the following combination of characters: scutellum sulcate, alula entirely microtrichose, hind tibia brown and black pilose, abdomen black.

Distribution. Known from French Guyana and Peru.

Hypselosyrphus pingo spec. nov.

Figs 83–89, 111.

Studied type specimens. HOLOTYPE. BRAZIL. Female. Label 1: "Brasilien / Nova Teutonia / 27°11'B. 52°23'L / Fritz Plaumann / I 1971 / 300 . 500 m"; label 2 (red): "HOLOTYPE / *Hypselosyrphus pingo* / Reemer". Coll. ZMAN.

PARATYPES: BRAZIL: 1 male [head missing], Nova Teutonia, 27°11'B. 52°23'L, XI.1968, leg. F. Plaumann, coll. RMNH; 1 female, Nova Teutonia, 27°11'B. 52°23'L, 20.II.1937, leg. F. Plaumann, coll. CNC; 1 female, Nova Teutonia, 27°11'B. 52°23'L, I.1968, leg. F. Plaumann, coll. USNM.

Redescription (based on holotype)

Adult female Body size: 10 mm.


Head. Face occupying 1/4 of head width in frontal view; black; entirely white pilose. Gena hardly developed. Oral cavity directly bordering eye margins; with lateral margins not produced. Frons black; sparsely black pilose medially, white pilose along lateral margins. Vertex convexly produced, shining black; black pilose; ocellar triangle not elevated compared to rest of vertex. Occiput black; very narrow, barely visible in lateral view; dorsally black pilose, except for small patch of yellowish white pile next to vertex; ventrally white pilose. Eye entirely with short, pale pile, a little longer than ommati diameter. Antennal fossa a little higher than wide. Antenna black, except basoflagellomere blackish brown; antennal ratio approximately as 5:1:2.

Thorax. Scutum, postpronotum and postalar callus pale brown; densely yellowish brown pilose. Scutellum trapezoid, with posterior margin apicomediaally slightly sulcate; pale brown; yellowish brown pilose. Anepisternum convex, without sulcus; dark brown; anterior part yellow pilose, posterior part black pilose. Anepimeron entirely yellow pilose. Katepisternum dark brown; yellow pilose dorsally; bare ventrally. Katatergum long black microtrichose. Anatergum short microtrichose. Other pleurae bare. Calypter and halter yellowish.

Wing: with dark brown fascia halfway wing, as wide as 1/5 of length of wing, and equally wide dark brown fascia at apex, with yellow fascia in between and also yellow between wing base and first brown fascia; colours most clear on anterior half, fading posteriorly. Microtrichose, except bare on 1st costal cell, basal 1/4 of cell br, basal 1/10 of cell bm.

Legs: Yellow; yellow pilose, except black pilose on basal 1/3 of hind femur.

Abdomen. More or less oval, but apical segments narrower than basal ones; a little wider than thorax, widest at posterior 1/4 of tergite 2. Tergite 1 yellowish brown. Tergite 2 yellowish brown with narrow dark brown median line and narrowly dark brown posterior margin. Tergite 3 dark brown, tergites 4 and 5 blackish. Tergites entirely yellow pilose. Basal sternites pale brown, apical sternites darker; yellow pilose.



FIGURES 85–89. 85–87. *Hypselosyrphus pingo* female (holotype). 85. head frontal; 86. head lateral; 87. wing. 88–89. *Hypselosyrphus pingo*. 88. female (paratype), habitus dorsal; 89. male (paratype), habitus dorsal. 90. *Hypselosyrphus plaumanni* male (holotype), habitus dorsal. Photo: American Museum of Natural History. 91–95. *Hypselosyrphus pseudorhoga* female (holotype). 91. habitus dorsal; 92. habitus lateral; 93. head frontal; 94. head lateral; 95. wing.

Male. In the only known male specimen the head is lost. Otherwise, this specimen agrees with the female, except in that all tergites are yellowish brown, except for three black vittae on tergite 4. Genitalia as in fig. 111.

Etymology. The Latin verb *pingo* means to colour or to paint. The name refers to the painted wings of the species. As a species name it is to be treated as a noun in apposition.

Diagnosis. The wing marks and the unwidened hind-tibiae of this species immediately distinguish it from other known *Hypselosyrphus*-species, except *H. vexillipennis*. From that species it differs by the completely yellowish brown pilose scutum and scutellum (black pilose in *H. vexillipennis*).

Notes. The colouration of the tergites seems to be sexually dimorphic: mostly yellow with three black vittae on tergite 4 in the male, mostly dark on tergites 3 and 4 in female. In two of the paratypes, the median dark fascia on the wing extends to the wing base, so the wing is dark brown on the basal 3/5 of the wing. In this species the hind tibiae are not widened or corbiculate, unlike in other species of *Hypselosyrphus* (except *H. vexillipennis*) and *Stipomorpha*. It is a matter of taste whether this species should be considered as a mimic of stingless bees or not. Nevertheless, it possesses all characters described as diagnostic for *Hypselosyrphus*.

Distribution. Only known from Brazil (Nova Teutonia).

Hypselosyrphus plaumanni (Curran, 1940)

Fig 90.

Microdon plaumanni Curran, 1940: 3.

Studied type specimens. HOLOTYPE. BRAZIL. Female. Label 1: "*Microdon plaumanni* Curran Holotype"; label 2: "Holotype"; label 3: "Brasilien, Nova Teutonia, 27°11' B, 52°23' L, Fritz Plaumann, 15.2.1937". Coll. AMNH.

Additionally studied specimens. BRAZIL: 1 male, Nova Teutonia, 27°11' S, 52°23' W, 300-500 m, 1968, leg. F. Plaumann, coll. RMNH; 2 males, same locality, 12-1970, leg. F. Plaumann, coll. ZMAN; 1 male, same locality, XI.1966, leg. F. Plaumann, coll. CNC; 1 female, E. Rio, Itatiaya, 700 m., 2.XI.1939, leg. J.F. Zikán, coll. USNM.

Redescription (based on holotype)

Adult female Body size: 9 mm.

Head. Face occupying 1/5 of head width in frontal view; black; entirely with long white pilosity except for black pile ventrolaterad of antennal fossa, with lateral strips of white pollinosity along eye margins. Gena black. Oral cavity with lateral margins a little produced. Frons black; black pilose medially, white pilose laterally. Vertex black; black pilose; in profile almost vertical directly anterior to anterior ocellus. Occiput black; with anterior row of dorsally orientated black pile on dorsal 1/3; with poster row of posteriorly orientated pale pile over entire length. Eye entirely with short, pale pile, a little longer than ommati diameter. Antennal fossa about as wide as high. Antenna yellow; antennal ratio 4:1:2; basoflagellomere parallel-sided with narrowly rounded apex, with large oval sensory pit located approximately in middle, occupying about 1/3 of height of basoflagellomere. Arista slender, slightly longer than length of basoflagellomere.

Thorax. Black, a little brownish on pleurae, along posterior margin of scutum and scutellum. Scutum densely black pilose, except for patches of white pile along transverse suture and along posterior margin. Postpronotum and postalar callus black pilose. Scutellum directed upward, deeply sulcate posteriorly, thus leaving two large 'mamiform' processes; long black pilose, except white pilose ventrally. Anepisternum a little convex, no clear division between anterior and posterior part; anterior part black pilose, posterior part black pilose along posterior margin. Anepimeron entirely black pilose. Katepisternum black pilose dorsally. Katatergum and anatergum long and short microtrichose, respectively. Other pleurae bare. Calypter dark grey, halter pale brown.

Wing: hyaline, tinged with brown over most of surface, less so posteriorly and apically. Microtrichose, except for basal 1/8 of cells BM and R.

Legs: Front- and mid-femora dark brown, black pilose; hind-femora brown with apical 1/2 yellow, with long black pile. Front- and mid-tibiae yellowish brown, black pilose. Hind-tibia strongly widened, with greatest width in the middle; about 1.5 times as wide as posterior femur; yellow, with long (about as long as maximal width of tibia) yellow pile on basal 1/2, long black pile on apical 1/2. Tarsi yellow, dorsally black pilose on basal two tarsomeres, dorsally yellow pilose on apical three tarsomeres; ventrally with dense, short, yellow pilosity. Coxae and trochanters brownish black, with black pile.

Abdomen. Blackish brown. First tergite laterally white pilose. Second tergite white pilose, except for black pile along extreme lateral margins. Tergite 3 short black pilose over most of surface, with longer white pile posterolaterally. Tergite 4 deeply emarginated along posterior margin; black pilose, except white pilose along posterior and lateral margins.; tergite 5 anteriomedially with convex bulge, which fits into the posterior emargination of tergite 4; white pilose. Sternite 1 with long white pile; other sternites with long black pile.

Male. As female, except for usual sexual differences. Genitalia (not drawn) almost identical to those of *M. analis*.

Diagnosis. A mostly black species with partly bare alula, apicomediaally sulcate scutellum and basoflagellomere maximally half as long as scape. Very similar to *H. anax*. Differences are: tergite and sternite 4 entirely blackish brown, scutellum black pilose, except pale pilose posteroventrally, wing entirely tinged brownish.

Notes. Placed as a synonym of *Ubristes flavitibia* Walker by Thompson *et al.* (1976), but here transferred to *Hypselosyrphus* and reinstated as valid species.

Distribution. Only known from Brazil.

***Hypselosyrphus pseudorhoga* spec. nov.**

Figs 91–95.

Studied type specimens. HOLOTYPE. PERU. Label 1: “Quincemil / Peru 24/31 X / 1962 / L.E. Pena”; label 2 (red): “CNC / Ottawa”; label 3: “Pseudorhoga / new genus! / n. sp.!”. Coll. CNC.

Redescription (based on holotype)

Adult female Body size: 7 mm.

Head. Face occupying about 1/5 of head width in frontal view; yellow; entirely yellow pilose. Gena hardly developed. Oral cavity directly bordering eye margins; with lateral margins not produced. Frons yellow; yellow pilose. Vertex flat; shiny dark brown with faint metallic hues; yellow pilose. Occiput black; entirely yellow pilose. Eye bare. Antennal fossa about as high as wide. Antenna yellow; antennal ratio approximately as 2.5:1:5.

Thorax. Mesoscutum yellow with four blackish maculae anterolaterally and posterolaterally, with wide, vague, brownish median vitta; entirely yellow pilose. Scutellum semicircular; shining blackish brown with faint metallic hues; yellow pilose. Postpronotum and posterior callus yellow; yellow pilose. Pleurae yellow and yellow pilose. Anepisternum convex, without sulcus; anterior and posterior parts pilose. Anepimeron entirely pilose. Katepisternum bare. Katatergum long yellow microtrichose. Anatergum short microtrichose. Other pleurae bare. Mediotergite blackish. Calypter and halter yellow.

Wing: hyaline. Microtrichose, except bare on 1st costal cell, basal 2/3 of 2nd costal cell, basal 1/5 of cell r1, along anterior and posterior margins of cell br, posterobasal 3/4 of cell bm, anterobasal 1/2 of cell cup.

Legs: Yellow, except hind tibia dark brown with yellow apices. All legs yellow pilose, except hind tibia with long black pile on dark parts.

Abdomen. Widest point at posterior margin of tergite 2. Tergites yellow, with oval dark macula medially on tergite 2 and three dark vittae on tergites 3-5; yellow pilose. Sternites yellow; yellow pilose, except sternite 1 bare.

Male. Unknown.

Etymology. Chris Thompson coined the name *pseudorhoga* because of the *Rhoga*-like appearance of this species, as evoked by the dark maculae on the mesoscutum and the three dark vittae on the abdomen. The name is to be treated as a noun in apposition.

Diagnosis. No other known species of *Hypselosyrphus* has a basoflagellomere that is twice as long as the scape. The abdominal colour pattern is also characteristic: yellow with three dark vittae. The unproduced vertex is only shared with *H. ulopodus*, from which it differs by the two characters just mentioned.



Notes. Only a female of this species is known. As this differs from other species of *Hypselosyrphus* in the unproduced vertex and the short scape, its placement in this genus is very tentative and preliminary. The phylogenetic affinities of this species should be revisited once additional specimens have been found, based on molecular studies and the male genitalia.

Distribution. Only known from Quince Mil in southern Peru, a place in the foothills of the Andes at around 600-700 m. above sea level.

***Hypselosyrphus trigonus* Hull, 1937**

Figs 96–99.

Hypselosyrphus trigonus Hull, 1937: 21.



FIGURES 96–99. 96–99. *Hypselosyrphus trigonus* female (holotype). 96. habitus dorsal; 97. habitus lateral; 98. head frontal; 99. head lateral. **100–104.** *Hypselosyrphus ulopodus* male (holotype). 100. habitus dorsal; 101. habitus lateral; 102. head frontal; 103. head lateral; 104. wing. **105–106.** *Hypselosyrphus vexillipennis* female (holotype). 105. habitus dorsal; 106. habitus lateral.

Studied type specimens. HOLOTYPE. PANAMA. Female. Label 1: "Canal Zone: Barro, Colorado. 16-VII-1924. N. Banks."; label 2 (red.): "M.C.Z. Type31169"; label 3 (large, red-bordered): "*Hypselosyrphus trigoniformis* Hull, F.M.H.". Coll. MCZ.

Additionally studied specimens. BELIZE. Female. No further data. Coll. RMNH. PANAMA: 1 female, Chiriqui, 15 km NW Hato del Volcan, 1200 m, 24-31.V.1977, Peck & Howden, coll. CNC.

Note on holotype: The holotype is labelled as *Hypselosyrphus trigoniformis* Hull, but this must be a mistake, because Hull has not described a species under that name. He did, however, describe *Hypselosyrphus trigonus*, the description of which agrees well with this specimen.

Redescription (based on holotype)

Adult female Body size: 7 mm.

Head. Face occupying 1/5 of head width in frontal view; shining blackish brown; with long white pilosity on lateral 1/3; with white pollinosity on lateral 1/5. Gena hardly developed. Oral cavity directly bordering eye margins; with lateral margins not produced. Frons shining dark brown, dark pilose, except white pilose directly along eye margin. Vertex convexly produced, shining dark brown; black pilose; ocellar triangle not elevated compared to rest of vertex. Occiput black; narrow; with anterior row of dorsally orientated short black pile on dorsal 1/2; with posterior row of posteriorly orientated pale pile over entire length; black pollinose on dorsal half, white pollinose on ventral half. Eye entirely with short, pale pile, about as long as ommati diameter. Antennal fossa about as wide as high. Antenna pale brown; antennal ratio 4:1:4. Basflagellomere with acute apex; with small sensory pit at half of the segment, situated in a wide groove from base of arista to near apex. Arista pale, about as long as basoflagellomere.

Thorax. Dark brown. Scutum densely black pilose, except for transverse fasciae of pale pile along suture and along posterior margin. Postpronotum black pilose, postalar callus white pilose. Scutellum apicomediaally sulcate, with convex posterior margin, without calcars; directed upward, making an angle with the scutum of about 45°; long white pilose anteriorly, long black pilose posteriorly, long white pilose posteroventrally. Anepisternum a little convex, no clear division between anterior and posterior part; anterior part black pilose, posterior part with a few black pile along posterior margin. Anepimeron entirely black pilose. Katepisternum black pilose dorsally; with a few pile ventrally. Katatergum long microtrichose. Anatergum short microtrichose. Other pleurae bare. Calypter grey, halter brown.

Wing: hyaline, vaguely darkened around stigmal crossvein, and vaguely yellowishly tinged around and posteriad of pterostigma (best seen before a dark background); microtrichose, except on 1st costal cell, basal 2/5 of 2nd costal cell, basal 1/10 of cell r1, basal 2/5 of cell br, posterobasal 2/5 of cell bm, basal 2/5 of cell cup.

Legs: Brown, except fore- and middle-tarsi yellow and apical four tarsomeres of hindleg yellow; black pilose, except yellow pilose on apical three tarsomeres. Metatibiae strongly widened, with greatest width slightly apically of middle, about 1.5 times as wide as posterior femur at largest width; with strong excavation at cicatrice (lateral view); pilosity about half as long as width of tibia. Hind-basitarsus enlarged; about 1.5 times as wide as apex of metatibia in dorsal view. Coxae and trochanters brown, with black pile.

Abdomen. Oval, wider than thorax, with largest width at posterior margin of tergite 2. Colours hard to assess in type specimen because of partial folding and darkening of the segments. Hull (1937) states: "Abdomen short, oval, flat, vitreous, dark reddish brown on first, second and narrowly on base of third segment. Remainder bright orange." Sternite 1 black and black pilose. Other sternites yellow and yellow pilose.

Male. Unknown.

Diagnosis. Recognized by the following combination of characters: scutellum sulcate, alula entirely microtrichose, hind tibia brown and black pilose, abdomen orange to reddish brown.

Distribution. Known from Belize and Panama.

***Hypselosyrphus ulopodus* (Hull, 1944)**

Figs 100–104, 112.

Ubristes ulopodus Hull, 1944: 34.

Studied type specimens. HOLOTYPE. PERU. Female. Label 1: "Lachoruria, Putmayo Distr. PERU. 17-20 Aug.

1920"; label 2: "Cornell Univ. Expedition. Lot 569"; label 3 (red): "Holotype ulopodus Hull"; label 4: "Holotype Ubristes ulopodus Hull"; label 5 (red): "Holotype Cornell U. No. 2196". Coll. CU.

Additionally studied specimens. PARAGUAY. 1 male. Label 1: "Paraguay, Vezenyi"; label 2: "Asuncion, 1904.x.5"; label 3: "Coll. Mus. Nat. Hung."; label 4: "Ceratophya analis Curr. male, det. v. Doesburg". Coll. RMNH.

Redescription (based on holotype)

Adult female Body size: 8.5 mm.

Head. Face occupying 1/3 of head width in frontal view; yellow, except narrowly black laterally on dorsal half and with vague brown median vitta on ventral half; yellow pilose laterally, black pilose medially, with pile longer and denser around oral margin; with narrow lateral strips of white pollinosity along eye margins. Gena brownish. Oral cavity with lateral margins a slightly produced and anterior margin slightly notched. Frons black, except yellow posterior to lunula; black pilose. Vertex black; black pilose. Occiput black; black pilose dorsally, white pilose ventrally. Eye entirely with dense, dark pile, a little longer than ommati diameter. Antennal fossa about as wide as high. Antenna dark brown; antennal ratio approximately as 4:1:3.5; basoflagellomere parallel-sided with narrowly rounded apex, with small sensory pit located at 2/5 from base. Arista slender, slightly shorter than length of basoflagellomere.

Thorax. Blackish brown. Scutum entirely with long, erect black pile; with two submedian and two lateral vittae of greyish pollinosity. Postpronotum and postalar callus black pilose. Scutellum semicircular, without calcars; long black pilose anteriorly, long yellow pilose posteriorly. Anepisternum a little convex, with very slight sulcus between anterior and posterior part; black pilose, with large bare medioventral area. Anepimeron entirely black pilose. Katepimeron pilose. Katepisternum black pilose dorsally, bare ventrally. Katatergum and anatergum long and short microtrichose, respectively. Other pleurae bare. Calypter and halter dark greyish brown.

Wing: hyaline with yellow veins; tinged with brown in costal cells. Microtrichose, except for basal 1/4 of cell br and basal 1/3 of cell bm.

Legs, including coxae and trochanters, blackish brown, except apical four tarsomeres brownish yellow; entirely with long, black pile.

Abdomen. Brown. Tergites 3-5 fused, sutures not visible. Tergites black pilose laterally, pale yellow pilose medially, except tergite 4 entirely black pilose. All sternites with long black pile.

Male. (based on 1 specimen in coll. RMNH) Differs from female in the following: face black; antennal ratio approximately 4:1:2.5; tarsi entirely brownish yellow; abdomen blackish brown with posterior margin of tergite 4 yellow. Genitalia as in fig. 112.

Diagnosis. This is the only known species of *Hypselosyrphus* with a pilose katepimeron and a ventrally widened occiput.

Notes. This species is aberrant from its congeners because of the pilose katepimeron, the ventrally widened occiput and the unproduced vertex (the latter character is only shared with *H. pseudorhoga* spec. nov.). A phylogenetic analysis based on morphological characters placed the species as a sister of the clade containing *Rhoga* and *Hypselosyrphus* (Reemer and Ståhls 2013b). The relationships within this clade are considered not well enough established to introduce another generic name, hence the current assignment of this species.

Distribution. Known from Peru and Paraguay.

***Hypselosyrphus vexillipennis* spec. nov.**

Figs 105–107.


Studied type specimens. HOLOTYPE. BRAZIL. Female. Label 1: "1.I.1955 / Barueri / S. Paulo / 3409"; label 2: "K. Lenko leg.". Coll. USNM.

PARATYPES. BRAZIL. Female. Label 1: "BRASIL Rio de Janeiro / D.F. Corcovado / XI.1957 / Seabra e Alvarenga". Coll. MZUSP.


Female. Label 1: "Barueri, / São Paulo, Brasil / 15.I.1966 / K. Lenke col.". Coll. MZUSP.

Redescription (based on holotype)


Adult female Body size: 10 mm.




107




108




109



110



111



112

FIGURES 107–112. 107. *Hypselosyrphus vexillipennis* female (holotype), head frontal. 108–112. *Hypselosyrphus*, male genitalia. 108. *H. amazonicus* (Peru. coll. RMNH); 109. *H. anax* (holotype); 110. *H. maurus* (holotype); 111. *H. pingo* (paratype); 112. *H. ulopodus* (holotype) (dashed lines indicate missing part in specimen).

Head. Face occupying 1/5 of head width in frontal view; black; entirely white pilose. Gena hardly developed. Oral cavity directly bordering eye margins; with lateral margins not produced. Frons black; sparsely black pilose

medially, white pilose along lateral margins. Vertex convexly produced, shining black; black pilose; ocellar triangle not elevated compared to rest of vertex. Occiput black; very narrow, barely visible in lateral view; dorsally black pilose; ventrally white pilose. Eye entirely with short, pale pile, a little longer than ommati diameter. Antennal fossa a little higher than wide. Antenna black, except basoflagellomere blackish brown; antennal ratio approximately as 5:1:2.5.

Thorax. Scutum, postpronotum and postalar callus dark brown; densely black pilose. Scutellum trapezoid, with posterior margin apicomediaally slightly sulcate; dark brown; black pilose. Anepisternum convex, without sulcus; dark brown; anterior part and posterior margin black pilose. Anepimeron entirely black pilose. Katepisternum dark brown; black pilose dorsally; bare ventrally. Katatergum long black microtrichose. Anatergum short microtrichose. Other pleurae bare. Calypter brown, halter yellowish white.

Wing: dark brown on basal 3/5 and apical 1/5, with yellow fascia of 1/5 of wing length in between, somewhat infuscated around vein DM-Cu; colours most clear on anterior half, fading posteriorly. Microtrichose, except bare on 1st costal cell, basal 1/6 of cell br, basal 1/10 of cell bm.

Legs: Brown; black pilose, except yellow pilose on apical four tarsomeres of front and mid tarsus.

Abdomen. More or less oval, but apical segments narrower than basal ones; a little wider than thorax, widest at posterior 1/4 of tergite 2. Tergites dark brown; black pilose, except tergites 1, 5 and posterolateral corners of tergite 4 yellow pilose. Basal sternites pale brown, apical sternites darker; yellow pilose.

Male. Unknown.

Etymology. The name *vexillipennis* (flag-winged) refers to the painted wings of this species.

Diagnosis. The wing marks and the unwidened hind-tibiae of this species immediately distinguish it from other known *Hypselosyrphus*-species, except *H. pingo*. From that species it differs by the completely black scutum and scutellum (yellowish brown pilose in *H. vexillipennis*).

Notes. Compared with the holotype, the paratype is darker, almost black, in overall colouration. In this species the hind tibiae are not widened or corbiculate, unlike in other species of *Hypselosyrphus* (except *H. pingo*) and *Stipomorpha*. It is a matter of taste whether this species should be considered as a mimic of stingless bees or not. Nevertheless, it possesses all characters described as diagnostic for *Hypselosyrphus*.

Distribution. Only known from Brazil (São Paulo & Rio de Janeiro).

***Mermerizon inbio* Reemer, 2013**

Figs 113–117, 124.

Studied type specimens. HOLOTYPE. COSTA RICA. Male. Label 1: "COSTA RICA. Prov. Guanacaste, P.N. / Rincón de la Vieja, Send. a las aguas / termales, 900-1000 m, 6-7 OCT / 2001. D. Briceño, Red con Aguamiel. / L_N_305843_392970 #64950"; label 2: "INB0003380896 / INBIOCRI COSTA RICA"; label 3 (red): "Ultimo espécimen en / BD A. Lépez / 2-7-2002" / other side: "?MCR-25". Coll. INBIO.

Redescription (based on holotype)

Adult male Body size: 7.5 mm.

Head. Face occupying about 1/4 of head width in frontal view; yellow; yellow pilose, with narrow bare median line on dorsal half. Gena yellow. Frons black; yellow pilose laterally, black pilose posteriorly. Vertex dark yellow, except black at and around ocellar triangle; black pilose. Occiput black, except yellow posteriorly of vertex; black pilose on dorsal half, yellow pilose on ventral half. Eye bare. Antennal fossa about as high as wide. Antenna with scape dark brown, pedicel and basoflagellomere yellowish brown; antennal ratio approximately as 4:1:4.

Thorax. Scutum blackish brown, except yellow on notopleuron and around postpronotum and postalar callus; black pilose. Postpronotum, postalar callus and scutellum yellow; black pilose. Scutellum semicircular, without calcar, Anepisternum blackish brown; convex, without sulcus; black pilose on anterior part and along posterior margin, widely bare in between. Anepimeron brown; black pilose on dorsal 1/4. Katepisternum yellow dorsally, brown ventrally; bare. Katepimeron yellow. Katatergum long black yellow microtrichose. Anatergum short pale microtrichose. Calypter blackish. Halter yellowish brown.

Wing: hyaline; microtrichose, except bare on 1st costal cell, basal 1/4 of cell br, basal 1/3 of cell bm, anterobasal 1/4 of cell cup.

Legs: Front and mid legs yellowish brown; black pilose. Hind leg blackish brown, except basal 1/2 of tibia and

apical four tarsomeres yellowish brown. Front and mid coxae and trochanters yellowish brown; yellow pilose apically. Hind coxa and trochanter dark brown; black pilose.

Abdomen. Tergites and sternites yellowish; yellow pilose, except sternite 1 bare. Genitalia as in fig. 124.

Female. Unknown.

Diagnosis. Distinguished from the other two known species of *Mermerizon* by the black pilose mesoscutum.

Distribution. Only known from Costa Rica.

***Mermerizon mellosus* spec. nov.**

Figs 118–120, 125.

Studied type specimens. HOLOTYPE. COSTA RICA. Male. Label 1: "COSTA RICA. Guan. / 3 km SE R. Naranjo / 4-6 Aug 1993 / F.D. Parker"; label 2: "Ubristes / sp [male symbol] / det.: M. Hauser 2007". Coll. INBIO. Redescription (based on holotype)

Adult male Body size: 8.5 mm.

Head. Face occupying about 1/4 of head width in frontal view; yellow; yellow pilose, with narrow bare median line on dorsal half. Gena yellow. Frons black medially, yellow laterally; yellow pilose. Vertex yellow, except black at and around ocellar triangle; black pilose. Occiput yellow posteriad of vertex, black on dorsolateral half, yellow on ventral half; black pilose anteriorly on dorsal half, yellow pilose posteriorly on both dorsal and ventral half. Eye bare. Antennal fossa about as high as wide. Antenna yellow, with scape darker dorsally; antennal ratio approximately as 4:1:4.

Thorax. Scutum blackish brown dorsally, with margins widely yellow; entirely yellow pilose. Postpronotum, postalar callus and scutellum yellow; yellow pilose. Scutellum semicircular, without calcars. Pleurae yellow, except anepisternum brownish anterodorsally and katepisternum brownish dorsally. Anepisternum convex, without sulcus; yellow pilose on anterior part and along posterior margin, widely bare in between. Anepimeron yellow pilose on dorsal 1/4, with pile along dorsal margin blackish. Katepisternum bare. Katatergum long black microtrichose. Anatergum short pale microtrichose. Calypter and halter yellow.

Wing: hyaline; microtrichose, except bare on 1st costal cell, basal 1/10 of cell br, basal 1/4 of cell bm, anterobasal 1/10 of cell cup.

Legs: Yellow and yellow pilose, except: hind tibia brown on apical 1/4 and black pilose on apical 1/3; first tarsomere of hind tarsus blackish and black pilose. Coxae and trochanters yellow; yellow pilose.

Abdomen. Tergites yellow, with vague, narrow, brown vitta medially; yellow pilose. Sternites yellow; yellow pilose, except sternite 1 bare. Genitalia as in fig. 125.

Female. Unknown.

Etymology. The specific epithet *mellosus* is a Latin adjective meaning 'honey-coloured'.

Diagnosis. Differs from both other known species of *Mermerizon* by its entirely yellow femora.

Distribution. Only known from Costa Rica.

***Mermerizon mesmerizus* spec. nov.**

Figs 121–123, 126.

Studied type specimens. HOLOTYPE. ARGENTINA. Male. Label 1: "ARGENTINA. Catamarca / Prov., 9 km N La Merced. / 28°06.43'S-65°36.96'W / Mal. trap in damp ravine / 24.X-12.XI.2003. 1041 m / M.E. Irwin & F.D. Parker". Coll. RMNH (preliminary deposition, to be deposited in appropriate Argentinian collection later).

PARATYPES: Two males from same locality and date as holotype. One in coll. RMNH, one in coll. CSCA.

Description (based on holotype, unless stated otherwise)

Adult male Body size: 5 mm.

Head. Face occupying about 1/3 of head width in frontal view; yellow; yellow pilose. Gena yellow. Frons and vertex brown; yellow pilose. Occiput brown; yellow pilose dorsally, white pilose ventrally. Eye bare. Antennal fossa about as high as wide. Antenna black; antennal ratio approximately as 2:1:4.



FIGURES 113–123. 113–117. *Mermerizon inbio* male (holotype). 113. habitus dorsal; 114. habitus lateral; 115. head frontal; 116. head lateral; 117. wing. 118–120. *Mermerizon mellosus* male (holotype). 118. habitus dorsal; 119. habitus lateral; 120. head frontal. 121–123. *Mermerizon mesmerizus* male (holotype). 121. habitus dorsal; 122. habitus lateral; 123. head frontal.

Thorax. Scutum black, with margins widely yellow; entirely yellow pilose. Postpronotum and postalar callus yellow; yellow pilose. Scutellum semicircular, without calcars; yellow, except black ventrolaterally; yellow pilose. Anepisternum convex, without sulcus; brown anteriorly, yellow posteriorly; yellow pilose anterodorsally and posterodorsally. Anepimeron brown with yellow margins; yellow pilose. Katepisternum brown; yellow pilose dorsally. Katepimeron yellow; bare. Katatergum yellow; long yellow microtrichose. Anatergum yellow; short yellow microtrichose. Calypter grey. Halter yellow.

Wing: hyaline, with costal cell grey and with vaguely defined greyish transverse fasciae from pterostigma to bm-cu, from appendix of R4+5 to dm-cu and at M1; microtrichose, except bare on 1st costal cell, basal 2/3 of cell br and narrowly along anterior margin of cell cup.

Legs: [Front legs missing in holotype. Description partly based on paratypes.] Femora black, except narrowly yellow at apex; black pilose, except partly yellow pilose basally. Tibiae yellow, except hind femur black at apical 1/3; yellow pilose. Tarsi black; black pilose. Coxae and trochanters blackish brown; pale pilose.

Abdomen. Tergites brown, with lateral and posterior margins a little paler. Tergites 1 and 2 yellow pilose. Tergite 3 black pilose on anterior 2/3, yellow pilose on posterior 1/3. Tergite 4 black pilose on anterior 1/3 and laterally, yellow pilose medially and posteriorly. Sternites brownish. Sternite 1 bare. Sternite 2 yellow pilose. Sternites 3-4 black pilose. Genitalia as in fig. 126.

Female. Unknown.

Etymology. The specific epithet *mesmerizus* is a Latinized adjective derived from the English 'mesmerizing', which means 'hypnotizing' in the sense of 'fascinating'.

Diagnosis. Body size 5-6.5 mm. Unlike the other two known species of *Mermerizon*, this is not clearly a stingless bee mimic. The pilosity of the hind tibia is short and appressed. The wings have a pattern of vaguely defined greyish transverse fasciae.

Distribution. Only known from northern Argentina.

***Stipomorpha apicula* (Curran, 1930)**

Fig 127, 230.

Microdon apiculus Curran, 1930: 5.

Studied type specimens. HOLOTYPE. PANAMA. Male. Label 1 (red): "*Microdon apicula* Curran Type"; label 2: "Barro Colo Isld., Canal Zone, 1-7-1929"; label 3: "Collector C.H. Curran". Coll. AMNH.


Redescription (based on holotype)

Adult male. Body size: 8 mm.


Head. Face occupying 1/5 of head width in frontal view; shining pale yellow with black median stripe which gradually narrows from entire width of face at level of antennae down to 1/6 of width of face at oral margin; face with white pilosity on yellow parts and just below antennae, bare on median stripe. Gena brown. Oral cavity with produced brown lateral margins and notched anterior margin. Frons and lunula black and short black pilose, except for bare triangular part posterior to lunula. Vertex black; short black pilose. Occiput black; black pilose on dorsal half, white pilose on ventral half. Eye bare. Antennal fossa about as wide as high. Antenna blackish brown, scape and pedicel black pilose; antennal ratio 5:1:5; basoflagellomere parallel-sided with narrowly rounded apex, with sensory pit located at 3/5 from base, within a vague groove that ranges from a little ventrad of base of arista to a little beyond sensory pit; arista slender, about 2/3 of length of basoflagellomere.

Thorax. Black, but paler on posteroventral part of anepisternum, posterior part of anepimeron and katepimeron. Colour of pilosity on dorsal part hard to assess because of glue that has spread over it; Curran (1930) describes this pilosity as follows: "Pile black, an undulate anterior band on the mesonotum, a small spot at the inner ends of the suture, a very broad prescutellar band and the scutellum wholly, golden-reddish pilose." Scutellum semicircular, without calcars. Anepisternum more or less flat, pilose anterodorsally. Anepimeron pilose posterodorsally. Katepisternum and katepimeron bare. Calypter blackish brown, halter pale brown.


Wing: hyaline, with pterostigma and surrounding veins pale yellow; microtrichose, except bare near the junction of veins R1 and RS, on basal 1/3 of cell br, posterobasal 1/3 of cell bm, anterobasal 1-5 of cell cup, basomedian 1/6 of alula.



124



125



126



127



128



129

FIGURES 124–129. 124–126. *Mermerizon*, male genitalia. 124. *M. inbio* (holotype); 125. *M. mellosus* (holotype); 126. *M. mesmerizus* (holotype). 127. *Stipomorpha apicula* male (holotype), habitus dorsal. 128–129. *Stipomorpha crematogastri* female (holotype). 128. habitus dorsal; 129. habitus lateral.

Legs: Brownish black, except last two tarsomeres of each leg yellow. Femora black pilose. Front tibia white pilose; mid-tibia white pilose on basal half, black pilose on apical half. Hind-tibia white pilose on basal 1/3, black pilose on apical 2/3. Tarsi dorsally black pilose, with some yellow pile intermixed on last two tarsomeres. Tarsi ventrally with short, dense, appressed yellow pile. Coxae and trochanters blackish brown, black pilose, anterior coxa also with some yellow pile.

Abdomen. Orange brown. Second tergite wider than thorax, widest point at half the length; third and fourth tergites strongly narrowing. Tergite 1 yellow pilose laterally, but black pilose along extreme lateral margin; with anterolateral 'ridges'; with anteromedian smooth, concave area. Tergite 2 yellow pilose, but black pilose along extreme lateral margin. Tergite 3 sparsely whitish yellow pilose. Tergite 4 quite densely yellow pilose, more whitish anterolaterally; with two oval marks of greyish pollinosity on the anterior half of the tergite. Sternite 1 bare, separated from sternite 2 by a wide membrane. Other sternites sparsely pale pilose. Genitalia as in fig. 230.

Female. According to Curran (1930) there is a female paratype. This has however not been studied.

Diagnosis. Distinguishable from other known species by the black legs and thorax in combination with the orange brown abdomen.

Stipomorpha crematogastris spec. nov.

Figs 130–133.

Studied type specimens. HOLOTYPE. BRAZIL. Female. Label 1: "BRAZIL: Mato Grosso / 12° 50'S., 51° 47'W. / 8-IV 1968 / O.W. Richards."; label 2: "R.S. & R.G.S. / Expedition / B.M. 1968-260"; label 3: "Dry / forest"; label 4: "bred nest / Crematogaster"; with empty puparium on piece of dry leaf attached to pin. Coll. BMNH.

PARATYPE. BRAZIL. Female. Label 1: "BRAZIL: Mato Grosso / 12° 50'S., 51° 47'W. / 26-III 1968 / O.W. Richards."; label 2: "R.S. & R.G.S. / Expedition / B.M. 1968-260"; label 3: "Dry / forest"; label 4: "Roadside"; label 5: "on leaf [illegible]"; label 6: "Microdon (Ubristes) / ? lanei Curran / N.P. Wyatt det. 1985". Coll. BMNH.

Description (based on holotype)

Adult female. Body size: 10 mm.

Head. Face occupying almost 1/2 of head width in frontal view; shining yellow; yellow pilose. Lateral oral margins produced. Frons yellow; yellow pilose laterally. Vertex yellow; yellow pilose, except for transverse fasciae of black pile anteriorly and posteriorly. Occiput yellow; yellow pilose. Eye bare. Antennal fossa about as wide as high. Antenna pale brown. Antennal ratio approximately as 5:1:7; basoflagellomere parallel-sided with rounded apex, with sensory pit at around 2/5 from base. Arista slender, almost as long as basoflagellomere.

Thorax. Scutum black dorsally, with margins widely yellow; yellow pilose, except for four patches of black pile: two anteriorly and two posteriorly of transverse suture. Postpronotum, postalar callus and scutellum yellow and yellow pilose. Anepisternum weakly convex, without distinction between anterior and posterior part; yellow; yellow pilose anteriorly; yellow pilose along posterior margin. Anepimeron yellow; yellow pilose dorsally, bare ventrally. Katatergum and anatergum yellow; short and long microtrichose, respectively. Katepimeron yellow; bare. Katepisternum yellow; bare. Calypter and halter yellow.

Wing: hyaline, tinged with yellow, especially anteriorly and basally; microtrichose, except bare on 1st costal cell and on very small basomedian patch on alula.

Legs: Yellow; yellow pilose, except: hind femur black pilose on ventrobasal 1/4, hind tibia dorsally with long black pile on apical 2/3, hind tarsus dorsally black pilose. Coxae and trochanters yellow; yellow pilose, except hind coxa and hind trochanter partly black pilose.

Abdomen. Yellow. Tergite 1 and 2 yellow pilose. Tergite 3 black pilose. Tergite 4 black pilose, except for median vitta of yellow pilosity on posterior 3/4, and yellow pilose along posterior margin. Tergite 5 black pilose, except yellow pilose posteromedially. Second tergite slightly wider than thorax, widest at approximately 1/2. Sternites yellow; yellow pilose; sternite 1 bare.

Etymology. The name refers to ants of the genus *Crematogaster*, in which the puparia of the type specimens were found.

Diagnosis. The yellow vertex with patches of black pile is shared with *S. goettei*, *S. guianica*, *S. lanei* and *S. maculipennis*. From these species, *S. crematogastris* spec. nov. differs by the evenly yellow coloured abdomen and the yellow tinged wings. For further characters see key.



130



131



132



133



134



135




136



137



138



139



140



141

FIGURES 130–141. 130–133. *Stipomorpha crematogastri* female (holotype). 130. head frontal; 131. head lateral; 132. puparium dorsal; 133. puparium lateral. 134–137. *Stipomorpha dichromata* female (holotype). 134. habitus dorsal; 135. habitus lateral; 136. head frontal; 137. head lateral. 138–141. *Stipomorpha elcopala* male (holotype). 138. head frontal; 139. habitus lateral; 140. habitus dorsal; 141. wing.

Notes. The labels of the type specimens indicate that they were collected in dry forest. The holotype was apparently found in and subsequently reared from a nest of a *Crematogaster* species (Hymenoptera: Formicidae). For the empty puparium see figs. 132-133.

Distribution. Only known from Brazil (Mato Grosso).

Stipomorpha dichromata spec. nov.

Figs 134–137.

Studied type specimens. HOLOTYPE. BRAZIL. Female. Label 1: " Nova Teutonia / 27°11'S-52°23'W / Brazil, 300-500 m. / XI.1969 / Fritz Plaumann". Coll. CNC.

Description (based on holotype)

Adult female. Body size: 8 mm.

Head. Face occupying about 1/3 of head width in frontal view; shining dark brown; white pilose. Gena hardly developed, eyes almost directly bordering oral margin; brown. Lateral oral margins slightly produced; not reaching below eye margin in lateral view. Frons blackish brown; golden yellow pilose. Vertex shining blackish brown; golden yellow pilose; ocellar triangle equilateral. Occiput black; golden yellow pilose dorsally, yellowish white pilose ventrally. Eye bare. Antennal fossa about as wide as high. Antenna brown. Antennal ratio approximately as 2:1:4; basoflagellomere parallel-sided with rounded apex, with sensory pit at 2/3 from base. Arista slender, about 3/4 of length of basoflagellomere.

Thorax. Scutum black, with metallic hues along margins; appressed golden yellow pilose. Postpronotum, and postalar callus blackish, yellow pilose. Scutellum apically shallowly sulcate; blackish brown with metallic hues; appressed golden yellow pilose. Anepisternum weakly convex, without distinction between anterior and posterior part; dark brown; golden yellow pilose anteriorly and posteriorly. Anepimeron shining brown; entirely long yellow pilose dorsally. Katatergum and anatergum brown; long and short microtrichose, respectively. Katepimeron brown; bare. Katepisternum brown; yellow pilose dorsally, bare ventrally. Calypter and halter yellow.

Wing: hyaline, with faint brownish tinge, especially anteriorly; microtrichose except bare on 1st costal cell, basal 1/5 of cell r1, basal 3/4 of cell br, posterobasal 1/2 of cell bm, anterobasal 1/3 of cell cup.

Legs: Yellowish brown, femora darker basally; all femora and entire hind leg black pilose, front and middle tibiae and tarsi yellow pilose. Coxae and trochanters dark brown; white pilose.

Abdomen. Tergites 1 and 2 shining yellowish brown, with posterior margin of tergite 2 narrowly pale yellow; yellow pilose. Tergites 3-5 dark brown; shining with metallic hues, except for large dull part medially on anterior 3/2 of tergite 3 and small, oval dull part medially on anterior 1/4 of tergite 4; golden yellow pilose, except black pilose on dull parts. Sternites yellowish brown, yellow pilose, except sternite 1 bare.

Etymology. The name *dichromata* is Greek for two-coloured.

Diagnosis. The contrasting colour pattern of the abdomen is unique among all known *Stipomorpha* species: tergites 1 and 2 yellowish brown, other tergites dark brown.

Distribution. Only known from one specimen from Brazil, Nova Teutonia.

Stipomorpha elcopala spec. nov.

Figs 138–141, 231.

Studied type specimens. HOLOTYPE. COSTA RICA. Male. Label 1: "COSTA RICA. Cartago, Cordillera / central, Pejivalle, Humo, El Copal / 1026 m asl. Malaise trap. / 09°46'55.9"N 83°45'09.2"W / 9-10.IX.2010. Leg. J.T. Smit.". Coll. INBIO.

PARATYPE. HONDURAS. Male. Label 1: "Honduras: Yoro, Palo de / Comba, 15°11'N, 87°39'W / 29.IX.1995 / leg. R. Cave". Coll. MZLU.

Description (based on holotype)

Adult male. Body size: 5 mm.

Head. Face occupying 1/4 of head width in frontal view; black with narrow yellow lateral margins; entirely white pilose. Gena black. Occiput black; black pilose dorsally, white pilose ventrally. Oral cavity with slightly

produced lateral margins. Frons black; short black pilose, except for bare triangular part posterior to lunula. Vertex black; black pilose. Eye bare. Antennal fossa about as wide as high. Antenna black; scape and pedicel black pilose; antennal ratio 3:1:6; basoflagellomere parallel-sided with narrowly rounded apex, with small sensory pit at 3/4 from base; arista slender, about 3/4 of length of basoflagellomere, appearing bare under low magnification.

Thorax. Black. Postpronotum, scutum and postalar callus short black pilose, except for two small patches of white pile along transverse suture and two small white pilose patches anterior to scutellum. Scutellum subrectangular, without calcars; white pilose. Anepisternum a little convex, without sulcus, black pilose anterodorsally and along posterior margin. Anepimeron black pilose dorsally, white pilose ventrally. Katepisternum white pilose dorsally. Katepimeron bare. Calypter brownish yellow, halter blackish.

Wing: hyaline, with very faint dark cloud between apex of costal cell and vena spuria, and with yellowish veins bordering and posterior to pterostigma; microtrichose except bare basally on cell r1 along vein RS, on posterior 1/2 of cell br, basal 5/6 of cell bm, anterior 1/2 of cell cup and mediobasal 1/5 of alula.

Legs: brownish black, except fifth tarsomeres of all legs yellow. Legs black pilose, except anterior four tibiae posteriorly pale pilose and hind-tibia pale pilose on basal 2/3; pile on hind-tibia about as long as width of tibia. Coxae and trochanters black pilose.

Abdomen. Reddish, except lateral margins of tergites 1 and 2 blackish brown. Tergite 2 wider than thorax, widest point at around half the length; tergites 3 and 4 strongly narrowing. Antetergite very large. Tergite 1 laterally mixed black and yellow pilose, bare medially. Tergite 2 black pilose laterally, yellow pilose anterolaterally and sublaterally; almost bare medially. Tergites 3 and 4 fused, without a visible suture; sparsely black pilose medially. Sternite 1 bare, separated from sternite 2 by a membrane of about the width of sternite 1. Sternite 2 bare, laterally more than twice as wide as medially, separated from sternite 3 by a membrane of twice the median width of sternite 2. Genitalia as in fig. 231.

Female. Unknown.

Etymology. The name of this species refers to its type locality: El Copal, a nature reserve in the Central Valley of Costa Rica. The name is to be treated as an adjective.

Diagnosis. Although the black thorax and legs in combination with the reddish abdomen remind of *Stipomorpha apicula* (Curran), this species is morphologically most similar to *Stipomorpha lacteipennis* (Shannon). These two species share similar length ratios of the antennal segments, a pilose posterior margin of the anepisternum, a pilose dorsal part of the katepimeron, a partially bare alula, a notably triangular abdomen, and similar morphology of the male genitalia. However, the reddish abdomen readily separates *S. elcopala* from *S. lacteipennis*.

Notes. The paratype differs from the holotype in its larger size (6.5 mm) and the entirely yellow tergites 1 and 2 (lateral margins not blackish brown as in holotype).

Distribution. Known from Costa Rica and Honduras.

***Stipomorpha fallax* spec. nov.**

Figs 142–144, 232.

Studied type specimens. HOLOTYPE. COSTA RICA. Male, Atenas, 18.IV–15.V.1995, leg. M.J. Sommeijer. Coll. ZMAN.

PARATYPE. PANAMA. Male. Label 1: "Museum Leiden, Canal Zone, 8 km NW Gamboa, Pipeline Rd. 9°10'N; 79°45'W."; label 2: "From *Luehea seemannii* (Tiliaceae). 30.III.1976. Y. Lubin & G. Montgomery". Coll. RMNH.

Description (based on holotype)

Adult male. Body size: 7.5 mm.

Head. Face occupying about 1/4 of the head width in frontal view; shining pale yellow; with whitish pilosity, most dense sublaterally and ventrally, very sparse medially; with narrow strip of white pubescence along eye margins. Gena hardly developed, eyes almost directly bordering oral margin; yellow. Lateral oral margins not produced; not reaching below eye margin in lateral view. Frons about as long as width of lunula; black; yellow pilose laterally. Vertex shining black; golden yellow pilose. Occiput black; golden yellow pilose dorsally, yellowish white pilose ventrally. Eye very sparsely and short pilose, with pili about as long as ommati diameter, appearing

bare under low magnification. Antennal fossa about as wide as high. Antenna brown. Antennal ratio approximately as 2:1:3; basoflagellomere parallel-sided with rounded apex. Arista slender, about 2/3 of length of basoflagellomere.

Thorax. Scutum black dorsally, with margins widely yellow; yellow pilose, except for four patches of black pile: two anteriorly and two posteriorly of transverse suture. Postpronotum, postalar callus and scutellum yellow and yellow pilose. Scutellum semicircular; without calcaria. Anepisternum weakly convex, without distinction between anterior and posterior part; brownish anteriorly and yellow pilose, whitish yellow posteriorly and bare. Anepimeron dark brown; yellow pilose dorsally, bare ventrally. Katatergum and anatergum brown; long and short microtrichose, respectively. Katepimeron yellow; bare. Katepisternum brown; bare. Calypter and halter yellow.

Wing: hyaline, without colouration, microtrichose except bare on 1st costal cell, basal 1/2 of cell br, posterobasal 1/5 of cell bm, basal 1/4 of cell cup, basomedian 1/2 of alula.

Legs: Yellow, except hind tibia on apical 1/2 and basal four tarsomeres of hind tarsus blackish brown.; yellow pilose, except tarsi dorsally mixed black and yellow pilose and hind tibia long black pilose on apical 1/2. Coxae and trochanters yellow and yellow pilose.

Abdomen. Yellow and short yellow pilose. Second tergite slightly wider than thorax, widest at 1/2; third and fourth tergites much narrower. Sternites yellow; sparsely yellow pilose; sternite 1 bare. Genitalia as in fig. 232.

Etymology. The name *fallax* (Latin for deceitful, false) was chosen in analogy of the names *fraudator*, *mendax* and *spuria*, which have approximately the same meaning, in order to stress the similarity of these species.

Diagnosis. Orange-brown species with black vertex, very similar in size and colouration to *S. fraudator*, *S. mendax* and *S. spuria*. Diagnostic characters: anepisternum with posterodorsal margin bare, alula partly bare, scutellum yellowish, vertex black, scape twice as long as pedicel.

Notes. In the paratype from Panama the hind legs are entirely yellow, only slightly darkened on apical 1/2 of tibia and basal tarsomeres.

In the holotype the label states "From *Luehea seemannii* (Tiliaceae)", suggesting flower visiting.

Distribution. Known from Costa Rica and Panama.

***Stipomorpha fraudator* (Shannon, 1927)**

Figs 145–148, 233.

Ubristes fraudator Shannon, 1927: 20.

Studied type specimens. HOLOTYPE. BRAZIL. Male. Label 1 (small, round, red-bordered): "Holotype"; label 2: "Amazon 66 53"; label 3: "*Microdon: Ubristes fraudator* Snn.". Coll. BMNH.

Redescription (based on holotype)

Adult male. Body size: 9 mm.

Head. Face occupying about 1/4 of head width in frontal view; shining yellow with whitish pilosity, most dense sublaterally and ventrally, very sparse medially; with narrow strip of white pubescence along eye margins. Gena hardly developed, eyes almost bordering oral margin, yellow anteriorly, black posteriorly. Oral cavity with lateral margins a little produced. Frons slightly longer than width of lunula; yellow; yellow pilose laterally. Vertex a little swollen; shining black; yellow pilose. Occiput black; yellow pilose dorsally, white pilose ventrally. Eye very sparsely and short pilose, with pili about as long as ommati diameter, appearing bare under low magnification. Antennal fossa about as wide as high. Antenna brown. Antennal ratio 5:1:4; basoflagellomere parallel-sided with rounded apex, with small sensory pit located at 3/5 from base. Arista slender, about 2/3 of length of basoflagellomere.

Thorax. Scutum yellow, with large dark brown marks on most of dorsal surface, separated by narrow yellow lines medially, submedially and along transverse suture; yellow pilose, except for two patches of black pile posterior to transverse suture. Postpronotum, postalar callus and scutellum yellow and yellow pilose. Scutellum without calcaria. Pleurae yellow dorsally, brownish ventrally. Anepisternum weakly convex, without distinction between anterior and posterior part; yellow pilose anterodorsally, bare posteriorly. Anepimeron yellow pilose dorsally. Katatergum and anatergum long and short yellow microtrichose, respectively. Other pleurae bare. Calypter and halter yellow.



FIGURES 142–153. 142–144. *Stipomorpha fallax* male (holotype). 142. habitus dorsal; 143. habitus lateral; 144. head frontal. 145–148. *Stipomorpha fraudator* male (holotype). 145. habitus dorsal; 146. habitus lateral; 147. head frontal; 148. head lateral. 149–153. *Stipomorpha goettei* female. 149. (lectotype), head frontal; 150. (lectotype), head lateral; 151. (Surinam, coll. RMNH), head dorsal; 152. sternites 1–3 lateral; 153. sternites 1–3 ventral.

Wing: hyaline, without colouration, microtrichose except bare on posterobasal 1/2 of cell br, basal 1/4 of cell bm and basal 1/4 of cell cup.

Legs: Yellow and yellow pilose, except: posterior tibia blackish with narrowly yellow base, white pilose on basal 3/4, black pilose on apical 1/4; first three tarsomeres of all tarsi blackish and blackish pilose dorsally, fourth tarsomeres a little paler, fifth tarsomere yellow and yellow pilose. Hind tarsus ventrally with dense, appressed yellow pile. Coxae and trochanters yellow and yellow pilose.

Abdomen. Yellow and short yellow pilose. Second tergite wider than thorax, widest at basal 1/3; third and fourth tergites much narrower. Tergite 1 with anteromedian smooth, concave area. Sternites yellow. Genitalia as in fig. 233.

Female. Unknown.

Diagnosis. Orange-brown species with black vertex, very similar in size and colouration to *S. fallax*, *S. mendax* and *S. spuria*. Diagnostic characters: anepisternum with posterodorsal margin bare, alula entirely microtrichose, wing without whitish cloud or fascia, front- and mid-legs with first 3-4 tarsomeres dark brown.

Distribution. Only known from the holotype from Brazil.

***Stipomorpha goettei* (Shannon, 1927) comb. nov.**

Figs 149–153, 234.

Microdon (Ubristes) goettei Shannon, 1927: 19.

Studied type specimens. BRAZIL. Four syntypes examined, one of which is here designated as lectotype. Old labels: "Syntype"; "Ega"*; "*Microdon: Ubristes goettei* Snn."; added label: "*Microdon goettei* Shannon, LECTOTYPE, design. M. Reemer". For label information of paralectotypes see table 2. For further notes see account of *Stipomorpha guianica*.

Additionally studied specimens. SURINAM: 1 male, Akintosoela, Mapane area, SME Q.22, 28.VII.1995, leg. B. De Dijn & A. Gangadin, coll. RMNH; 1 female, SE of Zanderij, road to Kraka, 16.III.2006, leg. M. Reemer, coll. RMNH; FRENCH GUYANA: 1 female, Roura, Kaw Road, PK 37, Relais Patawa, N 04°32'42"–W 52°09'09", malaise trap, August 2008, leg. J.A. Cerda, coll. RMNH.

Body size: 8–10.5 mm.

Redescription (based on lectotype)

Adult female. Body size: 10 mm.

TABLE 2. Syntypes of *Microdon goettei* Shannon, 1927 and their new type status. Two of these specimens are identified as belonging to *S. guianica* (Curran). All specimens are females and are kept in the collection of the BMNH.

Labels	New type status	New identity
Old labels: "Syntype"; "Ega"*; " <i>Microdon: Ubristes goettei</i> Snn." Added label: " <i>Microdon goettei</i> Shannon, LECTOTYPE, design. M. Reemer"	lectotype <i>U. goettei</i> Shannon	<i>Stipomorpha goettei</i> (Shannon)
Old labels: "Syntype"; "Amazon. 6653" Added label: " <i>Microdon goettei</i> Shannon, id. M. Reemer 2007"	paralectotype <i>U. goettei</i> Shannon	<i>Stipomorpha goettei</i> (Shannon)
Old label: "Syntype"; "Amazon. 6653"; " <i>Microdon: Ubristes goettei</i> Snn." Added label: " <i>Microdon guianicus</i> Curran, id. M. Reemer 2007"	paralectotype <i>U. goettei</i> Shannon	<i>Stipomorpha guianicus</i> (Curran)
Old labels: Syntype; Para Added label: " <i>Microdon guianicus</i> Curran, id. M. Reemer 2007"	paralectotype <i>U. goettei</i> Shannon	<i>Stipomorpha guianicus</i> (Curran)

*: Ega = Tefé.

Head. Face occupying about 1/3 of head width in frontal view; in dorsal half with tubercles along eye margin; yellow, a little darker on dorsal 1/3; yellow pilose, short dorsally and medially, long ventrally and ventrolaterally; with narrow strip of white pubescence along eye margin. Gena yellow. Oral cavity with lateral margins a little produced and notched anteriorly. Frons about as long as width of lunula; blackish posterior to lunula, otherwise yellow; yellow pilose, except for posterolateral patches of black pile. Vertex swollen; yellow; with fasciae of black pile from ocellar triangle to black pile patches on frons; with black pile on ocellar triangle and along posterior

margin; with yellow pile anteriorly and laterally. Occiput black; yellow pilose dorsally, more whitish pilose ventrally. Eye very sparsely and short pilose, with pili about as long as ommati diameter, appearing bare under low magnification. Antennal fossa about as wide as high. Antenna with scape yellow, pedicel and basoflagellomere brown. Antennal ratio 3:1:5; basoflagellomere parallel-sided with rounded apex. Arista slender, about 2/3 of length of basoflagellomere.

Thorax. Scutum black, with widely yellow margins; yellow pilose, except for four patches of black pile, two of which anteriorly and two posteriorly of transverse suture. Postpronotum, postalar callus and scutellum yellow and yellow pilose. Scutellum without calcaria. Pleurae yellow dorsally, brownish ventrally. Anepisternum weakly convex, without distinction between anterior and posterior part; yellow pilose anterodorsally, bare posteriorly. Anepimeron yellow pilose dorsally. Katatergum and anatergum yellow pilose and microtrichose, respectively. Other pleurae bare. Calypter and halter yellow.

Wing: hyaline, with a wide white fascia on apical half (best visible against dark background); microtrichose except bare on most of cell br, basal 1/5 of cell bm and basal 1/5 of cell cup.

Legs: Front- and mid-legs yellow; yellow pilose. Hind femur yellowish brown, vaguely blackish on median 1/3; yellow pilose. Hind tibia strongly widened; brownish yellow, vaguely blackish around cicatrice on apical half; with long, dense pilosity dorsally: white on basal 1/3, black on apical 2/3; short yellow pilose apicolaterally. Hind tarsus: first tarsomere yellow and yellow pilose dorsally, except for black pile apically; other tarsomeres dark brown and black pilose dorsally; all tarsomeres densely, short, golden pilose ventrally. Coxae and trochanters yellow and yellow pilose.

Abdomen. Yellowish brown with large, vaguely demarcated, dark brown to black lateral markings on tergites 2-5, leaving only narrow median yellow vittae and widely yellow apical margins. Tergite 1 with anteromedian smooth, concave area. Second tergite about as wide as thorax, widest at basal 1/3; third and fourth tergites narrower, more or less parallel-sided. Entirely short, yellow pilose. Sternites brownish. Genitalia as in fig. 234.

Diagnosis. The yellow vertex with patches of black pile is shared with *S. crematogastris*, *S. guianica*, *S. lanei* and *S. maculipennis*. From these species, *S. goettei* differs by the absence of pile on the posterior part of the anepisternum in combination with the whitish cloud on the wing (view against dark background). Differences between *S. goettei* and *S. guianica* are listed in table 3.

TABLE 3. Differences between *Stipomorpha goettei* (Shannon) and *S. guianica* (Curran). Unless stated otherwise, characters apply to both sexes.

<i>Stipomorpha goettei</i> (Shannon)	<i>Stipomorpha guianica</i> (Curran)
face in dorsal half with tubercles along eye margin	face smooth, without tubercles along eye margin
face in profile slightly concave at middle and clearly convex on lower half	face in profile straight or slightly convex
basoflagellomere almost 1.5 times longer than scape (antennal ratio 3:1:5)	basoflagellomere only slightly longer than scape (antennal ratio 4:1:(4-5))
posterior part of anepisternum without pile	posterior part of anepisternum with pile along posterodorsal margin
female: tergites 3 and 4 with pale median vitta	female: tergites 3 and 4 without pale median vitta

Notes. Shannon (1927) based his description of *Microdon goettei* on five females. Only four of these syntypes could be found in the BMNH-collection. This series of syntypes was found to consist of two closely similar species, each represented by two specimens. Two syntypes were found to agree with *Microdon guianicus* Curran. The other two are here considered as the 'real' *Microdon goettei* Shannon. Support for this view is provided by the original description, which states that in the antennae the 'third [joint is] a little longer than combined length of first and second' (Shannon 1927). To ensure the stability of the taxon, a lectotype is designated for *Microdon goettei* Shannon out of the syntype series (see table 2).

Distribution. Known from Brazil, French Guyana and Surinam. All records of "*Ubristes goettei*" from Surinam by Van Doesburg (1966) belong to *Stipomorpha guianica*, but specimens belonging to the 'real' *S. goettei* were collected in recent years (see additionally studied specimens).



FIGURES 154–162. 154–157. *Stipomorpha guianica* (Surinam, coll. RMNH). 154. male, habitus dorsal; 155. male, habitus lateral; 156. male, head frontal; 157. female, habitus dorsal. 158–161. *Stipomorpha inarmata* male (holotype). 158. habitus dorsal; 159. habitus lateral; 160. head frontal; 161. wing. 162. *Stipomorpha lacteipennis* male (holotype), habitus dorsal.

Stipomorpha guianica (Curran, 1925)

Figs 154–157, 235.

Microdon guianicus Curran, 1925: 340.

Studied type specimens. HOLOTYPE. GUYANA. Female. Bartica. Pictures of types studied from type database of the Museum of Comparative Zoology, accessible on the internet. Coll. MCZ.

Additionally studied specimens. ECUADOR: 1 female, Limon, 900 m., II.1948, Z. Muller, coll. AMNH. FRENCH GUYANA: 1 male, Montagnes Tortue, 04°15.007'N–52°21.512'W, 12.I.2003, leg. V. Soon, coll. RMNH. GUYANA: 1 male, Kurupkari: 4°4'N–8°40'W, malaise trap, IX–XI.1992, coll. BMNH. PERU: 1 female, Madre de Dios, Rio Tambopata, Sachavacayoc Centre, 12°51'S–69°22'W, 16–26.X.2008 (malaise trap), leg. & coll. J.T. Smit. SURINAM: 1 female, Perica, 20.VIII–3.IX.1997 malaise trap, leg. B. De Dijn, coll. RMNH; 1 female, Perica, 10–24.XII.1997, malaise trap, leg. B. De Dijn, coll. RMNH; 1 female, Paramaribo Leiding, 28.I–6.II.2006, malaise trap, leg. M. Reemer, coll. RMNH; 1 female, Peperpot, 2–9.II.2006, malaise trap, leg. M. Reemer, coll. RMNH; 1 male, Paramaribo Cultuurtuin, 7.III.2006, leg. M. Reemer, coll. RMNH; 1 female, Peperpot, 21–28.III.2006, malaise trap, leg. M. Reemer, coll. RMNH; 1 male, Peperpot, 28.III.2006, leg. M. Reemer, coll. RMNH; 1 female, Peperpot, 29.III–6.IV.2006, malaise trap, leg. M. Reemer, coll. RMNH; 2 females, Peperpot, 6–14.IV.2006, malaise trap, leg. M. Reemer, coll. RMNH; 2 females, 14–20.IV.2006, malaise trap, leg. M. Reemer, coll. RMNH; 1 female, Peperpot, 20–27.IV.2006, malaise trap, leg. M. Reemer, coll. RMNH.

Redescription (based on additionally studied material from Surinam)

Adult male. Body size: 7.5–10 mm.

Head. Face occupying about 1/4 of head width in frontal view; without tubercles along eye margin; yellow; yellow pilose, short dorsally and medially, long ventrally and ventrolaterally; with narrow strip of white pubescence along eye margin. Gena yellow. Oral cavity with lateral margins a little produced and notched anteriorly. Frons about as long as width of lunula; yellow; black pilose, except yellow pilose laterally. Vertex swollen; yellow, except black on ocellar triangle; black pilose, with pile more dense and appressed on anterior 1/2. Occiput black; yellow pilose, except black pilose anterodorsally. Eye very sparsely and short pilose, with pili about as long as ommati diameter, appearing bare under low magnification. Antennal fossa slightly higher than wide. Antenna with scape yellowish brown, pedicel and basoflagellomere brown. Antennal ratio 4:1:(4-)5; basoflagellomere parallel-sided with rounded apex. Arista slender, about 5/6 of length of basoflagellomere.

Thorax. Scutum black, with widely yellow margins; yellow pilose, except for fasciae of black pile, one of which anterior of and the other posterior of transverse suture. Postpronotum, postalar callus and scutellum yellow and yellow pilose. Scutellum without calcars. Pleurae yellow dorsally, brownish ventrally. Anepisternum weakly convex, without distinction between anterior and posterior part; yellow pilose anterodorsally and along posterodorsal margin. Anepimeron yellow pilose dorsally. Katatergum and anatergum yellow pilose and microtrichose, respectively. Other pleurae bare. Calypter and halter yellow.

Wing: hyaline, with a wide white fascia on apical half (best visible against dark background); microtrichose except bare on basal 1/2 of cell br, basal 1/5 of cell bm and basal 1/10 of cell cup, basomedian 2/3 of alula.

Legs: Front- and mid-legs yellow, third and fourth tarsomeres slightly darkened; yellow pilose. Hind femur yellowish brown, vaguely blackish on median 1/3; yellow pilose. Hind tibia strongly widened; brownish yellow, blackish on median 1/3; with long, dense pilosity posteriorly: white on basal 1/4, black on apical 3/4; short yellow pilose ventrally. Hind tarsus: first tarsomere yellow and black pilose dorsally; tarsomeres 2–4 blackish and black pilose dorsally; tarsomere 5 yellow and yellow pilose dorsally; all tarsomeres densely, short, golden pilose ventrally. Coxae and trochanters yellow and yellow pilose.

Abdomen. Dark brown, except tergite 1, anterior and lateral margins of tergite 2, posterolateral corner of tergite 3 and lateral and posterior margins of tergite 4 yellowish. Tergite 1 with anteromedian smooth, concave area. Second tergite slightly wider than thorax, widest at basal 1/4; third and fourth tergites narrower, more or less parallel-sided. Entirely short, yellow pilose. Sternites brownish. Genitalia as in fig. 235.

Diagnosis. The yellow vertex with patches of black pile is shared with *S. crematogastris*, *S. goettei*, *S. lanei* and *S. maculipennis*. From these species, *S. guianica* differs by the presence of pile on the posterior part of the anepisternum, the partly bare alula and the absence of a dark brown anteromedian spot on the wing. For further characters see key. For differences with *S. goettei* see table 3.

Notes. All records of "*Ubristes goettei*" from Surinam by Van Doesburg (1966) belong to *S. guianica*. In Surinam, the present author observed this species visiting flowers on two occasions: a male on 17.III.2006 and a male on 28.III.2006.

Distribution. Known from Ecuador, Guyana, Surinam, French Guyana and Peru.

Stipomorpha inarmata (Curran, 1925)

Figs 158–161, 236.

Microdon inarmatus Curran, 1925: 5.

Studied type specimens. HOLOTYPE. GUYANA. Male. Label 1: "Collection C.W. Johnson"; label 2: "Bartica, BG, IV.I 1901"; label 3 (red): "Type *Microdon inarmatus* Curran"; label 4 (red): "Type 7656"; label 5: "Jan.-July 2003, MCZ Image Database". Coll. MCZ.

Additionally studied specimens. BRAZIL: 1 male & 1 female, Boca do Cuminá-Miri, Oriximiná, PA [Pará], 19-26.I.1968, Exp. Perm. Amaz, coll. USNM. FRENCH GUYANA: 1 female, Kaw Mountains, 04°32.893'N-52°10.245'W, 8.XII.2002, leg. V. Soon, coll. RMNH; 1 female, Roura, Kaw Road, PK 37, Relais Patawa, N 04°32'42"- W 52°09'09", malaise trap, XII.2008, leg. J.A. Cerda, coll. RMNH. GUYANA: 1 male, Essequibo R., Moraballi Creek, 29.IX.1929, leg. Oxf. Univ. Expedn., coll. CNC.

Redescription

Adult male. Body size: 8 mm.

Head. Face occupying 1/4 of head width in frontal view; shining yellow with blackish brown median stripe from oral margin to antennal fossa; face with white pilosity, a little longer around oral margin, except bare on median stripe. Gena brown. Occiput black; black pilose dorsally, getting white laterally and ventrally. Oral cavity with produced lateral margins and notched anterior margin. Frons and lunula black and short black pilose, except for bare triangular part posterior to lunula. Vertex black; black pilose. Eye very sparsely and short pilose, with pili about as long as ommati diameter, appearing bare under low magnification. Antennal fossa about as wide as high. Antenna brown; scape and pedicel dark pilose; antennal ratio 4:1:3.5; basoflagellomere parallel-sided with narrowly rounded apex, with sensory pit located at 3/4 from base, within a vague groove that ranges from just before 1/2 to just after the pit; arista slender, about 2/3 of length of basoflagellomere, very shortly pilose, appearing bare under low magnification.

Thorax. Scutum black, postpronotum and postalar callus pale brown, pleurae dark brown, scutellum dark brown. Postpronotum, scutum, postalar callus and scutellum short black pilose, except scutum with lateral fasciae of white pile along transverse suture and lateral prescutellar patches of white pile. Scutellum semicircular, without calcars. Anepisternum more or less flat, yellow pilose anterodorsally, bare posteriorly. Anepimeron pilose posterodorsally. Katepisternum and katepimeron bare. Calypter grey, halter yellowish.

Wing: hyaline; microtrichose except bare on basal 3/4 of cell br, basal 1/2 of cell bm, anterobasal 1/4 of cell cup, basomedially on alula.


Legs: brownish black, except anterior four tarsi yellow with first tarsomeres darker, and hind tarsi with last three tarsomeres yellow. Femora black pilose; anterior four tibiae white pilose; hind tibia white pilose on basal 2/3, black pilose on apical 1/3; tarsi black pilose dorsally, yellow pilose ventrally. Pile on hind-atibia a little longer than half the width of the tibia. First tarsomere of hind tarsus as long as 1/3 of length of hind-tibia, a little wider than apex of tibia, twice as long as wide (dorsal view). Coxae and trochanters pale pilose.

Abdomen. Blackish brown, except tergites 1 & 2 pale brown. Second segment wider than thorax, widest point at half the length; third and fourth tergites strongly narrowing. Tergites pale pilose. Sternite 1 bare, sternite 2 probably bare, sternite 3 and 4 pilose. Genitalia as in fig. 236.

Female. Unknown.

Diagnosis. From other *Stipomorpha*-species with a black thorax, *S. inarmata* can be recognized by the following characters: face largely yellow with narrow median brown stripe, basoflagellomere slightly shorter than scape, alula partly bare, anepisternum only pilose anterodorsally, katepisternum bare, structure of male genitalia. The tarsi may be darker than in holotype.

Distribution. Known from Guyana, French Guyana and northern Brazil.



FIGURES 163–172. 163–164. *Stipomorpha lacteipennis* male (holotype). 163. habitus lateral; 164. head frontal. 165. *Stipomorpha lanei* male (Surinam, coll. RMNH), habitus dorsal. 166–167. *Stipomorpha lanei* female (French Guyana, coll. RMNH). 166. head frontal; 167. head dorsal. 168. *Stipomorpha lanei* female (holotype), habitus dorsal. Photo: American Museum of Natural History. 169–170. *Stipomorpha litoralis* male (holotype). 169. habitus dorsal; 170. habitus lateral. 171–172. *Stipomorpha mackiei* male (holotype). 171. habitus dorsal; 172. habitus lateral.

***Stipomorpha lacteipennis* (Shannon, 1927)**

Figs 162–164, 237.

Microdon lacteipennis Shannon, 1927: 18.

Microdon triangularis Curran, 1940: 6. Syn. nov.

Studied type specimens. LECTOTYPE *Microdon lacteipennis* Shannon. BRAZIL. Male. Label 1 (blue label): "Syntype"; label 2: "Amazon. 66.53"; label 3: "*Microdon Ubristes lacteipennis* Snn.". Coll. BMNH.

HOLOTYPE *Microdon triangularis* Curran. BRAZIL. Male. Label 1 (red label): "Microdon triangularis Curran Holotype"; label 2: "Abril 1937"; label 3: "Servico Febre Amarela, M.E.S., Bras."; label 4: Douradas, Mato Grosso, Brasil". Coll. AMNH.

PARATYPE *Microdon triangularis* Curran. BRAZIL. Male. Label 1: "Dourados, Mato Gross, Brasil", label 2: "Abril 1937", label 3: "Servico Febre, Amazon, M.E.S., Bras.", label 4: "R.C. Shannon collection", label 5: "From type series", label 6: "*Microdon triangularis* Cur., det. F.M. Hull". Coll. USNM.

Additionally studied specimens. BOLIVIA: 1 male, Santa Cruz distr. 4 km N Bermejo, Refugio Los Volcanes, 1000 m, 18°06'S, 63°36'W, 25–30.X.2007, leg. A.R. Cline, coll. RMNH; BRAZIL: 1 male, Mato Grosso, Dourados, IV.1937, leg. Servico Febre Amarela, coll. CNC; PERU: 1 male, Madre de Dios, Tambopata, Sachavacayoc centre, 12°51'20"S–69°22'20"W, 25.VI–4.VIII.2010, leg. J.T. Smit, coll. J.T. Smit; 1 male, same data as previous except date 23.III–28.IV.2011; SURINAM: 1 male, Blakawatra, 13.VI.1963, leg. J. v.d. Vecht, coll. RMNH; 1 female, Peperpot, 20–27.IV.2006, leg. M. Reemer, coll. RMNH. VENEZUELA: 1 male, T.F. Amaz., Cerro de la Neblina, Basecamp, 140 m., 0°50'N, 66°10'W, 10–20.II.1985, malaise trap in rainforest, leg. P.J. & P.M. Spangler, R.A. Faitoute, W.E. Steiner colrs, coll. USNM.

Redescription (based on lectotype)

Adult male. Body size: 5.5–7 mm.

Head. Face occupying 1/5 of head width in frontal view; black with narrow yellow lateral margins; entirely white pilose. Gena black. Occiput black; black pilose dorsally, white pilose ventrally. Oral cavity with slightly produced lateral margins. Frons black; short black pilose, except for bare triangular part posterior to lunula. Vertex black; black pilose. Eye appearing bare under low magnification. Antennal fossa about as wide as high. Antenna black, with basoflagellomere a little brownish; scape and pedicel black pilose; antennal ratio 3:1:6; basoflagellomere parallel-sided with narrowly rounded apex, with small sensory pit at 2/3 from base; arista slender, about 3/4 of length of basoflagellomere, appearing bare under low magnification.

Thorax. Black, postalar callus a little brownish. Postpronotum, scutum, postalar callus and scutellum short black pilose, except for two small patches of white pile along transverse suture and two small white pilose patches anterior to scutellum. Scutellum subrectangular, without calcars. Anepisternum a little convex, without sulcus, black pilose anterodorsally and along posterior margin. Anepimeron black pilose dorsally, white pilose ventrally. Katepisternum white pilose dorsally. Katepimeron bare. Calypter brownish yellow, halter blackish.

Wing: hyaline, with faint dark cloud between apex of costal cell and vena spuria, and with faint yellowish cloud on and posterior to pterostigma; microtrichose except bare on posterobasal 1/2 of cell br, posterobasal 1/2 of cell bm, anterobasal 1/2 of cell cup and mediobasal 1/5 of alula.

Legs: brownish black, except fifth tarsomeres of all legs yellow. Legs black pilose, except anterior four tibiae pale pilose and hind-tibia pale pilose on basal 1/2; pile on hind-tibia about as long as width of tibia. First tarsomere of hind-tarsus as long as 1/3 of length of hind-tibia, clearly wider than apex of tibia, twice as long as wide. Coxae and trochanters black pilose.

Abdomen. Black. Tergite 2 wider than thorax, widest point at around half the length; tergites 3 and 4 strongly narrowing. Antetergite very large. Tergite 1 laterally mixed black and white pilose, bare medially. Tergite 2 black pilose laterally, white pilose anterolaterally and sublaterally; almost bare medially. Tergites 2 and 3 separated by a yellowish membrane of almost the median length of tergite 2. Tergites 3 and 4 fused, without a visible suture; black pilose dorsally and posteriorly, bare laterally. Sternite 1 bare, separated from sternite 2 by a membrane of about the width of sternite 1. Sternite 2 bare, laterally more than twice as wide as medially, separated from sternite 3 by a membrane of twice the median width of sternite 2. Genitalia as in fig. 237.

Female. As Van Doesburg (1927) already noted, the female is quite similar to the male, except for usual sexual dimorphism and wing pattern more pronounced.

Diagnosis. *Stipomorpha lacteipennis* shares its pilose posterior anepisternum and pilose dorsal part of the

katepisternum only with *S. litoralis*. From this species it differs by the presence of a whitish fascia in the wing and by the male genitalia.

Notes. Shannon (1927) only described the male, based on two male syntypes. In the BMNH collection there is only one syntype left, which is hereby designated as lectotype in order to stabilize nomenclature. The holotype and a paratype of *Microdon triangularis* Curran have been examined and were found to be conspecific with *S. lacteipennis* Shannon.

Distribution. Known from Bolivia, Brazil, Peru, Surinam and Venezuela.

***Stipomorpha lanei* (Curran, 1936)**

Figs 165–168, 238.

Microdon lanei Curran, 1936: 5.

Studied type specimens. HOLOTYPE. BRAZIL. Female. Label 1: "Juquia - S.P., J. Lane, XI, 1929"; label 2 (red): "*Microdon lanei* Curran Holotype". Coll. AMNH.

Additionally studied specimens. FRENCH GUYANA: 1 female, Kaw mountains, 04°33.562'N–52°12.425'W, 21.X.2002, leg. V. Soon, coll. RMNH; 1 male, Roura, Kaw road, PK 37, Relais Patawa, N 04°32'42"–W 52°09'09", malaise trap, XII.2009, leg. J.A. Cerda, coll. RMNH; SURINAM: 1 male, Paramaribo, 9.XII.1957, leg. P.H. van Doesburg Jr., coll. RMNH; 1 female, Kwatta, 8.II.1964, leg. D.C. Geijskes, coll. RMNH; 1 male, Paramaribo, Leiding, 05°17'03"W, 28.I–6.II.2006, leg. M. Reemer, coll. RMNH. VENEZUELA: 1 male, T.F. Amaz., Cerro de la Neblina, 140 m., 0°50'N–66°10'W, 21–28.II.1985 (malaise trap in rain forest), leg. P.J. & P.M. Spangler, R.A. Faitoute & W.E. Steiner, coll. USNM.

Redescription (based on holotype)

Adult female. Body size: 8 mm.

Head. Face occupying about 1/4 of head width in frontal view; shining yellow with yellow pilosity, except on a bare median line occupying 1/2 of the width of the face. Gena yellow. Oral cavity with lateral margins not produced and not notched anteriorly. Frons yellow; short black pilose, laterally and posteriorly. Vertex yellow, except brown on ocellar triangle and posterior to it; short black pilose. Occiput yellow; black pilose dorsally, yellow pilose ventrally and over entire posterior surface. Eye appearing bare under low magnification. Antennal fossa about as wide as high. Scape and pedicel dark brown, basoflagellomere reddish brown. Antennal ratio 4:1:7. Basoflagellomere parallel-sided with narrowly rounded apex; with oval sensory pit, occupying 1/5 of height of basoflagellomere, located at 2/3 from base, within a vague 'sensory groove' that ranges from ventrad of the base of the arista almost to the apex. Arista slender, about 2/3 of length of basoflagellomere.

Thorax. Yellowish brown, except for two wide, vaguely demarkated dark brown lateral vittae on the scutum, narrowly divided in two along the transverse suture; medially the scutum is yellow anteriorly and dark brown posteriorly. Scutum short black pilose, except for two submedian vittae of short yellow pile and narrow lateral fasciae of yellow pile along the transverse suture. Postpronotum yellow pilose. Postalar callus and scutellum black pilose. Scutellum semicircular, without calcars. Anterior and posterior part of anepisternum not divided by a sulcus; anterior part yellow pilose, posterior part bare. Anepimeron yellow pilose on dorsal half. Katatergum and anatergum pilose and microtrichose, respectively. Other pleurae bare. Calypter and halter yellow.

Wing: hyaline, with veins yellowish anteriorly and apically. Microtrichose except bare along vein RS between veins R1 and R2+3, on basal 2/3 of cell br, posterobasal 1/4 of cell bm, anterobasal 1/6 of cell cup.

Legs: Entirely yellow and yellow pilose, including coxae and trochanters.

Abdomen. Yellowish brown. Tergite 2 wider than thorax, with widest point at half the length; tergites 3 and 4 narrower, with tergite 4 strongly narrowing posteriorly. Tergites 3, 4 and 5 fused, with sutures vaguely visible. Tergites entirely yellow pilose; shining, tergite 3 dull on anterior 2/5, because of a fascia of microtrichia, which occupies most of the tergite's width; tergite 4 with two anterolateral dull oval markings of microtrichia of 1/3 of the length of the tergite. Sternites yellow. Sternite 1 bare. Pilosity of other sternites hard to assess in type specimen.

Male (based on specimens from Surinam): More or less as female, except hind femur, hind tibia and abdomen black pilose. Genitalia as in fig. 238.

Diagnosis. Instantly recognizable by the unproduced yellow vertex (compare e.g. *S. guianica*, in which the

vertex is strongly produced). The male is unique among *Stipomorpha* species in the fact that the abdomen is somewhat constricted, with its smallest width at the transition between tergites 3 and 4. In the female this constriction is also present to some extent, but less pronounced.

Distribution. Known from Brazil, Surinam and Venezuela.

Stipomorpha litoralis (Papavero, 1964)

Figs 169, 170, 239.

Ubristes litoralis Papavero, 1964: 21. Type locality: Brazil, São Paulo, Caraguatatuba.

Studied type specimens. HOLOTYPE (in bad condition). BRAZIL. Male. Label 1: "Caraguatatuba - SP, (Res. Flor. - 40 m.), Brasil, 2.IV.1962, K. Lemko col."; label 2 (red): "Holotipo"; label 3: "28.646"; label 4: "*Ubristes litoralis*, sp. n., N. Papavero det. 1962". Coll. MZUSP.

Redescription (based on holotype)

Adult male. Body size: 7 mm.

Head. (head of holotype in bad condition, not all characters can be assessed) face black on median 1/2 to 3/4, with yellow laterally; entirely white pilose. Gena black. Occiput black; black pilose dorsally, white pilose ventrally. Oral cavity with produced lateral margins and notched anterior margin. Frons and vertex black and black pilose. Eye bare. Antennal fossa about as wide as high. Antenna (basoflagellomere missing in holotype) black; scape about twice as long as wide and twice as long as pedicel.

Thorax. Black, pleurae brown. Postpronotum, scutum, postalar callus and scutellum black pilose. Scutellum without calcar. Anepisternum a little convex, without sulcus, black pilose anterodorsally and along posterior margin. Anepimeron black pilose dorsally, white pilose ventrally. Katepisternum white pilose dorsally. Katepimeron bare. Calypter and halter blackish brown.

Wing: hyaline, except a little darker anterobasally; microtrichose except bare on basal 1/5 of cell br, posterobasal 1/4 of cell bm, basal 1/5 of cell cup.

Legs: brownish black, except fifth tarsomeres of middle leg yellow (other tarsi missing in holotype). Legs black pilose, except trochanter and tibia of foreleg and dorsobasal 1/2 of hind tibia pale pilose.

Abdomen. Shining blackish brown. Second segment wider than thorax, widest point at around half the length; third and fourth tergites strongly narrowing. Tergite 1 mostly black pilose, with patch of white pile at posterior 1/2 of lateral margin. Other tergites black pilose. Sternites bare. Genitalia as in fig. 239.

Diagnosis. *Stipomorpha litoralis* shares its pilose posterior anepisternum and pilose dorsal part of the katepisternum only with *S. lacteipennis*. From this species it differs by the lack of a whitish fascia in the wing and by the male genitalia.

Notes. According to Thompson *et al.* (1976) this is a synonym of *Microdon triangularis* Curran, but that taxon is here treated as a synonym of *M. lacteipennis* Shannon.

Distribution. Only known from the holotype, which is from São Paulo, Caraguatatuba, in Brazil.

Stipomorpha mackiei (Curran, 1940)

Figs 171, 240.

Microdon mackiei Curran, 1940: 5.

Studied specimens. HOLOTYPE. GUYANA. Male. Label 1: "Rockstone Br Guiana, June 2 1929 (A. Mackie)"; label 2 (red label): "Microdon mackiei Curran Holotype". Coll. AMNH.

Additionally studied specimens. PERU: 1 male, Madre de Dios, Rio Manu, B iolat biol. sta., Pakitza, 356 m., 11°56'47"S-071°17'W, 9.VII.1992, leg. T.E. Erwin, E. & F. Pfuno, coll. USNM.

SURINAM: 1 female, Blakawatra, 11.VI.1963, leg. J. v.d. Vecht, coll. RMNH; 1 male, Paramaribo, Charlesburg, 21.I.1964, leg. D.C. Geijskes, coll. RMNH; 1 female, Oost-West verbinding, 70 km E of Paramaribo, 21.X.1995, leg. B. De Dijn, coll. RMNH; 2 females, Marowijne, near Perica, malaise trap, 6–20.VIII.1997, leg. B. De Dijn, coll. RMNH; 1 female, Marowijne, near Perica, malaise trap, 21.I.–4.II.1998, leg. B. De Dijn, coll.

RMNH; 1 female, Brownsberg, malaise trap, 21.VI–5.VII.2001, leg. A. Gangadin, coll. RMNH; 1 female, Commewijne, Peperpot, 05°46'08"N–55°07'33"W, malaise trap, 24.II–7.III.2006, leg. M. Reemer, coll. RMNH; 1 female, Zanderij, 05°26'19"N–55°12'20"W, leg. M. Reemer, coll. RMNH; 1 female, Commewijne, Peperpot, 05°46'08"N–55°07'33"W, 29.III–6.IV.2006, leg. M. Reemer, coll. RMNH; 1 female, Commewijne, Peperpot, 05°46'08"N–55°07'33"W, 6–14.IV.2006, leg. M. Reemer, coll. RMNH.

Redescription (based on additional material from Surinam)

Adult male. Body size: 9 mm.

Head. Face occupying about 1/3 of head width in frontal view; yellow, with vague, narrow, brownish median vitta on dorsal 2/3; yellow pilose; with narrow strip of white pubescence along eye margin. Gena; yellow; yellow pilose. Lateral oral margins produced; anterior margin notched. Frons about as long as width of lunula; black, except yellow along lunula; yellow pilose. Vertex shining black; golden yellow pilose, except black pilose posteriad of ocelli. Occiput black; yellow pilose. Eye very sparsely and short pilose, with pili about as long as ommati diameter, appearing bare under low magnification. Antennal fossa about as wide as high. Antenna yellowish brown. Antennal ratio approximately as 4:1:6; basoflagellomere parallel-sided with rounded apex. Arista slender, about 2/3 of length of basoflagellomere.

Thorax. Scutum brownish dorsally, with margins widely yellow, transitions between yellow and brown parts vague; yellow and golden pilose, except for two patches of black pile posterior to transverse suture. Postpronotum, postalar callus and scutellum yellow and yellow pilose. Scutellum without calcars. Pleurae yellow. Anepisternum weakly convex, without distinction between anterior and posterior part; yellow; yellow pilose anterodorsally and along posterodorsal margin. Anepimeron yellow pilose dorsally, bare ventrally. Katatergum and anatergum long and short microtrichose, respectively. Katepimeron bare. Katepisternum bare. Calypter and halter yellow.

Wing: hyaline, tinged with yellow anterobasally; microtrichose, except bare on cell r1 basally along vein RS, on basal 1/2 of cell br except microtrichose along vena spuria, on basal 1/6 of cell bm, on basal 1/10 of cell cup and on basomedian 1/2 of alula.

Legs: Yellow (but see notes on variation below); yellow pilose, except brown pilose on apicodorsal 1/2 of hind tibia and dorsally on basal tarsomeres of hind leg. Coxae and trochanters yellow and yellow pilose.

Abdomen. Yellow; yellow pilose. Second tergite about as wide as thorax, widest at 1/2 of its length; third and fourth tergites much narrower. Sternites yellow; sparsely yellow pilose; sternite 1 bare. Genitalia as in fig. 240.

Diagnosis. Distinguishable from the very similar *S. tenuicauda* reliably only by the male genitalia. The character of the distribution of microtrichia on the alula, as mentioned in the key, could be verified in a few males only. Whether it works for all specimens (including females) is uncertain.

Notes. *Stipomorpha mackiei* (Curran, 1940) is described from the male (contrary to Thompson *et al.* (1976) who incorrectly state that the type is a female), which has a dark medial stripe on the face. The only known male from Surinam has a yellow face. In most females from Surinam the face is yellow, but some specimens have a faint dark medial stripe. In other characters the specimens from Surinam agree perfectly with the type of *Ubristes mackiei* (Curran). The studied male from Peru has a dark facial stripe, and also differs from the Surinam specimens in the almost entirely dark hind legs and metanotum (yellow in the Surinam specimens). However, in morphological characters, like the genitalia, the specimen is very similar and therefore is considered to belong to the same species. Apparently, colouration of hind legs and metanotum is not a reliable character.

Distribution. Known from Guyana, Peru and Surinam.



***Stipomorpha maculipennis* spec. nov.**

Figs 177–181, 241.

Studied type specimens. HOLOTYPE. ARGENTINA. Male. Label 1: "ARGENTINA: Mis., / Iguazu Nat. Park, / hosteria Hoppe. / c. 140 m. Malaise trap / 10–11.iv.1974. C.R. Vardy / B.M. 1974–204"; label 2 (red): "HOLOTYPE / *Stipomorpha* / *maculipennis* / M. Reemer". Coll. BMNH.

Description (based on holotype)

Adult female. Body size: 9 mm.



FIGURES 173–185. 173–174. *Stipomorpha mackiei* male (holotype). 173. head frontal; 174. head lateral. 175–176. *Stipomorpha*, alula. 175. *S. mackiei*; 176. *S. tenuicauda*. 177–181. *Stipomorpha maculipennis* male (holotype). 177. habitus dorsal; 178. habitus lateral; 179. head frontal; 180. head lateral; 181. wing. 182–185. *Stipomorpha mendax* male (holotype). 182. habitus dorsal; 183. habitus lateral; 184. head frontal; 185. head lateral.

Head. Face occupying slightly more than 1/3 of head width in frontal view; yellow; yellow pilose, short dorsally and medially, long ventrally and ventrolaterally. Gena yellow. Oral cavity with lateral margins produced. Frons about as long as width of lunula; blackish posterior to lunula, otherwise yellow; yellow pilose, except for posterolateral patches of black pile. Vertex swollen, in profile produced for about 1/3 of height of eye; yellow; yellow pilose, except for two large anterior patches of black pile and black pile along posterior margin. Occiput yellow, except blackish over short dorsolateral stretch; yellowish pilose. Eye bare. Antennal fossa about as wide as high. Antenna with scape yellowish brown, pedicel and basoflagellomere brown. Antennal ratio approximately as 3:1:5; basoflagellomere parallel-sided with rounded apex, with sensory pit at about 2/5 from base. Arista slender, about 3/4 of length of basoflagellomere.

Thorax. Scutum black, with widely yellow margins; black pilose, except yellow pilose along anterior and posterior margins and with medially interrupted fascia of yellow pile along transverse suture. Postpronotum, postalar callus and scutellum yellow and yellow pilose. Pleurae yellow, except posterior part of anepimeron, dorsal and ventral part of katapisternum, and meron brown. Anepisternum yellow pilose anterodorsally, bare posteriorly. Anepimeron yellow pilose dorsally. Katatergum and anatergum long and short microtrichose, respectively. Other pleurae bare. Metanotum black. Calypter and halter yellow.

Wing: hyaline, with a wide tinge on apical half and with a brown macula anteromedially, between costal vein and base of vein R2+3; microtrichose, except bare on 1st costal cell, extreme base of 2nd costal cell, entirely on cell br (except microtrichose along vena spuria), basal 2/5 of cell bm, basal 1/5 of cell cup, basomedian 1/2 of alula. Legs: Yellow, except hind femur and tibia mostly brownish; yellow pilose, except most of hind leg black pilose. Hind tibia strongly widened; with long, dense black pilosity posteriorly. Coxae and trochanters yellow and yellow pilose.

Abdomen. Tergites 1–2 pale brown. Tergites 3–4 dark brown, except yellow along posterior margin. Tergites yellow pilose, except tergite 2 laterally with black pile intermixed. Sternites 1–3 dark brown. Sternite 4 yellow, except dark brown anteriorly. Sternite 1 bare, other sternites yellow pilose. Genitalia as in fig. 241.

Etymology. The specific epithet is composed of the Latin words *macula* (spot, mark) and *penna* (wing). This name refers to the brown macula on the wing of this species.

Diagnosis. This is the only known species of *Stipomorpha* with a dark brown macula anteromedially on the wing.

Distribution. Only known from Argentina.

***Stipomorpha mendax* spec. nov.**

Figs 182–185, 242.

Studied type specimens. HOLOTYPE. SURINAM. Male. Label 1: “Marowijne, near Perica, E-W verbinding, road 21. 25 km E of Commewijne river. Malaise trap. 6–20.VIII.1997. Leg. B. de Dijn. Coll. RMNH.

PARATYPES (all in coll. RMNH). SURINAM: 1 male, Paramaribo, Charlesburg, Krepi, 21.I.1964, leg. D.C. Geijskes; 1 male, Paramaribo, Ma Retraite, 14.I.1964, leg. D.C. Geijskes; 1 male, Paramaribo, Ma Retraite, 9.II.1964, leg. D.C. Geijskes; 1 female, Kwatta, 1.II.1964, leg. D.C. Geijskes.

Additionally studied material. FRENCH GUYANA, 1 female, Montagne de Kaw, Camp Patawa, 11.XII.2002, leg. V. Soon, coll. RMNH.

Description (based on holotype)

Adult male. Body size: 6 mm.

Head. Face occupying about 1/4 of head width in frontal view; shining yellow with vague, narrow median dark vitta; with whitish pilosity, most dense sublaterally and ventrally, very sparse medially; with narrow strip of white pubescence along eye margins. Gena hardly developed, eyes almost directly bordering oral margin; yellow. Lateral oral margins not produced; not reaching below eye margin in lateral view. Frons about as long as width of lunula; brown; yellow pilose laterally. Vertex shining black; black pilose. Occiput black; yellow pilose dorsally, white pilose ventrally. Eye very sparsely and short pilose, with pili about as long as ommati diameter, appearing bare under low magnification. Antennal fossa about as wide as high. Antenna brown. Antennal ratio approximately as 3:1:3.5; basoflagellomere parallel-sided with rounded apex. Arista slender, about 2/3 of length of basoflagellomere.

Thorax. Scutum black dorsally, with margins widely yellow; yellow pilose, except for two patches of black pile posterior to transverse suture. Postpronotum, postalar callus and scutellum yellow and yellow pilose. Scutellum without calcars; slightly sulcate apicomediaally. Anepisternum weakly convex, without distinction between anterior and posterior part; yellow; yellow pilose anteriorly, bare posteriorly. Anepimeron brownish; yellow pilose dorsally, bare ventrally. Katatergum and anatergum brown; long and short microtrichose, respectively. Katepimeron yellow; bare. Katepisternum brown; bare. Calypter and halter yellow.

Wing: hyaline, without colouration, microtrichose except bare basally on cell r1 along vein RS, on basal 3/4 of cell br, posterobasal 1/2 of cell bm, basomedian 1/6 of alula.

Legs: Yellow, except hind tibia blackish with narrowly yellow base and first three tarsomeres of hind tarsus brown; yellow pilose, except tarsi dorsally black pilose and hind tibia long black pilose on apical 3/4. Coxae and trochanters yellow and yellow pilose.

Abdomen. Yellow and short yellow pilose. Second tergite slightly wider than thorax, widest at basal 1/3; third and fourth tergites much narrower. Sternites yellow; sparsely yellow pilose; sternite 1 bare. Genitalia as in fig. 242.

Etymology. The name *mendax* (Latin for lying, deceiving) was chosen in analogy of the names *fraudator* (Latin for cheating, deceiving) and *spuria* (Latin for false), two very similar-looking species of *Stipomorpha*.

Diagnosis. Orange-brown species with black vertex, very similar in size and colouration to *S. fallax*, *S. fraudator* and *S. spuria*. This species is keyed out in the key three times, because of variability in microtrichosity on the alula, variability in overall colouration and colouration of pilosity, and because of sexual dimorphism in wing colouration. In some specimens the alula is entirely microtrichose, while in others it has a small bare area basomedially. The only female identified as this species (see additionally studied material) differs from the males in the presence of a faint whitish cloud in the apical half of the wing (view against dark background). In other characters it is similar to the male.

Other varying characters: in some specimens the vertex is black pilose (as in the holotype), in others it is yellow pilose. The extent of black pile on the scutum also varies, as well as the extent of dark colouration on the abdomen. However, no discrete differences were found and the genitalia are similar in all male specimens.

Distribution. Only known from Surinam.

***Stipomorpha micromidas* (Shannon, 1925)**

Figs 186–190, 243.

Microdon micromidas Shannon, 1925: 112.

Studied type specimens. HOLOTYPE. PANAMA. Female. Label 1: "Taboga, Panama"; label 2: "23 Febr. 1912"; label 3: "A. Busck coll."; label 4: "Type no. 27833 U.S.N.M"; label 5: "Microdon micromidas Snn." Coll. USNM.

Additionally studied specimen. COSTA RICA. Male. Label 1: "COSTA RICA. Prov. Guanacaste, P.N. / Rincón de la Vieja, Send. a las aguas / termales, 900-1000 m. 6–7 OCT / 2001. D. Briceño. Red con Aguamiel. / L_N_305843_392970 #64950"; label 2 (barcode): "INB0003380896 / INBIOCRI COSTA RICA"; label 3 (red): "Ultimo especimen en / BD A. Lépez / 2-7-2002" / other side: "? MCR-25". Coll. INBIO.

Redescription (based on holotype)

Adult female. Body size: 6.5 mm.

Head. Face occupying slightly less than 1/3 of head width in frontal view; shining yellow; entirely yellow pilose, most densely laterally and ventrally, very sparse medially. Gena hardly developed, eyes almost directly bordering oral margin; yellow. Lateral oral margins not produced; not reaching below eye margin in lateral view. Frons and lunula black; frons yellow pilose. Vertex convex; shining black; yellow pilose. Occiput black; yellow pilose dorsally, white pilose ventrally. Eye very sparsely and short pilose, appearing bare under low magnification. Antennal fossa about as wide as high. Antenna brown; antennal ratio 3:1:5.5; basoflagellomere parallel-sided with narrowly rounded apex, with sensory pit located at 3/4 from base; arista slender, about 3/4 of length of basoflagellomere.

Thorax. Scutum shining black; yellow pilose, except for three patches of black pile on anterior half and two patches of black pile on posterior half. Postpronotum and postalar callus black; yellow pilose. Scutellum black anteriorly, yellow along posterior margin; yellow pilose. Pleurae blackish. Anepisternum yellow pilose anteriorly

and posteriorly, widely bare in between. Anepimeron entirely yellow pilose. Katepisternum yellow pilose dorsally, bare ventrally. Katepimeron bare. Katatergum and anatergum short and long microtrichose, respectively. Calypter and halter yellow.

Wing: hyaline, tinged with yellow, with greyish tip (apical of vein M1) and greyish fascia along vein dm-cu; microtrichose, except bare on basal 1/8 of cell cup.

Legs: Front and mid legs yellow and yellow pilose. Hind leg yellow, except apical 1/2 of tibia and basal two tarsomeres of tarsus blackish (3rd tarsomere intermediately coloured); yellow pilose, except black pilose on blackish parts and on entire dorsal part of hind tarsus. Coxae and trochanters blackish yellow; yellow pilose.

Abdomen. Brownish yellow; short yellow pilose. Second tergite slightly wider than thorax, widest at basal 1/3; third and fourth tergites much narrower. Sternites yellow; bare, except sternite 4 yellow pilose.

Description of male. As female, except for following differences.

Adult male. Body size: 8 mm.

Head. Face occupying about 1/4 of head width in frontal view. Frons about as long as width of lunula. Vertex black; golden yellow pilose, except black pilose posteriad of ocellar triangle.

Thorax. Scutellum black; yellow pilose.

Wing: greyish fascia along vein dm-cu hardly visible.

Legs: Hind femur yellow and yellow pilose, except for vaguely demarkated dark ring around middle 1/3, with patch of black pile anteriorly. Hind tibia black and black pilose, except yellow and yellow pilose on basal 1/5. Hind tarsus with tarsomeres 1–2 black, tarsomere 3 brownish, tarsomeres 4–5 yellow; black pilose.

Abdomen. Genitalia as in fig. 243.

Diagnosis. The only known *Stipomorpha*-species with a pilose dorsal part of the katepisternum and a completely yellowish abdomen. The wing tip is greyish, in contrast with the yellowish basal 2/3 of the wing.

Distribution. Costa Rica and Panama.

***Stipomorpha mixta* (Curran, 1940)**

Figs 191–195, 244.

Microdon mixtus Curran, 1940: 6.

Studied type specimens. HOLOTYPE. GUYANA. Female. Label 1 (small round, red-bordered label): "Holotype"; label 2: "British Guyana: Cuyuni R., Kamaria Landing. 22.XI.1929. Oxf. Univ. Expedn. B.M. 1929-485."; label 3: "6250"; label 4 (red): "*Microdon mixtus* Curran Holotype female"; label 5: "*Microdon mixtus* Curran Det. C.H. Curran"; label 6: "Note 389". Coll. BMNH.

Additionally studied specimens. FRENCH GUYANA: 1 male & 1 female, Kaw Mountains, 04°32.893'N–52°10.245'W, 30.XII.2002, leg. V. Soon, coll. RMNH; GUYANA: 1 male, Kurupkari: 4°40'N, 58°40'W, sept.-nov. 1992, leg. BMNH(E)2006-132, coll. BMNH; SURINAM: 1 female, Brownsberg, 04°56'45"N–55°10'59"W, 2.VI.2006, leg. M. Reemer, coll. RMNH.

Redescription (based on holotype)

Adult female. Body size: 8.5 mm.

Head. Face occupying almost 1/3 of head width in frontal view; shining yellow with yellow pilosity, a little longer around oral margin. Gena yellow. Oral cavity with lateral margins not produced and not notched anteriorly. Frons black; yellow pilose, except for bare triangular part posterior to lunula. Vertex black; yellow pilose. Occiput black; black pilose dorsally, yellow pilose laterally, gradually getting white ventrally. Eye very sparsely and short pilose, with pili about as long as ommati diameter, appearing bare under low magnification. Antennal fossa about as wide as high. Antenna brown, scape yellow on basal 3/4. Antennal ratio 6:1:6; basoflagellomere parallel-sided with narrowly rounded apex, with sensory pit located at 2/3 from base, just ventrad of a groove that ranges from the base of the arista to the apex. Arista slender, about 2/3 of length of basoflagellomere.

Thorax. Yellow, except blackish brown on scutum, leaving wide yellow lateral margins. Scutum yellow pilose, except for lateral patches of dark pile posterior to transverse suture. Postpronotum, postalar callus and scutellum yellow and yellow pilose. Scutellum without calcars. Anterior and posterior part of anepisternum not differentiated; anterior part yellow pilose, posterior part bare. Anepimeron yellow pilose dorsally. Katatergum and anatergum pilose and microtrichose, respectively. Other pleurae bare. Calypter and halter yellow.



FIGURES 186–200. 186–187. *Stipomorpha micromidas* male (Costa Rica, coll. INBIO). 186. habitus dorsal; 187. habitus lateral. 188–190. *Stipomorpha micromidas* female (holotype). 188. habitus dorsal; 189. habitus lateral; 190. head lateral. 191–195. *Stipomorpha mixta*. 191. male (Surinam, coll. RMNH), habitus dorsal; 192. female (holotype), habitus dorsal; 193. female (holotype), habitus lateral; 194. female (holotype), head frontal; 195. female (holotype), head lateral. 196–200. *Stipomorpha panamana* male (holotype). 196. habitus dorsal; 197. habitus lateral; 198. head frontal; 199. head lateral; 200. wing.

Wing: hyaline, tinged with yellow in and posterior to costal cell and in and posterior to pterostigma. Microtrichose except bare on posterobasal 1/4 of cell br, basal 1/2 of cell bm.

Legs: Yellow and yellow pilose, except with long black pile on apical 1/2 of hind-tibia and dorsally on hind tarsi. Hind-tibia strongly widened, widest point around 1/2. Coxae and trochanters yellow and yellow pilose.

Abdomen. Yellow. Tergite 2 wider than thorax, widest at posterior margin; tergites 3 and 4 about as wide as thorax. Tergite 1 and 2 black pilose. Tergite 1 with anteromedian smooth, concave area. Tergite 3 black pilose, with some yellow pile laterally. Tergite 4 black pilose anteriorly and medially (in the shape of a T with a wide cross-bar), yellow pilose laterally and posteriorly. Sternite 1 narrow and bare, separated from sternite 2 by a membrane of about the width of sternite 1. Sternite 2 pilose, laterally twice as wide as medially, separated from sternite 3 by a membrane of about the median width of sternite 2. Sternite 3 and 4 pilose, not separated by membrane

Male. (See also notes below!) Body size: 7 mm. As female, except for following characters (based on 1 specimen). Face occupying about 1/4 of total head width in frontal view. Frons yellow anteriorly, black posteriorly; mixed yellow and black pilose. Antenna yellowish brown; basoflagellomere slightly longer than scape; antennal ratio approximately as 4:1:5. Wing bare on basal 1/3 of cell r1, entirely on cells R and BM, basal 1/3 of cell cup. Tergite 3 yellow pilose, except black pilose laterally. Tergite 4 yellow pilose along anterior margin, on median 1/3 and along posterior margin, black pilose on lateral 1/3. Genitalia as in fig. 243.

Diagnosis. From other *Stipomorpha* species with a yellow scutellum and abdomen, *S. mixta* differs by the following characters: posterior part of anepisternum bare, alula entirely microtrichose, wing uniformly hyaline, vertex black, tarsi entirely yellow, anterior margin of tergite 2 not curled around tergite 1 laterally.

Notes. A very variable species in colouration of pilosity, antennal ratio and extent of microtrichosity on wings. Possibly this variability indicate that there is more than one species involved, but this could not be determined with the available material.

The female specimen from Surinam was captured after the collector saw it tumbling down (from the canopy?) on shrub leaves along a narrow path in dense primary forest.

Distribution. Known from Guyana, Surinam and French Guyana.

Stipomorpha panamana spec. nov.

Figs 196–200, 245.

Studied type specimens. HOLOTYPE. PANAMA. Male. Label 1: "El Cermeno / Pan. I-IV.41 / Fly trap / Zetek no. 4775"; label 2: "LotNo / 41-7233". Coll. USNM.

Redescription (based on holotype)

Adult male. Body size: 8 mm.

Head. Face occupying about 2/5 of head width in frontal view; yellowish brown, with oblique blackish line from antennal fossa to eye margin; yellow pilose, short dorsally and medially, long ventrally and ventrolaterally. Gena yellow. Oral cavity with lateral margins a little produced and notched anteriorly. Frons about as long as width of lunula; blackish posterior to lunula, yellow laterally; yellow pilose. Vertex swollen, in profile produced for about 1/4 of height of eye; yellowish brown; yellow pilose, except for fascia of black pile anteriorly and medially interrupted fascia of black pile along posterior margin. Occiput yellow, except black dorsolaterally; yellow pilose. Eye bare. Antennal fossa about as wide as high. Antenna brown, with scape yellowish on basal half. Antennal ratio 4:1:6; basoflagellomere parallel-sided with rounded apex, with sensory pit at approximately half its length. Arista slender, about 3/4 of length of basoflagellomere.

Thorax. Scutum black, with widely yellow margins; yellow pilose, except for four patches of black pile, two of which anterior of and two posterior of transverse suture. Postpronotum, postalar callus and scutellum yellow and yellow pilose. Scutellum without calcars. Pleurae yellow dorsally, brownish ventrally. Anepisternum weakly convex, without distinction between anterior and posterior part; yellow pilose anterodorsally and along posterodorsal margin. Anepimeron yellow pilose dorsally. Katatergum and anatergum yellow pilose and microtrichose, respectively. Other pleurae bare. Calypter and halter yellow.

Wing: hyaline, with whitish tinge on apical half (only visible against dark background), costal and subcostal cells brownish; microtrichose except bare on basal 1/6 of cell r1, on basal 1/2 of cells br and bm, on basal 1/10 of cell cup and basomedially on alula.

Legs: yellow; yellow pilose, except hind tibia dorsally mixed black and yellow pilose, and hind tarsus dorsally black pilose. Coxae and trochanters yellow and yellow pilose.

Abdomen. Tergites yellowish brown, except tergite 3 dark brown on anterior 2/3 and tergite 4 dark brown over most of dorsal surface (yellow along posterior and lateral margins); yellow pilose. Sternites brown; sternites 1 and 2 bare, sternites 3 and 4 yellow pilose. Genitalia as in fig. 245.

Diagnosis. Yellowish brown species with swollen, yellow vertex. Anepisternum with pile along posterodorsal margin, katepisternum dorsally bare, alula partly bare, wing without anteromedial spot, face wider than an eye.

Distribution. Only known from Panama.

Stipomorpha puerilis (Doesburg, 1966)

Figs 201–203.

Ubristes puerilis Doesburg, 1966: 86.

Studied type specimens. HOLOTYPE. SURINAM. Female. Label 1 (yellow): "HOLOTYPE"; label 2: "Zanderij Surin. / 23-9-1960 / P.H. v. Doesburg Jr."; label 3: "Ubristes / nanus Dsb. [female sign] / det. v. Doesburg"; label 4: see notes. Coll. RMNH.

Additionally studied specimens. VENEZUELA: 1 female, T.F. Amaz., Cerro de la Neblina, 140 m., 0°50'N–66°10'W, 21–28.II.1985 (malaise trap in rain forest), leg. P.J. & P.M. Spangler, R.A. Faitoute & W.E. Steiner, coll. USNM.

Notes. No specimen labelled as this species could be found in the RMNH collection, but there is a specimen labelled as '*Ubristes nanus* Doesburg'. This specimen agrees with the description of *Ubristes puerilis* and is from the same date and locality as the type of *U. puerilis*. Presumably Van Doesburg first intended to name the species *nanus*, but later changed his mind without correcting the label. Under this assumption, a new label was added to the pin by the present author: 'HOLOTYPE *Ubristes puerilis* Van Doesburg, 1966'.

Distribution. Known from Surinam and Venezuela.

Stipomorpha simillima (Hull, 1950)

Figs 204–208, 246.

Microdon simillimus Hull, 1950: 611.

Studied type specimens. HOLOTYPE. GUYANA. Male. Label 1 (small round, red-bordered): "Holotype"; label 2: "Dark forest"; label 3: "A in cop. with B"; label 4: "British Guyana: Essequibo R., Moraballi Creek. 17.X.1929. Oxf. Univ. Expedn. B.M. 1929-485."; label 5 (red): "Holotype *Microdon simillima* Hull". Coll. BMNH.

PARATYPES. GUYANA. Male. Label 1 (small, round, yellow-bordered): "Paratype"; label 2: "Dark forest."; label 3: "British Guyana: Essequibo R., Moraballi Creek. 17.X.1929. Oxf. Univ. Expedn. B.M. 1929-485."; label 4 (yellow): "Paratype *Microdon simillima* Hull". Coll. BMNH.

Male. Label 1: "British Guiana: / Essequibo R., / Moraballi Creek / 25.ix.1929. / Oxf. Univ. Expedn. / B.M. 1929-485."; label 2 (yellow): "Paratype / Microdon / simillima / Hull". Coll. CNC.

Additionally studied specimens. BRAZIL: 1 male, Belém, Pará, 17.V.1967, leg. Y. Sedman, coll. USNM; FRENCH GUYANA: 1 male, Crique Sapokal Degrad Laurens, 95 m., N 3°16' / W 52°41', 26.viii–2.ix.2000, malaise trap, leg. A.E.I. Guyane, coll. J.T. Smit; PERU: 1 female, Madre de Dios, Rio Tambopata, Sachavacayoc centre, N 12°51'20.1" / W 69°22'20.1", 9–14.X.2010, malaise trap, leg. & coll. J.T. Smit; 1 female, Madre de Dios, Rio Tambopata, Sachavacayoc centre, N 12°51'46.4" / W 69°21'46.6", 14–20.X.2010, malaise trap, leg. & coll. J.T. Smit.

Redescription (based on holotype)

Adult male. Body size: 6–7 mm.

Head. Face occupying 1/4 of head width in frontal view; shining yellow with a pale brown median stripe from oral margin to just below antennae, gradually narrowing upward; face with white pilosity, a little longer around oral margin, except bare on median stripe. Gena blackish. Occiput black; black pilose dorsally, getting white laterally

and ventrally. Oral cavity with hardly produced lateral margins. Frons and lunula black and short black pilose, except for bare triangular part posterior to lunula. Vertex black; black pilose. Eye very sparsely and short pilose, with pili about as long as ommati diameter, appearing bare under low magnification. Antennal fossa about as wide as high. Antenna blackish brown, except scape yellow on interior sides; scape and pedicel dark pilose; antennal ratio 4:1:3.5; basoflagellomere parallel-sided with narrowly rounded apex, with sensory pit located at 3/4 from base, within a vague groove that ranges from just before 1/2 to just after the pit; arista slender, about 2/3 of length of basoflagellomere, very shortly pilose, appearing bare under low magnification.

Thorax. Black, more brownish on pleurae. Postpronotum, scutum, postalar callus and scutellum short black pilose, except a few pale pile on scutellum (and possibly along transverse suture of scutum, but not well visible in type specimen). Scutellum semicircular, without calcars. Anepisternum more or less flat, pilose anterodorsally. Anepimeron pilose posterodorsally. Katepisternum and katepimeron bare. Calypter grey, halter yellowish.

Wing: hyaline, with faint yellowish cloud on and posterior to pterostigma and with faint dark cloud between apex of costal cell and vena spuria, and microtrichose except bare on basal 3/4 of cell br, basal 1/2 of cell bm, anterobasal 1/4 of cell cup.

Legs: brownish black, except anterior four tarsi yellow with first tarsomere more brownish. Legs black pilose, except anterior four tibiae pale pilose and hind-tibia pale pilose on basal 1/3; pile on hind-tibia a little longer than half the width of the tibia. First tarsomere of hind-tarsus as long as 1/3 of length of hind-tibia, a little wider than apex of tibia, twice as long as wide (dorsal view). Front-coxae and trochanters pale pilose; mid- and hind-coxae and -trochanters black pilose.

Abdomen. Blackish (but see diagnosis). Second segment wider than thorax, widest point at half the length; third and fourth tergites strongly narrowing. Tergite 1 black pilose laterally; with anterolateral 'ridges'; with anteromedian smooth, concave area. Tergite 2 black pilose laterally, pale pilose dorsally. Tergite 3 black pilose. Tergite 4 black pilose, lateral margins with pale pilosity, which is posteriorly connected with two submedial lines of pale pile on posterior half of the tergite. Sternite 1 narrow and bare, separated from sternite 2 by a membrane of twice the width of sternite 1. Sternite 2 bare, laterally twice as wide as medially, separated from sternite 3 by a membrane of 1.5 times the median width of sternite 2. Sternite 3 and 4 pilose, not separated by membrane. Genitalia as in fig. 246.

Female. As male, except for usual sexual differences.

Diagnosis. From other *Stipomorpha*-species with a black thorax, *S. simillima* can be recognized by the following characters: face largely yellow with narrow median brown stripe, basoflagellomere about as long as or slightly shorter than the scape, anepisternum only pilose anterodorsally, katepisternum bare, structure of male genitalia. The colouration of the abdomen appears to be quite variable, from blackish in the holotype to pale brown in the paratype.

Notes. The fourth label on the holotype reads "A in cop. with B". Apparently the male holotype has been collected in copula with a female that has later been labelled as specimen B. The whereabouts of this female are unknown. In the original description Hull (1950) only mentions two male paratypes, one of which is present in the BMNH collection.

Distribution. Known from Guyana, French Guyana, Brazil and Peru.

Stipomorpha spuria spec. nov.

Figs 209–211, 247.

Type specimens: HOLOTYPE. SURINAM. Male. Commewijne, Peperpot. 05°46'08"N-55°07'33"W. 20.IV.2006. Leg. M. Reemer. Coll. RMNH.

Description (based on holotype)

Adult male. Body size: 7 mm.

Head. Face occupying about 1/3 of head width in frontal view; shining yellow with whitish pilosity, most dense sublaterally and ventrally, almost bare medially. Gena developed, about as wide as 2nd antennal segment; yellow. Oral margins laterally a little produced; in lateral view reaching below ventral eye margin. Frons about as long as lunula; yellow; yellow pilose laterally. Vertex a little swollen; shining black; yellow pilose. Occiput black; yellow pilose dorsally, whitish pilose ventrally. Eye bare. Antennal fossa about as wide as high. Antenna pale

brown. Antennal ratio approximately as 3:1:4; basoflagellomere parallel-sided with rounded apex, with small sensory pit at 2/3 from base. Arista black; slender; about 3/4 of length of basoflagellomere.



FIGURES 201–211. 201–203. *Stipomorpha puerilis* female (holotype). 201. habitus dorsal; 202. habitus lateral; 203. head frontal. 04–208. *Stipomorpha simillima* male (holotype). 204. habitus dorsal; 205. habitus lateral; 206. head frontal; 207. head lateral; 208. wing. 209–211. *Stipomorpha spuria* male (holotype). 209. habitus dorsal; 210. habitus lateral; 211. face frontal.

Thorax. Scutum black, but widely yellow along margins and narrowly along transverse suture; golden yellow pilose, except for medially interrupted fascia of black pile posteriorly of transverse suture. Postpronotum, postalar callus and scutellum yellow and yellow pilose. Scutellum semicircular, without calcars. Pleurae yellow, except ventral part of katapisternum, meron, dorsal part of katatergum and metanotum blackish. Anepisternum weakly convex, without distinction between anterior and posterior part; yellow pilose anterodorsally and along posterior margin. Anepimeron entirely yellow pilose. Katatergum and anatergum long and short microtrichose, respectively. Other pleurae bare. Calypter and halter yellow.

Wing: hyaline, without colouration, microtrichose except bare on basal 1/4 of cell br, basal 1/4 of cell bm, basal 1/5 of cell cup and basomedian 1/6 of alula.

Legs: Front- and mid-legs yellow and yellow pilose, except mid-femur partly black pilose posteriorly. Hind leg yellow, except apical 1/2 of tibia and first two tarsomeres darkened; femur yellow pilose, except black pilose anteriorly on basal 1/2; tibia yellow pilose on basal 1/3, black pilose on apical 2/3; first three tarsomeres black pilose dorsally, last two tarsomeres yellow pilose. Coxae and trochanters yellow and yellow pilose, except hind coxa black pilose anteriorly.

Abdomen. Yellow; entirely yellow pilose. Second tergite about as wide as thorax, widest at basal 1/3; third and fourth tergites much narrower. Sternites yellow; sparsely yellow pilose, except sternite 1 bare. Genitalia as in fig. 247.

Female. Unknown.

Etymology. The name *spuria* (Latin for false) was chosen in analogy of the names *fallax*, *fraudator* and *mendax*, which have approximately the same meaning, in order to stress the similarity of these species.

Diagnosis. Orange-brown species with black vertex, very similar in size and colouration to *S. fallax*, *S. fraudator* and *S. mendax*. Diagnostic characters: anepisternum with posterodorsal margin pilose, katapisternum dorsally bare, alula partly bare, wing without distinctly yellow parts.

Distribution. Only known from the holotype from Surinam.

***Stipomorpha tenuicauda* (Curran, 1925)**

Figs 212–215, 248.

Microdon tenuicaudus Curran, 1925: 339.


Studied type specimens. HOLOTYPE. PERU. Female. Label 1: "El Campamiento. Col. Perene. PERU. Jult '20"; label 2: "Cornell Univ. Expedition Lot 607"; label 3 (red): "Type *Microdon tenuicauda* Curran"; label 4 (red): "Holotype Cornell U. No. 1736"; label 5: "*Microdon tenuicauda* Curran Det. C.H. Curran". Coll. CU.

Additionally studied specimens. BOLIVIA: 1 male, La Paz Prov., Mapiro, Arroyo Tuhiri, 15°17'26"S–68°15'46"W, 12.IV.2004, leg. & coll. M. Hauser; 1 male with same locality info, but 13.IV.2004, coll. RMNH; BRAZIL: 1 male, Belém, Pará, 26.IV.1967, leg. Y. Sedman, coll. USNM; COSTA RICA: 4 females, Puntar, Golfo Dulce, 3 km SW Rincón, 10 m, 1989–1990, leg. Hanson, coll. USNM; ECUADOR: 1 female, Napo, Jatun Sacha Res., 6 km E Misahualli, 450 m, 30.IV–8.V.2002, leg. S.A. Marshall, DNA voucher specimen no. S259, G. Ståhls, FMNH Helsinki, coll. USNM; FRENCH GUYANA: 1 male, Patawa, 4°32.658'N–52°9.132'W, leg. O. Morvan, coll. CNC; PERU: 1 female, Cusco, River Ceosnibata, 12.XII.1951, leg. Woytkowski, coll. CNC; 1 female, Avispas, Madre de Dios, 400 m, 20–30.IX.1962, leg. L. Pena, coll. USNM.

Redescription (based on holotype)

Adult female. Body size: 9 mm.

Head. Face occupying almost 1/3 of head width in frontal view; shining yellow with yellow pilosity, a little longer around oral margin. Gena yellow. Oral cavity with lateral margins produced and notched anteriorly. Frons black; yellow pilose, except for yellow bare triangular part posterior to lunula. Vertex black; yellow pilose; convexly produced. Occiput black; yellow pilose. Eye very sparsely and short pilose, appearing bare under low magnification. Antennal fossa about as wide as high. Antenna yellowish brown, pedicel and basoflagellomere a little darker. Antennal ratio 4:1:5; basoflagellomere parallel-sided with narrowly rounded apex, with sensory pit located at apical 1/4. Arista slender, about 3/4 of length of basoflagellomere.



FIGURES 212–221. 212–215. *Stipomorpha tenuicauda* female (holotype). 212. habitus dorsal; 213. habitus lateral; 214. head frontal; 215. head dorsal. 216–219. *Stipomorpha trigoniformis* male (holotype). 216. habitus dorsal; 217. habitus lateral; 218. head frontal; 219. wing. 220–221. *Stipomorpha wheeleri* male (paratype). 220. habitus dorsal; 221. habitus lateral.

Thorax. Scutum shining black, except yellow along margins, widely so around postpronotum and postalar callus; appressed golden yellow pilose, except erect along anterior and posterior margin and with lateral fasciae of half-erect black pile along transverse suture. Postpronotum and postalar callus yellow and yellow pilose. Scutellum yellow; yellow pilose anteriorly, otherwise black pilose. Pleurae yellow, except katepimeron posteriorly and katatergum anteriorly blackish. Anterior and posterior part of anepisternum not differentiated, more or less convex; anterior part yellow pilose, posterior part yellow pilose along posterior margin. Anepimeron yellow pilose dorsally. Katatergum and anatergum long and short microtrichose, respectively. Other pleurae bare. Metanotum shining blackish, except yellow on dorsal 1/3. Calypter and halter yellow.

Wing: hyaline, tinged yellow, especially on anterobasal 1/2; microtrichose except bare on basal 1/2 of cell br, posterobasal 1/4 of cell bm, basal 1/4 of cell cup and on at least 90% of alula.

Legs (including coxae and trochanters) yellow and yellow pilose, except: hind tibia whitish pilose on basal 3/5 and with black ground colour and black pilose on apical 2/5; hind tarsus with first two tarsomeres black and black pilose. Metatibia strongly widened, widest point around 1/2.

Abdomen. Yellow and yellow pilose. Second tergite about as wide as thorax, widest at posterior 1/3; other tergites clearly narrower. Sternite 1 narrow and bare, separated from sternite 2 by a membrane of about the width of sternite 1. Sternite 2 pilose, laterally twice as wide as medially, separated from sternite 3 by a membrane of about twice the median width of sternite 2. Sternite 3 and 4 pilose, not separated by membrane. Genitalia as in fig. 248.

Male. As female, except for usual sexual differences.

Diagnosis. Distinguishable from the very similar *S. mackiei* reliably only by the male genitalia. The character of the distribution of microtrichia on the alula, as mentioned in the key, could be verified in a few males only. Whether it works for all specimens (including females) is uncertain.

Distribution. Known from Brazil, Costa Rica, Ecuador, French Guyana and Peru.

***Stipomorpha trigoniformis* (Shannon, 1927)**

Figs 216–219, 249.

Microdon (Ubristes) trigoniformis Shannon, 1927: 19. Type locality:

Studied type specimens. HOLOTYPE. BRAZIL. Male. Label 1 (small, round, red-bordered): "Holotype"; label : "Villa Nova", other side: "55 37"; label 3: "*Microdon Ubristes trigoniformis* Snn.". Coll. BMNH.

Additionally studied specimens. FRENCH GUYANA: 1 male, Kaw mountains, 04°33.562'N–52°12.425'W, 21.X.2002, leg. V. Soon, coll. RMNH.

Redescription (based on holotype)

Adult male. Body size: 7.5 mm.

Head. Face occupying 1/4 of head width in frontal view; shining black with yellow lateral margins, the black part occupying 2/3 of face; face with white pilosity, bare on median 1/3. Gena hardly developed, directly bordering oral cavity. Oral margins black, laterally produced and anteriorly notched. Frons and vertex black and short black pilose, except for bare triangular part posterior to lunula. Occiput black; black pilose on dorsal half, white on ventral half. Eye sparsely and short pilose, with pili about as long as ommati diameter, appearing bare under low magnification. Antennal fossa about as wide as high. Antenna blackish brown; antennal ratio 3:1:3.5; basoflagellomere parallel-sided with narrowly rounded apex; arista slender, about 3/5 of length of basoflagellomere.

Thorax. Black. Postpronotum, scutum, postalar callus and scutellum short black pilose, except a few pale pile along transverse suture. Scutellum semicircular, without calcars. Anepisternum more or less flat, pilose anterodorsally. Anterior part of anepimeron pilose dorsally. Katepisternum and katepimeron bare. Calypter grey, halter blackish.

Wing: hyaline, with faint brownish hue all over; microtrichose except bare on basal 2/3 of cell br, basal 1/8 of cell bm, anterobasal 1/4 of cell cup.

Legs: brownish black, except fifth tarsomeres of all tarsi yellow. Legs black pilose, except tibiae pale pilose basally (extent hard to assess in holotype); pile on hind-tibia a little longer than half the width of the tibia. First tarsomere of hind-tarsus as long as 1/3 of length of hind-tibia, a little wider than apex of tibia, 1.5 times as long as wide (dorsal view). Coxae and trochanters black pilose.

Abdomen. Blackish brown. Second segment slightly wider than thorax, widest point at anterior 1/3; third and fourth tergites strongly narrowing. Tergite 1 black pilose anterolaterally; with anteromedian smooth, concave area. Tergite 2 black pilose anterolaterally, pale pilose laterally and dorsally. Tergite 3 and 4 pale pilose. Sternite 1 bare. Genitalia as in 249.

Female. Unknown.

Diagnosis. From other *Stipomorpha*-species with a black thorax, *S. trigoniformis* can be recognized by the following characters: katapisternum bare, alula entirely microtrichose, wing without whitish transverse fascia.

Distribution. Known from Brazil and French Guyana.

***Stipomorpha wheeleri* (Mann, 1928)**

Figs 220–224, 250.

Microdon wheeleri Mann, 1928.

Studied type specimens. Two specimens labelled as types (red labels) in USNM-collection: 1 male & 1 female; both with same labels: label 1: "no. 147"; label 2: "Red Tank, C.Z. 2.27.23, W.M. Wheeler". The female has an additional label stating: "*Microdon wheeleri* Mann type". Mann (1928) stated that he designated a 'type and allotype'. As his description is based primarily on the female, which also carries a label stating 'type', this is regarded as the holotype. There are also two specimens (same locality and date) labelled as paratypes on blue labels, as well as four empty puparia, from which the specimens were reared.

Redescription (based on holotype)

Adult female. Body size: 8 mm.

Head. Face occupying slightly more than 1/3 of head width in frontal view; shining pale yellow; yellow pilose on lateral 1/4, bare medially. Gena yellow. Oral margins not produced laterally, not notched anteriorly. Frons and vertex yellow; yellow pilose; ocellar triangle black, elevated; frontal ocellus split in two. Occiput yellow; yellow pilose. Eye bare. Antennal fossa about as wide as high. Antenna pale brown; antennal ratio 3.5:1:3; basoflagellomere parallel-sided with rounded apex, with sensory pit at 3/4 from base; arista slender, about 2/3 of length of basoflagellomere.

Thorax. Scutum shining dark brown, except yellow along margins; yellow pilose, except for a medially interrupted fascia of black pile posterior to transverse suture. Postpronotum and postalar callus yellow; yellow pilose. Scutellum yellow; yellow pilose. Anepisternum more or less flat, pilose anteriorly and posteriorly, widely bare in between; brown anteriorly, yellow posteriorly. Anterior anepimeron yellow pilose dorsally, bare ventrally. Posterior and dorsomedial anepimeron yellow; bare. Katapisternum and katepimeron yellow; bare. Meron, katatergum and anatergum brown; bare. Katatergum long microtrichose, anatergum short microtrichose. Calypter blackish, halter yellow.

Wing: hyaline, tinged with yellow; microtrichose except bare on basal 1/10 of cell r1, on most of cell br except microtrichose along vena spuria, on posterobasal 2/3 of cell bm, on basal 1/3 of cell cup.

Legs: Yellow; yellow pilose, except dorsal surface of hind tibia and basal three tarsomeres of hind tarsus black pilose. Coxae and trochanters yellow; yellow pilose.

Abdomen. Yellow; yellow pilose. Second tergite wider than thorax, widest point at 2/3 from base; third and fourth tergites strongly narrowing. Tergites 3 and 4 fused, with abrupt lateral transition (view from dorsal). Sternite 1 bare, other sternites sparsely yellow pilose.


Male (based on paratype). As female, except for following differences.

Head. Face occupying about 1/3 of head width in frontal view. Frons dark brown; yellow pilose. Vertex dark brown, black pilose. Occiput brown; black pilose dorsally, yellow pilose ventrally. Basoflagellomere with sensory pit at 2/3 from base; arista slender, about 2/3 of length of basoflagellomere.

Thorax. Scutum entirely blackish brown; entirely black pilose. Postpronotum, postalar callus and scutellum black pilose. Pilosity of pleurae as in female, but black instead of yellow.

Legs: Brown, more extensively black pilose.

Abdomen. Brown and black pilose on tergites 1, 2 and basal half of 3, then gradually getting yellow and yellow pilose. Genitalia as in fig. 250 (drawn from paratype).



FIGURES 222–232. 222–224. *Stipomorpha wheeleri*. 222. male (paratype), head frontal. 223. female (holotype), habitus dorsal; 224. female (holotype); habitus lateral. 225–227. *Stipomorpha zophera* male (holotype). 225. habitus dorsal; 226. habitus lateral; 227. head frontal. 228–229. *Stipomorpha*, tergites 1 & 2 dorsal. 228. *S. mendax*; 229. *S. mixta*. 230–232. *Stipomorpha*, male genitalia lateral. 230. *S. apicula* (holotype); 231. *S. elcopala* (holotype); 232. *S. fallax* (holotype).

Notes. The male paratype is much darker in colouration than the female holotype. Mann (1928) writes that Wheeler, who reared the specimens from their pupae, told him that all specimens were yellow at the time of emergence and darkened gradually. So, possibly the female is yellow because it is teneral.

According to Mann (1928) the pupae from which the type series was reared, were found in nests of *Crematogaster* (*Orthocrema*) *brevispinosa* Mayr subsp. *tumulifera* Forel in *Cordia alliodora* Ruiz & Pavon (Boraginaceae).

Distribution. Only known from Panama.

***Stipomorpha zophera* spec. nov.**

Figs 225–227, 251.

Studied type specimens. HOLOTYPE. FRENCH GUYANA. Male. Label 1: "FRENCH GUIANA / Montagnes Tortue / 4°15.007'N 52°21.512'W / leg. V. Soon 11.01.2003"; label 2 (red): "HOLOTYPE / *Stipomorpha zophera* / M. Reemer". Coll. RMNH.

PARATYPES: FRENCH GUYANA: 1 male same locality & leg. as holotype, but with date 9.IX.2002; 1 male, Kaw Mountains, 04°32.893'N-52°10.245'W, 8.XII.2002, leg. V. Soon, coll. RMNH.

Additionally studied specimens. ECUADOR: 1 male, Napo, Limoncocha, 15.VI.1977, leg. P.J. Spangler & D.R. Givens, coll. USNM. GUYANA: 1 male, Mazaruni-Potaro District, Takutu Mountains, 6°15'N, 59°5'W, 9.XII.1983, leg. W.E. Steiner, coll. USNM.

Description (based on holotype)

Adult male. Body size: 7 mm.

Head. Face occupying 1/4 of head width in frontal view; shining yellow with a blackish brown median stripe from antennal fossa to slightly below middle; face with yellow pilosity, replaced by longer black setae around oral margin, except bare on median stripe. Gena brown. Occiput black; black pilose dorsally, getting white laterally and ventrally. Oral cavity with lateral margins not produced. Frons and lunula black, short black pilose, except for bare triangular part posterior to lunula. Vertex black; black pilose. Eye very sparsely and short pilose, with pili about as long as ommati diameter, appearing bare under low magnification. Antennal fossa about as wide as high. Antenna brown; scape and pedicel dark pilose; antennal ratio 6:1:8; basoflagellomere with apical 1/3 clearly narrower than basal 2/3, with sensory pit located at about 3/5 from base, within a vague groove; arista slender, about 3/5 of length of basoflagellomere, very shortly pilose, appearing bare under low magnification.

Thorax. Scutum black; black pilose, except for medially interrupted fascia of yellow pile along transverse suture. Postpronotum and postalar callus pale brown; black pilose. Scutellum brown; black pilose.

Anepisternum more or less flat, black pilose anterodorsally, bare posteriorly. Anepimeron black pilose dorsally. Katepisternum and katepimeron bare. Calypter grey, halter yellowish.

Wing: hyaline; microtrichose except bare along vein RS on basal part of cell r1, entirely on cell br, on basal 4/5 of cell bm, anterobasal 1/4 of cell cup, basomedially on alula.

Legs: brownish black, except anterior four tarsi yellow and hind tarsi with last three tarsomeres yellowish brown. Legs black pilose, except hind coxa and trochanter mixed black and white pilose. Pile on metatibia a little longer than half the width of the tibia. First tarsomere of posterior tarsus as long as 2/5 of length of metatibia, a little wider than apex of tibia, twice as long as wide (dorsal view).

Abdomen. Blackish brown. Second segment wider than thorax, widest point at half the length; third and fourth tergites strongly narrowing. Tergites pale pilose, except tergites 1 and 2 laterally black pilose. Sternite 1 bare, other sternites pilose. Genitalia as in fig. 251.

Female. Unknown.


Etymology. The specific epithet *zophera* (Gr., dusky, gloomy) refers to the dark colour of this species.

Diagnosis. From other *Stipomorpha*-species with a black thorax, *S. zophera* can be recognized by the following characters: face largely yellow with narrow median brown stripe, basoflagellomere longer than scape, alula partly bare, anepisternum black pilose anterodorsally, bare posteriorly, katepisternum bare, structure of male genitalia.


Very similar to *S. inarmata*, from which it differs by: basoflagellomere longer than scape, anterior part of anepisternum black pilose, front- and mid-tibiae black pilose,

Notes. According to the label, the studied specimen from Guyana was collected "at blacklight in forest clearing near streams".


Distribution. Known from Ecuador, Guyana and French Guyana.




233




234




235




236



237




238




239


FIGURES 233–239. *Stipomorpha*, male genitalia. 233. *S. fraudator* (holotype); 234. *S. goettei* (Surinam, coll. RMNH); 235. *S. guianica* (Surinam, coll. RMNH); 236. *S. inarmata* (holotype); 237. *S. lacteipennis* (lectotype); 238. *S. lanei* (Surinam, coll. RMNH); 239. *S. litoralis* (holotype).




240




241




242



243




244




245


FIGURES 240–245. *Stipomorpha*, male genitalia. 240. *S. mackiei* (Surinam, coll. RMNH); 241. *S. maculipennis* (holotype); 242. *S. mendax* (holotype); 243. *S. micromidas* (Costa Rica, coll. INBIO); 244. *S. mixta* (Guyana, coll. BMNH); 245. *S. panamana* (holotype).




246




247




248



249



250



251

FIGURES 246–251. *Stipomorpha*, male genitalia latera. 246. *S. simillima* (holotype); 247. *S. spuria* (holotype); 248. *S. tenuicauda* (Bolivia, coll. RMNH); 249. *S. trigoniformis* (holotype); 250. *S. wheeleri* (paratype); 251. *S. zophera* (holotype).

***Ubristes flavitibia* Walker, 1852**

Figs 252–255, 265.

Ubristes flavitibia Walker, 1852: 217.

Microdon procedens Curran, 1941: 251. Syn. nov.

Microdon procteri Curran, 1941: 251. Syn. nov.

Studied type specimens. HOLOTYPE *Ubristes flavitibia*. BRAZIL. Male. Label 1: "Holotype"; label 2: "Type"; label 3: "*Ubristes flavitibia*"; label 4: "*Ubristes flavitibia* Wlk." Coll. BMNH.

HOLOTYPE *Microdon procedens*.—Male. Label 1 (red): "*Microdon procedens* Curran Holotype"; label 2: "Brasilien, Nova Teutonia, 27°11' B, 52°23' L. Fritz Plaumann. 27.10.1939". Coll. AMNH.

PARATYPE *Microdon procedens*. BRAZIL. Male. Label 1: "Brasilien, Nova Teutonia, 27°11' B, 52°23' L. 24.10.1939. Fritz Plaumann"; label 2: "Paratype, male, *Microdon procedens* Curran". Coll. USNM.

HOLOTYPE *Microdon procteri*. BRAZIL. Male. Label 1 (red): "*Microdon procteri* Curran Holotype"; label 2: "Brasilien, Nova Teutonia, 27°11' B, 52°23' L. Fritz Plaumann. 28.9.1939". Coll. AMNH.

Additionally studied specimens. BRAZIL: 1 female, Tijuca forest near Rio, 7-30.ix.1993, leg. T. Pape, coll. ZMUC.

Redescription (based on holotype *U. flavitibia*)

Adult male. Body size: 11 mm.

Head. Face occupying 1/3 of head width in frontal view; black, with two yellow submedian vittae on upper half, reaching antennal fossa, and a small yellow mark along eye margin on ventral half; with long white pilosity and a dense patch of black pile anterior to oral margin. Gena black. Oral cavity with lateral margins produced. Frons and vertex black and black pilose, with some white pile along eye margins and along transition between frons and vertex. Occiput black; grey pollinose; black pilose dorsally, white pilose ventrally. Eye very sparsely and short pilose, with pili about as long as ommati diameter, appearing bare under low magnification. Antennal fossa about as wide as high. Antenna brown, scape a little paler; antennal ratio 8:1:10; basoflagellomere parallel-sided with narrowly rounded apex, with sensory pit located at 3/4 from base, within a groove that ranges from the base of the arista to close to the apex. Arista slender, about 2/3 of length of basoflagellomere.

Thorax. Black, a little brownish on postpronotum, postalar callus, scutellum and pleurae. Scutum densely black pilose, except for some white pile along transverse suture and a small patch of white pile anterior to the scutellum. Postpronotum, postalar callus and scutellum black pilose. Scutellum without calcars. Anterior and posterior part of anepisternum divided by a weak sulcus; anterior part black pilose, posterior part black pilose along posterior margin. Anepimeron black pilose on dorsal 2/3. Katatergum and anatergum long and short microtrichose, respectively. Other pleurae bare. Calypter greyish, halter yellow with grey knob.

Wing: hyaline, brownly infuscated along anterior margin. Microtrichose except bare on posterobasal 1/2 of cell br and almost entirely on alula (only a narrow basal strip with microtrichia).

Legs: Front- and mid-femora brown, black pilose; hind-femur brown with apical 1/4 yellow, with long black pile. Front-tibia brown, black pilose; mid-tibia yellowish brown, with long black pile; hind-tibia yellow, with very long (some longer than maximal width of tibia) yellow pile on basal 3/4, long black pile on apical 1/4. Tarsi yellow, with basal tarsomeres a bit darker, black pilose dorsally, mid- and hind-tarsi ventrally with dense, short, yellow pile. Coxae and trochanters brownish black, with white pile.

Abdomen. Black. Tergite 1 with anterior half concave, laterally with white pile. Tergite 2 about as wide as thorax, with two lateral 'bulges' halfway, which mark the maximum width of the abdomen; with long white pile anterolaterally, rather long black pile on the lateral 'bulges' and short, appressed pale pile on posterior half of tergite. Tergites 3 and 4 with short black pile over entire surface and long black pile along posterior margin. Sternites 1 and 2 with long white pile; sternites 3 and 4 with mixed long black and white pile. Hypopygium yellowish. Genitalia as in fig. 265.

Female. As male, except for usual sexual differences.

Diagnosis. Within *Ubristes* s.s. this is the only known entirely black coloured species.

Notes. The types of *Microdon procedens* and *M. procteri* (8-9 mm) are smaller than the type of *Ubristes flavitibia* (11 mm). Besides that, the types of *M. procedens* differ from *U. flavitibia* only in the colour of the pile on the hind tibia: mostly brown (not black, as stated by Curran 1941) in *M. procedens*, yellow in *U. flavitibia*. However, it appears that the pilosity on the tibia of the type specimens of *M. procedens* has lost its natural colour

and turned brown. No differences could be found in external morphology or genitalia. The type of *M. procteri* is even more similar to that of *U. flavitibia*. Therefore, both *M. procedens* and *M. procteri* are here considered as junior synonyms of *U. flavitibia*.

Distribution. Known from southern parts of Brazil.

***Ubristes ictericus* spec. nov.**

Figs 256–260.

Studied type specimens. HOLOTYPE. BRAZIL. Male. Label 1: "Belem, Para / Brazil / 1-VI-1967 / Coll. Y. Sedman"; label 2 : "IN APEG"; label 3 (red): "HOLOTYPE [male sign] / Ubristes ictericus / M. Reemer". Coll. USNM.

PARATYPE. BRAZIL. Female. Same label data as holotype, except label 3: "new species 3"; label 4 (yellow): "PARATYPE [female sign] / Ubristes ictericus / M. Reemer". Coll. USNM.

PARATYPE. ECUADOR. Female. Label 1: "ECUADOR: Sucumbios / Sacha Lodge, 0.5°S. / 76.5°W. 270 m. 27VIII–10XI / 1994, Hibbs, ex: malaise"; label 2: "PARATYPE [female sign] / Ubristes ictericus / M. Reemer". Coll. SEMC.

Description (based on holotype)

Adult male. Body size: 10 mm.

Head. Face occupying about 1/2 of head width in frontal view; yellow, with very narrow, brown median line on lower 2/3; entirely white pilose. Gena yellow. Lateral oral margins produced. Frons and vertex yellow, except darkened posterior to lunula and on ocellar triangle; yellow pilose, except black pilose on ocellar triangle. Occiput black dorsally, yellow ventrally; yellow pilose dorsally, white pilose ventrally; whitish pollinose. Eye very sparsely and short pilose, with pili about as long as ommati diameter, appearing bare under low magnification. Antennal fossa about as wide as high. Antenna brown, except scape yellowish; antennal ratio 6:1:9; basoflagellomere parallel-sided with narrowly rounded apex, with sensory pit located at 5/6 from base, within a groove that ranges from the base of the arista to close to the apex. Arista slender, about 2/3 of length of basoflagellomere.

Thorax. Scutum black, except margins widely yellowish brown; medially appressed golden yellow pilose, except for pair of black pilose patches on anterior 1/4, erect yellow pilose along margins. Postpronotum and postalar callus and scutellum yellow; yellow pilose. Scutellum yellow; yellow pilose basally, black pilose apically. Anepisternum yellowish brown; yellow pilose anteriorly and along posterior margin. Anepimeron yellow pilose on dorsal half. Katatergum and anatergum long and short microtrichose, respectively. Other pleurae bare. Calypter and halter yellow.

Wing: hyaline, with faint yellowish tinge. Microtrichose, except bare on 1st costal cell, basal 1/10 of 2nd costal cell, basally on cell r1 along vein RS, entirely on cell br except microtrichose along vena spuria, basal 2/3 of cell bm, basal 1/5 of cell cup, almost entirely on alula (only a narrow basal strip with microtrichia).

Legs: yellow, except basal 3 tarsomeres of hind leg dark brown; yellow to white pilose, except black pilose on anteroventral part of hind femur, apical 1/2 of hind tibia and dorsal part of hind tarsus.

Coxae and trochanters yellow; yellowish white pilose, except for sparse black pile apically on hind coxa.

Abdomen. Yellowish brown; yellow pilose, except black pilose posterolaterally on tergite 3 and on posterior 1/4 of tergite 4. Tergite 2 about as wide as thorax, with two lateral 'bulges' halfway, which mark the maximum width of the abdomen; with long yellow pile anterolaterally, rather long yellow pile on the lateral 'bulges' and short, appressed pale pile on posterior half of tergite. Sternites with long yellow pile. Hypopygium yellowish.

Female. Body size: 9–10 mm. The female is very similar to the male, except for usual sexual differences. The paratype from Ecuador is slightly darker in colouration, with the posterior margin of the vertex dark and the hind femur and tibia somewhat darkened.

Diagnosis. Within *Ubristes* s.s. this is the only known species with an entirely yellow abdomen.

Distribution. Known from northern Brazil and Ecuador.



252



253



254




255



256



257



258





259



260

FIGURES 252–260. 252–255. *Ubristes flavitibia* male (holotype). 252. habitus dorsal; 253. habitus lateral; 254. head frontal; 255. head lateral. 256–260. *Ubristes ictericus* (holotype). 256. habitus dorsal; 257. habitus lateral; 258. tergite 2 dorsal; 259. head frontal; 260. head lateral.



FIGURES 261–266. 261–264. *Ubristes jaguarinus* male (holotype). 261. habitus dorsal; 262. habitus lateral; 263. head frontal; 264. head lateral. 265–266. *Ubristes*, male genitalia lateral. 265. *S. flavitibia* (holotype); 266. *S. jaguarinus* (holotype).

***Ubristes jaguarinus* spec. nov.**

Figs 261–264, 266.

Studied type specimens. HOLOTYPE. COSTA RICA. Male. Label 1: "COSTA RICA. Prov. Guanacaste, Z.P. / Nosara. Hojanca, R.F. Monte Alto, / Send. La Ceiba, 500-600 m., 17 AUG / 2001. I. Jimenez, Libre / L_N_221650_382750#64060"; label 2: "INB0003350605 / INBIOCRI COSTA RICA"; label 3 (red): "Ultimo especimen en / B.D. A. Lépez / 2.7.2002" / other side: "?MCR-10". Coll. INBIO.

Description (based on holotype)

Adult male. Body size: 10 mm.

Head. Face occupying slightly less than 1/2 of head width in frontal view; yellow, with narrow, brown median line on lower 1/2; yellow pilose laterally and ventrally, black pilose medially and dorsally, except narrowly bare medially; narrowly pollinose along eye margin. Gena yellow, except for narrow brown line from eye margin to oral margin. Lateral oral margins produced. Frons yellow; mixed yellow and black pilose laterally. Vertex yellowish brown, blackish at and anterior of ocellar triangle; black pilose. Occiput yellow; black pilose dorsally, yellow pilose laterally and ventrally. Eye very sparsely and short pilose, with pili about as long as ommati diameter, appearing bare under low magnification. Antennal fossa about as wide as high. Antenna yellowish; antennal ratio 6:1:7; basoflagellomere parallel-sided with narrowly rounded apex, with sensory pit located at 3/4 from base, within a vague groove that ranges from the base of the arista to close to the apex. Arista slender, about 2/3 of length of basoflagellomere.

Thorax. Scutum black, except margins widely yellow; appressed golden pilose along anterior margin, along transverse suture and on median vitta as wide as 1/3 of width of scutum; black pilose on other parts. Postpronotum yellow; yellow pilose. Postalar callus yellow; black pilose. Scutellum brownish; black pilose along margins, yellow pilose medially. Anepisternum dark brown anteriorly, yellow posteriorly; yellow pilose anteriorly and along posterior margin. Anepimeron yellow pilose on dorsal half. Katatergum and anatergum long and short microtrichose, respectively. Other pleurae bare. Calypter and halter yellow.

Wing: hyaline, veins in anterior half yellowish brown. Microtrichose, except bare on 1st costal cell, basal 1/3 of 2nd costal cell, basally on cell r1 along vein RS, on basal 3/4 of cell br, anterobasal 1/2 of cell bm, basal 1/6 of cell cup, almost entirely on alula (only a narrow basal strip with microtrichia).

Legs: yellow; yellow pilose, except hind leg mixed black and yellow pilose. Front and mid coxae and trochanters yellow; yellow pilose. Hind coxa and trochanter dark brown; yellow pilose, except coxa black pilose apically.

Abdomen. Tergite 1 brown medially, yellow laterally; yellow pilose. Tergite 2 halfway with lateral bulge-like tubercles; yellow with median brown vitta on anterior 3/4 and lateral brown vittae on anterior 3/4; long yellow pilose laterally on anterior half, short black pilose on posterior half, long black pilose on lateral tubercles. Tergite 3 narrowly yellow along anterior margin and on posterior 1/3, brown in between; black pilose. Tergite 4 brown on anterior 3/5 and on narrow median line extending almost to posterior margin, yellow on posterior 2/5; yellow pilose on much of brown parts, black pilose on yellow parts. Sternite 1 yellow; bare. Sternite 2 yellow; yellow pilose. Sternite 3 and 4 brown; yellow pilose. Genitalia as in fig. 266.

Female. Unknown.

Etymology. With a little imagination, the maculate colour pattern of *Ubristes jaguarinus* reminds of that of the jaguar, a large, feline carnivore occurring in the new world tropics.

Diagnosis. Within *Ubristes* s.s. this is the only known species with a maculate abdomen.

Distribution. Only known from Costa Rica.

REST GROUP


The species below were placed in *Microdon* subgenus *Ubristes* s.l. by Thompson *et al.* (1976), but do not fit into the concepts of the species groups treated in the present paper. They are classified into other genera following the classification of Reemer and Ståhls (2013a). Redescriptions and notes are given below.

Microdon (Chymophila) angulatus Hull, 1943

Figs 267–269.

Microdon angulatus Hull, 1943: 715.

Studied type specimens (see notes below). HOLOTYPE. Male. Label 1 (round, blue): "Ega"; label 2: "PARATYPE / *Microdon* / *angulata* / Hull"; label 3: "HOLOTYPE / *Microdon angulatus* / Hull, 1943". Coll. CNC. PARATYPE. BRAZIL. Male. Label 1: "Paratype"; label 2: "Amazon 66 53"; label 3: "Holotype *Microdon angulata* Hull". Coll. BMNH.



FIGURES 267–271. 267–269. *Microdon (Chymophila) angulatus* male (paratype). 267. habitus dorsal; 268. habitus lateral; 269. genitalia lateral. *Peradon chrysopygus* female (holotype), habitus dorsal. Photo: Luca Picciau (MRSN). 271. Number of species (y-axis) per number of known specimens (x-axis) for the genera revised in the present paper.

PARATYPE. BRAZIL. Male. Label 1: "Paratype"; label 2: "Amazon 66 53"; label 3: "Paratype *Microdon angulata* Hull". Coll. BMNH.

Notes on type specimens. The description of Hull (1943) was based on a male holotype, labelled "Ega" and two male paratypes, labelled "Amazon, 66 53". The holotype is now in the CNC collection, carrying a yellow label stating that it's a paratype. The BMNH collection holds the two paratypes, one of which is labelled as holotype. As Hull (1943) stated that the holotype is from Ega, the specimen in the BMNH collection labelled as holotype can not be regarded as such. A new red label with the text "HOLOTYPE / *Microdon angulatus* / Hull, 1943" has been added to the CNC specimen by the present author.

Redescription

Adult male. Body size: 11 mm.

Head. Face occupying 1/3 of head width in frontal view; shining dark brown; white pilose, except for a dense brush of black pile at anterior oral margin. Gena blackish brown. Oral cavity with lateral margins produced. Frons blackish, white pilose; vertex blackish, black pilose. Occiput black; black pilose dorsally, white pilose ventrally. Eye sparsely but clearly pilose. Antennal fossa about as wide as high. Antenna with scape dark brown, pedicel pale brown except for dark brown base; ratio of scape:pedicel as 4:1. [N.B.: in the specimen labelled as 'holotype' the antennae are missing, while in the other paratype the basoflagellomeres are missing. Hull (1943) describes the antennae as follows: "Antennae very slender, a little longer than the depth of the face. Third joint about as long as

the first joint and two-and-one-half times as long as the second; all of the antennae brown in colour, except blackish base of the second joint. Arista slender, not quite as long as third joint.”]

Thorax. Scutum black with dark brown lateral margins; black pilose, with pale pile along transverse suture. Postpronotum and postalar callus dark brown, black pilose. Scutellum dark brown, black pilose basally, pale pilose apically; with two apical pale brown calcars about 1/3 as long as length of the scutellum, with mutual distance about 1/2 the width of the scutellum; scutellum concave between calcars, but not sulcate. Anterior and posterior part of anepisternum divided by a clear sulcus; black pilose anteriorly and posteriorly, these patches of pilosity dorsally connected. Anepimeron black pilose. Katepisternum black pilose dorsally. Katatergum and anatergum pilose and microtrichose, respectively. Other pleurae bare. Calypter greyish yellow, halter brownish.

Wing: brown infuscated, especially in and posterior to costal cell, with pales part in apical 1/4. Entirely microtrichose.

Legs: Blackish, with tarsi brown: first tarsomeres dark brown, last two tarsomeres yellowish brown. Legs black pilose, except yellow pilose dorsally on last two tarsomeres and ventrally on all tarsomeres. Hind tibia and first tarsomeres of hind tarsus very dense and long pilose, with longest pile longer than width of tibia. Coxae and trochanters brownish, with mostly pale and some black pile.

Abdomen. Blackish brown. Tergite 1 with anterior half concave, laterally with black pile. Tergite 2 wider than thorax; with long pale pile laterally and anteriorly, short black pile posteriorly. Tergites 3 and 4 fused, with suture vaguely visible over most of width; wider than thorax; short black pilose, except longer pale pilose laterally. Abdomen laterally strongly depressed between 3rd and 4th tergite. Tergite 4 with very large lateral bulges. Sternites blackish brown and pale pilose. Genitalia as in fig. 269 (based on the paratype in the BMNH collection labelled as ‘holotype’).

Female. Unknown

Notes. In its wing venation (shape of vein M1; fig. 267) and structure of the male genitalia (fig. 269), this species clearly fits into the concept of *Chymophila* Osten Sacken, 1875, a subgenus of *Microdon* (see Reemer and Ståhls 2013a). As far as currently known, this is the only stingless bee mimicking species belonging to this subgenus.

***Ubristes chrysopygus* Giglio-Tos, 1892**

Fig. 270.

Ubristes chrysopygus Giglio-Tos, 1892: 1.

Studied type specimens. HOLOTYPE. MEXICO. Female. Label 1: "836."; label 2 (green): "Orizaba"; label 3: "Ubristes chrysopyga / Giglio-Tos". Coll. MRSN. [only photographs of the holotype were studied]

Notes. Following the classification of Reemer and Ståhls (2013a), this species belongs to the *flavofascium*-group of the genus *Peradon* Reemer. A photograph of the holotype is given in fig. 270.

Discussion

Of the 52 species treated in this paper, 22 are described as new, which raises the number of valid species of Microdontinae to 476 (Reemer and Ståhls 2013a). When comparing this ratio of described vs. undescribed species with those found in other recent revisions of Neotropical Syrphidae, it is somewhat intermediate. Rotheray *et al.* (2007) also described 22 new species for species of *Copestylum* with larvae that develop in bromeliads, but among a total number of 23. In contrast, Morales and Marinoni (2009) revise 24 species of *Palpada*, only one of which they describe as new. These varying ratios result from differences per genus in average numbers of available specimens per species: the more specimens of a genus are collected, the higher the proportion of described species. Generally, species of Microdontinae are represented in collections by far fewer specimens than, for instance, species of *Palpada*. To illustrate this: for 32 out of 52 species treated in the present paper only one or two specimens are known (fig. 271). This suggests that there may be many additional species awaiting description. Therefore, it is important that identification of newly collected specimens is verified in as many ways as possible,

not only by using the key, but also by checking thoroughly the descriptions and figures. The male genitalia differ distinctly between all species of which the males are known, so these provide a good aid in identification.




FIGURE 271. Number of species (y-axis) per number of known specimens (x-axis) for the genera revised in the present paper.

As it now appears, *Stipomorpha* is the most speciose group of Microdontiinae mimicking stingless bees, containing many highly similar (but morphologically distinct) species. It contains (by subjective judgement) very good mimics of Neotropical Meliponini, especially of the genera *Trigona* Jurine, 1807 and *Tetragona* Lepeletier & Audinet-Serville, 1828. These mimics bear close resemblance to their supposed models, not only in colouration of wings, colour patterns of head, thorax and abdomen, patterns of pilosity, but also in their flight behaviour, sticking up their abdomens and leaving their ‘corbiculate’ hind legs dangling. Several species of these bee groups seem to be specifically mimicked by certain *Stipomorpha* species. For instance, *Stipomorpha goettei* and *S. guianica* are good mimics of certain *Tetragona* species, *S. mackiei* seems to mimic *Tetragona dorsalis* (Smith, 1854), while *S. lacteipennis* possibly mimics certain *Trigona* species (pers. comm. D. Roubik upon showing him pictures of these taxa). This suggests that the mimicry is Batesian rather than Müllerian (following the definitions in e.g. Gilbert 2005); the appearance of the harmless mimics seems to match that of the noxious models near-perfectly, instead of the mimics being noxious themselves and resembling each other in a more general way. The apparent rarity of the adult mimics (most species are known from few or even single specimens) also supports a Batesian model of evolution; Müllerian mimics usually are more abundant (Gilbert 2005).

A question that arises upon considering this group of flies, is whether mimicry has stimulated speciation. Would there have been as many *Stipomorpha*-species if there had been fewer species of stingless bees? In general terms: does the number of mimicking species depend on the number of possible models? A possible mechanism for mimicry-driven speciation, a version of standard allopatric speciation, could be as follows: 1. mimic A resembles model A; 2. mimic A disperses and founds an isolated population in an area where model A does not occur, so selective advantage of mimicking model A no longer exists for this population; 3. restricted gene flow results in mimic A in the new population gradually developing a resemblance to model B; 4. if population come into contact again, members of the original population of mimic A no longer recognize members of the second population as sexual partners, so the populations are reproductively isolated, thus two species have evolved.

A case in which mimicry indeed seems to be the drive for speciation was described by Naisbit *et al.* (2003) in *Heliconius* Kluk, 1780 butterflies. Apart from butterflies, very little is known about mimetic relationships of tropical insects (Gilbert 2005). The many microdentine mimics of stingless bees offer an interesting case for further examination of 'tropical mimicry'. In order to do this, it will be necessary to link the species to supposed models, examine the question whether they are Batesian or Müllerian mimics, estimate the intra-generic phylogeny by DNA-sequencing (of both mimics and models), and analyze their biogeography. This will possibly be the subject of a future paper.

The observations on flower-visiting specimens of *Stipomorpha guianica*, and possibly also of *S. fallax*, suggest that the common idea that Microdentinae do not visit flowers (e.g. Cheng and Thompson 2008) is not true for all species. Another Neotropical species, *Microdon tigrinus* Curran, 1940, has also been reported to visit flowers (Morales and Köhler 2006). Obviously, the subject of feeding by adult Microdentinae needs further attention. Perhaps investigation of gut contents could provide a first clue as to whether they feed on pollen or not.

ACKNOWLEDGEMENTS

I am grateful to Kees van Achterberg (RMNH) for allowing me to use his photographic and microscopic equipment and for his kind assistance while using it. The following persons are thanked for providing me with material or for helping me with examining specimens from the collections they are responsible for: Ben Brugge (ZMAN), Rune Bygebjerg (MZLU), David Grimaldi (AMNH), Z.H. Falin (SEMC), Martin Hauser, E. Richard Hoebeke (CU), Uwe Kallweit (SNSD), Stephen Marshall, Ximo Mengual, Tam Nguyen (AMNH), Philip Perkins (MCZ), Luca Picciau (MRSN), Peter Sehnal (NMW), Jeff Skevington (CNC), John T. Smit, Villu Soon (Museum of Zoology, Tartu), Wouter van Steenis, F. Christian Thompson (USNM), Rob de Vries (RMNH), Nigel Wyatt (BMNH), Manuel Zumbado (INBIO). Thanks are also due to Koos Biesmeijer and Francis Gilbert, who commented on an earlier version of the manuscript. Chris Thompson had recognized earlier that some of the species described in this paper were new, so I thank him for his generosity in sharing these specimens with me. He also commented on the manuscript in an early stage. Finally, a word of gratitude to Gunilla Ståhls for all her help and critical comments.

REFERENCES

- Albrecht, C.W. (1993) Arsenic and old collections. Local History Notebook March/April 1993, OHIO association of Historical Societies and Museums, Columbus. Available from: <http://www.ohiohistory.org/local-history-office/ohio-local-history-alliance> (Accessed 25 June 2013)
- Cheng, X.-Y. & Thompson, F.C. (2008) A generic conspectus of the Microdentinae (Diptera: Syrphidae) with the description of two new genera from Africa and China. *Zootaxa*, 1879, 21–48.
- Costa, M.A., del Lama, M.A., Melo, G.A.R. & Sheppard, W.S. (2003) Molecular phylogeny of the stingless bees (Apidae: Apinae, Meliponini) inferred from mitochondrial 16S rDNA sequences. *Apidologie*, 34, 73–84.
- Curran, C.H. (1925) New American Diptera. II. *Annals and Magazine of Natural History*, 9th series, 16, 338–354. <http://dx.doi.org/10.1080/00222932508633318>
- Curran, C.H. (1930) New Syrphidae from Central America and the West Indies. *American Museum Novitates*, 416, 1–11.
- Curran, C.H. (1934) Diptera of Kartabo, Bartica District, British Guiana. *Bulletin of the American Museum of Natural History*, 66, 287–532.
- Curran, C.H. (1940) Some new Neotropical Syrphidae (Diptera). *American Museum Novitates*, 1086, 1–14.
- Curran, C.H. (1941) New American Syrphidae. *Bulletin of the American Museum of Natural History*, 78, 243–304.
- Doesburg Sr.P.H. van (1966) Syrphidae from Suriname. Additional records and descriptions. *Studies on the Fauna of Suriname and other Guyanas*, 35, 61–107.
- Gilbert, F.S. (2005) The evolution of imperfect mimicry in hoverflies. In: Fellowes, M., Holloway, G. & Jens, R. (Eds.), *Insect Evolutionary Ecology*. Oxford University Press, Oxford, 448 pp.
- Hippa, H. & Ståhls, G. (2005) Morphological characters of adult Syrphidae: descriptions and phylogenetic utility. *Acta Zoologica Fennica*, 215, 1–72.
- Hull, F.M. (1937) New species of exotic syrphid flies. *Psyche*, 44, 12–32. <http://dx.doi.org/10.1155/1937/46960>
- Hull, F.M. (1943) Some flies of the genus *Microdon* in the British Museum (Natural History). *Annals and Magazine of Natural History*, 10, 702–720. <http://dx.doi.org/10.1080/03745481.1943.9728054>

- Hull, F.M. (1945) Some undescribed syrphid flies. *Proceedings of the New England Zoological Club*, 23, 71–78.
- Hull, F.M. (1949) The morphology and inter-relationship of the genera of syrphid flies, recent and fossil. *Transactions of the Zoological Society of London*, 26, 257–408.
<http://dx.doi.org/10.1111/j.1096-3642.1949.tb00224.x>
- Mann, W.M. (1928) A new *Microdon* from Panama. *Psyche*, 35, 168–170.
<http://dx.doi.org/10.1155/1928/73806>
- Marshall, S.A. (2012) *Flies: The natural history and diversity of Diptera*. Firefly Books, Ontario, 616 pp.
- McAlpine, J.F. (1981) Morphology and terminology - Adults. *Manual of Nearctic Diptera*, 1, 9–63.
- Morales, M. & Köhler, A. (2006) Espécies de Syrphidae (Diptera) visitantes das flores de *Eryngium horridum* (Apiaceae) no Vale do Rio Pardo, RS, Brasil. *Iheringia, Sér. Zool.*, 96, 41–45.
<http://dx.doi.org/10.1590/s0073-47212006000100006>
- Morales, M. & Marinoni, L. (2009) Cladistic analysis and taxonomic revision of the *scutellaris* group of *Palpada* Macquart (Diptera : Syrphidae). *Invertebrate Systematics*, 23, 301–347.
<http://dx.doi.org/10.1071/is09006>
- Naisbit, R.E., Jiggins, C.D. & Mallet, J. (2003) Mimicry: developmental genes that contribute to speciation. *Evolution and development*, 5, 269–280.
<http://dx.doi.org/10.1046/j.1525-142x.2003.03034.x>
- Reemer, M. & Ståhls, G. (2013a) Generic revision and species classification of the Microdontinae (Diptera: Syrphidae). *ZooKeys*, 288, 1–213.
<http://dx.doi.org/10.3897/zookeys.288.4095>
- Reemer, M. & Ståhls, G. (2013b) Phylogenetic relationships of Microdontinae (Diptera: Syrphidae) based on molecular and morphological characters. *Systematic Entomology*. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/syen.12020/pdf> (Accessed 1 Aug. 2013)
- Rotheray, G.E., Hancock, E.G. & Marcos García, M.A. (2007) Neotropical *Copestylum* (Diptera, Syrphidae) breeding in bromeliads (Bromeliaceae) including 22 new species. *Zoological Journal of the Linnean Society*, 150, 167–317.
<http://dx.doi.org/10.1111/j.1096-3642.2007.00288.x>
- Roubik, D.W., Smith, B.H. & Carlson, R.G. (1987) Formic acid in caustic cephalic secretions of stingless bee, *Oxytrigona* (Hymenoptera: Apidae). *Journal of Chemical Ecology*, 13, 1079–1086.
<http://dx.doi.org/10.1007/bf01020539>
- Sack, P. (1941) Syrphidae (Dipt.). In: Titschack, E. (Ed.), *Beiträge zur Fauna Perus. Band 1B. Nach der Ausbeute der Hamburger Südperu-Expedition 1936, anderer Sammlungen, wie auch auf Grund von Literaturangaben*. Verlag Conrad Behre, Hamburg, 403 pp.
- Schummel, T.E. (1842) Verzeichniss und Beschreibung der vom Verfasser bis jetzt in der Schleisen gefangenen Zweiflügler der Syrphen Familie. *Uebersicht der schlesische Gesellschaft für Vaterländische Kultur*, 1841, 112–126, 163–170.
- Shannon, R.C. (1925) An extraordinary adult myrmecophile from Panama. *Proceedings of the National Academy of Sciences*, 10, 211–213.
- Shannon, R.C. (1927) A review of the South American two-winged flies of the family Syrphidae. *Proceedings of the United States National Museum*, 70, 1–33.
<http://dx.doi.org/10.5479/si.00963801.70-2658.1>
- Thompson, F.C. (1999) A key to the genera of the flower flies (Diptera: Syrphidae) of the Neotropical region including descriptions of new genera and species and a glossary of taxonomic terms. *Contributions on Entomology, International*, 3, 321–378.
- Thompson, F.C., Vockeroth, J.R. & Sedman, Y.S. (1976) Family Syrphidae. *A catalogue of the Diptera of the Americas south of the United States*, 46, 1–195.
- Wiedemann, C.R.W. (1830) *Aussereuropäische zweiflügelige Insekten. Zweiter Theil*. Schulz, Hamm, xii + 684 pp, 5 pls.