

## Phylum Onychophora Grube, 1853\*

GEORG MAYER<sup>1</sup> & IVO DE SENA OLIVEIRA<sup>2</sup>

<sup>1</sup> Institute of Biology, Animal Evolution & Development, University of Leipzig, Talstraße 33, D-04103 Leipzig, Germany.  
E-mail: gmayer@onychophora.com

<sup>2</sup> Institute of Biology, Animal Evolution & Development, University of Leipzig, Talstraße 33, D-04103 Leipzig, Germany.  
E-mail: ivo.de\_sena\_oliveira@uni-leipzig.de

\* In: Zhang, Z.-Q. (Ed.) Animal Biodiversity: An Outline of Higher-level Classification and Survey of Taxonomic Richness (Addenda 2013). *Zootaxa*, 3703, 1–82.

Extant representatives of Onychophora (velvet worms) have been classified into two major subgroups or families, the Peripatidae and the Peripatopsidae. According to our previous estimate, a total of 179 nominal species and twelve additional infraspecific taxa of Onychophora had been described, including numerous “subspecies”, “variations” and “mutations (Mayer & Oliveira 2011). However, the validity of most of these taxa was uncertain. Since this last estimate, a comprehensive world checklist of Onychophora has been published, along with a revision of the nomenclatural status of names (Oliveira *et al.* 2012a). This study revealed that many previous synonyms and infraspecific taxa might constitute valid species. At the same time, it also has shown that ~10% of the described species of Onychophora show major taxonomical problems and therefore should be regarded as *nomina dubia*, following the provisions of the International Code of Zoological Nomenclature (ICZN 1999).

In addition to extant species of Onychophora, the world checklist includes three unambiguous fossil representatives, the relationship of which to the extant species is uncertain (Grimaldi *et al.* 2002; Oliveira *et al.* 2012a; Poinar 2000). A recent study further revealed that †*Helenodora inopinata* Thompson & Jones, 1980 (see Hou & Bergström 1995 for nomenclatural discussion) from Mazon Creek (Illinois, USA, ~296 million years ago) might be a representative of Onychophora as well, which is indicated by the presence of slime papillae in this species (Haug *et al.* 2012). As with most other fossils, the affinity of †*Helenodora inopinata* to the extant onychophoran taxa remains obscure.

In addition to these findings, six species and a new genus of extant Onychophora have been described in the last two years (Oliveira *et al.* 2012b, 2013; Ruhberg & Daniels 2013). This ongoing progress has necessitated an update of the estimated numbers of described species of Onychophora, which is provided below. In this updated list, we discriminate between valid taxa, *nomina dubia* and fossil taxa (indicated by a †).

Phylum **Onychophora** Grube, 1853 [5 families (3†), 54 genera (4†), 187 valid species (4†), 20 *nomina dubia*]

Family **Peripatidae** Evans, 1901 [12 genera (1†), 76 valid species (1†), 9 *nomina dubia*]

Family **Peripatopsidae** Bouvier, 1905 [39 genera, 108 valid species, 11 *nomina dubia*]

Family †**Succinipatopsidae** Poinar, 2000 [1† genus, 1† valid species]

Family †**Tertiapatidae** Poinar, 2000 [1† genus, 1† valid species]

Family †**Helenodoridae** Poinar, 2000 [1† genus, 1† valid species]

## References

- Bouvier, E.L. (1905) Monographie des Onychophores. *Annales des Sciences Naturelles Zoologie et Biologie Animale [9e Série]*, 2, 1–383.
- Evans, R. (1901) On two new species of Onychophora from the Siamese Malay States. *Quarterly Journal of Microscopical Science*, 44, 473–538.
- Grimaldi, D.A., Engel, M.S. & Nascimbene, P.C. (2002) Fossiliferous cretaceous amber from Myanmar (Burma): its rediscovery, biotic diversity, and paleontological significance. *American Museum Novitates*, 3361, 3–72.  
[http://dx.doi.org/10.1206/0003-0082\(2002\)361<0001:fcafmb>2.0.co;2](http://dx.doi.org/10.1206/0003-0082(2002)361<0001:fcafmb>2.0.co;2)
- Grube, E. (1853) Über den Bau von *Peripatus edwardsii*. *Müller's Archives of Anatomy and Physiology*, [1853], 322–360.
- Haug, J.T., Mayer, G., Haug, C. & Briggs, E.G. (2012) A Carboniferous non-onychophoran lobopodian reveals long-term survival of a Cambrian morphotype. *Current Biology*, 22, 1673–1675.  
<http://dx.doi.org/10.1016/j.cub.2012.06.066>
- Hou, X.G. & Bergström, J. (1995) Cambrian lobopodians - ancestors of extant onychophorans? *Zoological Journal of the Linnean Society*, 114, 3–19.  
<http://dx.doi.org/10.1111/j.1096-3642.1995.tb00110.x>
- International Commission on Zoological Nomenclature (1999) *International Code of Zoological Nomenclature, the fourth edition*. The International Commission for Zoological Nomenclature, London, 336 pp.
- Mayer, G. & Oliveira, I.S. (2011) Phylum Onychophora Grube, 1853. In: Zhang, Z.-Q. (Ed.) *Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness*. Zootaxa, 3148, 98.
- Oliveira, I.S., Read, V.M.S.J. & Mayer, G. (2012a) A world checklist of Onychophora (velvet worms), with notes on nomenclature and status of names. *ZooKeys*, 211, 1–70.  
<http://dx.doi.org/10.3897/zookeys.211.3463>
- Oliveira, I.S., Franke, F.A., Hering, L., Schaffer, S., Rowell, D.M., Weck-Heimann, A., Monge-Nájera, J., Morera-Brenes, B. & Mayer, G. (2012b) Unexplored character diversity in Onychophora (velvet worms): a comparative study of three peripatid species. *PLoS ONE*, 7, 1–20.  
<http://dx.doi.org/10.1371/journal.pone.0051220>
- Oliveira, I.S., Schaffer, S., Kvartalnov, P.V., Galoyan, E.A., Palko, I.V., Weck-Heimann, A., Geissler, P., Ruhberg, H. & Mayer, G. (2013) A new species of *Eoperipatus* (Onychophora) from Vietnam reveals novel morphological characters for the South-East Asian Peripatidae. *Zoologischer Anzeiger*, 252, 495–510.  
<http://dx.doi.org/10.1016/j.jcz.2013.01.001>
- Poinar, G. (2000) Fossil onychophorans from Dominican and Baltic amber: *Tertiapatus dominicanus* n.g., n.sp. (Tertiapatidae n.fam.) and *Succinipatopsis balticus* n.g., n.sp. (Succinipatopsidae n.fam.) with a proposed classification of the subphylum Onychophora. *Invertebrate Biology*, 119, 104–109.  
<http://dx.doi.org/10.1111/j.1744-7410.2000.tb00178.x>
- Ruhberg, H. & Daniels, S. (2013) Morphological assessment supports the recognition of four novel species in the widely distributed velvet worm *Peripatopsis moseleyi sensu lato* (Onychophora: Peripatopsidae). *Invertebrate Systematics*, 27, 131–145.  
<http://dx.doi.org/10.1071/is12069>
- Thompson, I. & Jones, D.S. (1980) A possible onychophoran from the Middle Pennsylvanian Mazon Creek Beds of northern Illinois. *Journal of Paleontology*, 54, 588–596.