

<http://dx.doi.org/10.11646/zootaxa.4039.4.10>
<http://zoobank.org/urn:lsid:zoobank.org:pub:8348B8C1-EFF2-4D4F-9C21-F8C60C2F3FE6>

MARIANNA ISABELLA ROSA RODRIGUES DE OLIVEIR, LUIZ NORBERTO WEBER & MARCELO FELGUEIRAS NAPOLI (2013) Internal oral morphology in larvae of the genus *Rhinella* Fitzinger, 1826 (Amphibia, Anura, Bufonidae). *Zootaxa*, 3745 (5): 501–523.

On page 519 of this publication the first part of the Table 2 was erroneously replaced by a copy of the second part of the Table 2. The correct first part of the Table 2 is presented here:

	<i>R. abei</i>	<i>R. arenarum</i>	<i>R. crucifer</i>	<i>R. granulosa</i>	<i>R. henseli</i>	<i>R. hoogmoedi</i>
Prenarial arena	Sinuous ridge Irregular apex Laterally there are three pustules	Transversal ridge Irregular apex	Sinuous ridge Irregular apex	Sinuous ridge Irregular apex	Transversal ridge Irregular apex	Three central projections higher and a lower lateral projection
Prenarial papillae	3	2	2	2	3	6
Narial valve projections	-	-	+	+	-	+
Postnarial papillae number	6	3	4	4	4	Ridge apparently formed by two or three papillae joined by the base
Postnarial papillae shape	Two anterior are conical Four posterior are digitiform	Conical	Four conical papillae on each side Two small conical papillae laterally	Digitiform	Conical	Ridge
Postnarial papillae apex	Most anterior irregular apex Most posterior tapered apex	Tapered	Bifid	Irregular	Rounded	Irregular
Postnarial papillae arrangement	Four posterior are aligned transversely	Inverted V	Inverted V	Aligned on the side of the area	Inverted V	Inverted V
Postnarial pustules	0	6	0	4	0	0
Median ridge	Trapezoid or triangular-shaped Irregular margin	Trapezoid-shaped Irregular margin	Triangular-shaped Irregular margin	Triangular-shaped Irregular margin	Trapezoid-shaped Irregular margin	Trapezoid-shaped Irregular margin
Lateral ridge papillae	Apex as wide as the base Three digitiform secondary branches	Apex as wide as the base Three digitiform secondary branches	Apex wider than the base Four digitiform secondary branches	Apex wider than the base Three digitiform secondary branches	Apex as wide as the base Three digitiform secondary branches	Apex wider than the base Three to four digitiform secondary branches

.....continued on the next page

TABLE 2. (Continued)

	<i>R. abei</i>	<i>R. arenarium</i>	<i>R. crucifer</i>	<i>R. granulosa</i>	<i>R. henseli</i>	<i>R. hoogmoedi</i>
Number of BFA papillae	Nearly 4-6	Nearly 5	Nearly 3-4	Nearly 4-5	Nearly 3-4	Nearly 3
BRA papillae shape	Digitiform elongated	Digitiform elongated	Digitiform elongated	Digitiform elongated	Digitiform short	Digitiform short
Pustules of the BRA	Nearly 10	Nearly 75	Nearly 50	Nearly 55	Nearly 55	Nearly 20
BRA papillae arrangement	Arranged on the side	Lower side	"V" design	Aligned on the side	"V" design	Lower side
Velar margin	Undulated Lateral fold with glandular aspect Without projections	Undulated provided with lateral fold with glandular appearance Without projections	Undulated Lateral fold with glandular appearance Without projections	Undulated Lateral fold with glandular aspect Elongated	Undulated Lateral fold with glandular aspect projections in the median region	Undulated Lateral fold with glandular aspect Without projections