



Two new species and new records of chiggers (Acari: Ixodidae, Trombiculidae) from birds in Vietnam

STANISLAV KALUZ¹, NGUYEN MANH HUNG², MIROSLAV CAPEK³ & IVAN LITERAK^{4,5}

¹Institute of Zoology, Slovak Academy of Sciences, Dubravská cesta 9, 845 06 Bratislava, Slovakia; e-mail: stanislav.kaluz@savba.sk

²Department of Parasitology, Institute of Ecology and Biological Resources (IEBR), Vietnamese Academy of Science and Technology (VAST), 18-Hoang Quoc Viet, Hanoi, Vietnam; e-mail: hung_iebr@yahoo.com

³Institute of Vertebrate Biology, Academy of Sciences of the Czech Republic, v. v. i., Kvetná 8, 603 65 Brno, Czech Republic; e-mail: capek@ivb.cz

⁴Department of Biology and Wildlife Diseases, Faculty of Veterinary Hygiene and Ecology, University of Veterinary and Pharmaceutical Sciences, Palackého tr. 1, 612 42 Brno, Czech Republic; e-mail: literaki@vfu.cz

⁵Corresponding author. E-mail: literaki@vfu.cz

Abstract

A total of 12 chigger species (Acariformes: Trombiculidae) occurred on 7 bird species in Vietnam. Two new species, namely *Neoschoengastia vietnamensis* sp. nov. and *Hypogastia stekolnikovi* sp. nov. are described, figured and compared with similar species placed in relevant genera using differential diagnoses of related species. Figures and diagnosis of *Leptotrombidium taiyuanense* Tian and Wen, 1984 are added. *Odontacarus audy* (Radford, 1946), *Leptotrombidium allossetum* Wang, Liao and Lin, 1981, *L. taiyuanense*, *Leptotrombidium hansenii* Traub and Lakshana, 1966, *Leptotrombidium kunshui* Wen and Xiang, 1984, *Leptotrombidium paradoxum* Vercammen-Grandjean and Langston, 1976, *Leptotrombidium turdicola* Vercammen-Grandjean and Langston, 1976, *Neotrombicula elegans* Schluger, 1966 and *Neoschoengastia longitarsalis* Schluger and Belskaya, 1966 were recorded in Vietnam for the first time.

Key words: birds, chiggers, taxonomy, *Leptotrombidium*, *Neoschoengastia*, *Hypogastia*, new species

Introduction

Chiggers or larvae of Trombiculoidea (Acariformes) parasitize mostly terrestrial vertebrates. The host-parasite relationships include combinations of diverse hosts with various chiggers (Shatrov & Kudryashova 2008). Birds constitute the preferred group of hosts for chiggers belonging for example to the genera *Neoschoengastia* Ewing, 1929, and *Hypogastia* Vercammen-Grandjean, 1967. These genera parasitize mainly water birds or birds nesting in various ground or tree holes (Nadchatram & Upham 1966). Small forest or mountain birds, too, are frequently parasitized (Brennan 1948, 1951, 1962, 1965). A good example of various host distribution is seen in the largest genus, *Leptotrombidium* Nagayo, Miyagawa, Mitamura and Imamura, 1916, which includes about 340 species (Stekolnikov 2013). Many of these are associated with rodents, whereas nearly 80 species have been found on other animals, including birds (Kudryashova 1998; Shatrov & Kudryashova 2008). The genus *Leptotrombidium* has been studied for decades and many papers have appeared in various parts of the world, including extensive publications elaborating the entire genus (Vercammen-Grandjean 1968; Vercammen-Grandjean & Langston 1976; Kudryashova 1998; Stekolnikov 2013).

Chiggers occurring on birds in South Asia and South-East Asia have been studied by several authors (e.g. Vercammen-Grandjean *et al.* 1970; Goff 1983; Stan Fernandes & Kulkarni 2003). Wharton & Hardcastle (1946) and Brennan & Amerson (1971) dealt with chiggers on birds in the Pacific area, while Nadchatram & Upham (1966) studied chiggers associated with birds in Malaysia. Few papers were aimed on chiggers in Vietnam (Schluger *et al.* 1960a, 1960b, 1960c, 1961, 1963).

This paper describes two new chigger species of the genera *Neoschoengastia* and *Hypogastia* collected from

small birds in a tropical forest in Vietnam. Also included here are figures and characters of *Leptotrombidium taiyuanense*, which was originally described insufficiently, as well as a list of new records of known chigger taxa documented in this study.

Material and methods

Wild birds were captured and examined at two locations within the Cuc Phuong National Park in northern Vietnam in 2010. The first location was situated in the tropical forest around the tourist centre and ranger station in the centre of the National Park, at 350 m above sea level (20°21' N 105°35' E, 1-5 February 2010)—location A; the second was in the botanical garden on the southeastern part of the National Park, at 140 m above sea level (20°15' N 105°42' E, 6-8 February 2010)—location B. Birds were mist-netted, identified based on Robson (2005), then visually checked for the presence of chiggers. Bird taxonomy follows Gill & Donsker (2015). Mammal taxonomy follows Mammal Species of the World (2015). Once examined, the birds were released back into the wild.

Chiggers were collected by Ivan Literak with tweezers and stored in 70% ethanol. We examined 161 birds belonged to 36 species. Chiggers were found on 48 individual birds belonging to seven species of passerines: *Copsychus malabaricus* (Scopoli), *Hemixos flava* Blyth, *Larvivora sibilans* (Swinhoe), *Niltava davidi* La Touche, *Pellorneum ruficeps* Swainson, *Turdus cardis* Temminck, and *Turdus dissimilis* Blyth. Chigger larvae were mounted onto slides using Swann's medium, microscopically identified, and the new species were described. All drawings were produced using standard light microscopy and enhanced with computer software (GIMP 2.8). The measurements in micrometres (µm) used in the text and figures were taken from the slide-mounted specimens using a stage-calibrated ocular micrometre. The measurements, morphological features of specimens, and anatomical nomenclature in this paper follow the classification by Vercammen-Grandjean (1968), Vercammen-Grandjean & Langston (1976), Kudryashova (1998) and Stekolnikov (2013).

The descriptions of the new species are attributed to the first author of this paper. The types of the new species described in this paper are deposited in the Department of Zoology Museum of the Institute for Ecology and Biological Resources, Vietnamese Academy of Science and Technology, Hanoi, Vietnam (IEBR VAST); in the Slovak National Museum, Bratislava, Slovakia (SNM); and in the Natural History Museum, London, United Kingdom (BMNH).

Systematics

A total of 12 chigger species occurred on seven species of passerines (Table 1).

Family Leeuwenhoekidae Womersley

Genus *Odontacarus* Ewing, 1929

Odontacarus audyi (Radford, 1946)

Material examined. Location A: 1 larva from *Turdus cardis* (Passeriformes: Turdidae), 1 February 2010; 3 larvae from *Niltava davidi* (Passeriformes: Muscicapidae), 4 February 2010; 6 larvae from *Niltava davidi*, 5 February 2010; 5 larvae from *Larvivora sibilans* (Passeriformes: Muscicapidae), 5 February 2010. Location B: 12 larvae from *Larvivora sibilans*, 7 February 2010; 8 larvae from *Pellorneum ruficeps* (Passeriformes: Pellorneidae), 7 February 2010; 1 larva from *Hemixos flava* (Passeriformes: Pycnonotidae), 7 February 2010; 14 larvae from *Copsychus malabaricus* (Passeriformes: Muscicapidae), 8 February 2010.

Distribution and hosts. *O. audyi* is widely distributed in South-East Asia (India, Malaysia, Thailand) and infests various birds (Nadchatram 1963). This author found *O. audyi* on *Clamator coromandus* (Linnaeus) (Cuculiformes: Cuculidae), *Centropus sinensis* (Stephens) (Cuculiformes: Cuculidae), *Pitta brachyura* (Linnaeus) (Passeriformes: Pittidae), *Pellorneum ruficeps* (Passeriformes: Pellorneidae), *Luscinia cyane* (Pallas)

(Passeriformes: Muscicapidae), *Saxicola ferreus* Gray (Passeriformes: Muscicapidae), *Garrulax moniliger* (Hodgson) (Passeriformes: Leiothrichidae), *Copsychus malabricus* (Passeriformes: Muscicapidae), *Anthus hodgsoni* Richmond (Passeriformes: Motacillidae), *Lanius collurio* Lesson (Passeriformes: Laniidae), and *Cyornis banyumas* (Horsfield) (Passeriformes: Muscicapidae). *O. audyi* is recorded in Vietnam for the first time. *Turdus cardis*, *Niltava davidi* and *Larvivora sibilans* are new hosts for this species.

Trombiculidae Ewing

Tribe Trombiculini Vercammen-Grandjean

Genus *Leptotrombidium* Nagayo, Miyagawa, Mitamura and Imamura, 1916

Leptotrombidium alloseium Wang, Liao and Lin, 1981

Material examined. Location A: 7 larvae from *Pellorneum ruficeps* (Passeriformes: Pellorneidae), 1 February 2010; 5 larvae from *Pellorneum ruficeps*, 4 February 2010; 12 larvae from *Turdus dissimilis* (Passeriformes: Turdidae), 1 February 2010; 1 larva from *Larvivora sibilans* (Passeriformes: Muscicapidae), 3 February 2010.

Distribution and hosts. *L. alloseium* is reported from *Apodemus agrarius* (Pallas) (Rodentia: Muridae) in China (Wang *et al.* 1981). We found *L. alloseium* in Vietnam for the first time. *Pellorneum ruficeps* and *Larvivora sibilans* are new hosts for this species.

Leptotrombidium taiyuanense Tian and Wen, 1984

(Figs. 1–5)

Material examined. Location A: Larva from *Turdus cardis* (Passeriformes: Turdidae); 4 February 2010.

Description. Larva, 1 individual measured. SIF = 7B-B.3.2111.0000; fsp = 7-7-7; fPp = N.N.BNN; Ga = B; Pc = 3; Gn = 2; fD = 2H.12.12.12.7.3.3 = 51; fSc: AM = AL = PL; SB/PL; fCx = 1.1.1; fSt = 2.2; DS = 51; VS = 44; NDV = 95.

Idiosoma (Figs. 1, 2). Scutum roughly rectangular, posterior margin weakly sinuous with very shallow concave median depression, two times wider than long, with AL, AM, PL setae and two flagelliform sensilla (Fig. 3). Position of AL and PL setae laterally marginal. Scutal puncta spread and cover most of scutum. Sensilla absent. Rounded eyes (2 + 2), anterior larger than posterior, ocular plates not visible. Humeral setae 46 long, dorsal anterior 36–40 long, central 34–38 and posterior setae 36–39 long, densely ciliated, dorsal setae arranged in regular rows, the rows slightly varying in position of setae. Scutal measurements: AW = 63, PW = 71, SB = 29, ASB = 28, PSB = 15, SD = 43, AP = 29, AM = 45, AL = 45, PL = 45, PL = AM = AL, SD < AW < PW, SB situated slightly anteriorly from level of PL. Ventral side with two pairs of sternal setae (fSt = 2.2), fCx = 1.1.1. Ventral side with 44 irregularly arranged ciliated setae. Lengths of ventral setae vary, increase from anterior 21–24 to median 23–26 and posterior 34–39.

Gnathosoma (Fig. 4). Palps 42 (+ claw) long, palpotibial claw unmeasurable, claw with 3 deeply indented prongs, galeala nude. Chelicera anteriorly with tricuspid cap.

Legs (Fig. 5). With pretarsus, claws and normal empodia. All tactile setae on legs barbed, slender. Specialized setae on leg segments—Leg I: Tarsus—S₁ 17, f₁ 2, nude pretarsala 11 (PT' = N), nude ST 21 and pST 11, 2 tibialae (15 and 15), microtibialae 3, 2 genualae—anterior 19, posterior 15, microgenuala 3; Leg II: Tarsus—nude pretarsala 11 (PT'' = N), S₂ 17, f₂ 2, 2 tibialae—anterior 13, posterior 12, genuala 14; Leg III: tibialae 14 and genuala 15. Number of barbed setae on leg segments (leg formula from coxa to tarsus): Leg I: 1-1-1-5-4-8-18; Leg II: 1-1-2-4-3-6-16; Leg III: 1-1-2-3-3-6-14. Length/width of tarsi I–III: I—57/20, II—49/17, III—69/14. Leg length: pa 251, pm 228, pp 258, lp 737.

Distribution and hosts. The species was originally described from a single specimen from *Tscherskia triton* (Winton) (Rodentia: Cricetidae) in China (Tian and Wen, 1984). We found *L. taiyuanense* in Vietnam for the first time. *Turdus cardis* is a new host for this chigger.

TABLE 1. Occurrence of chigger species on birds collected in Cuc Phuong National Park. Lieuwenhoeikiidae: OA – *Odontacarus audyi*; Trombiculidae, Trombiculini: LA – *Leptotrombidium alloseosum*, LTA – *Leptotrombidium taiyuanense*, LH – *Leptotrombidium hansenii*, LK – *Leptotrombidium kunshui*, LP – *Leptotrombidium paradox*, LT – *Leptotrombidium turdicola*, NE – *Neotrombicula elegans*; Trombiculidae, Schoengastini: NV – *Neoschoengastia vietnamensis* **sp. n.**, NL – *Neoschoengastia longitarsalis*, HSC – *Helenicula scanloni*, HST – *Hypogastia stekolnikovi* **sp. n.**

Number of chiggers (Number of birds parasitized)													
Birds	Number of birds examined	OA	LA	LTA	LH	LK	LP	LT	NE	NV	NL	HSC	HST
<i>Copsychus malabaricus</i>	2	14 (1)									1 (1)		
<i>Hemixos flavala</i>	1	1 (1)											
<i>Larvivora sibilans</i>	11	17 (3)	1 (1)			1 (1)			1 (1)	55 (4)			1 (1)
<i>Niltava davidi</i>	7	9 (4)								3 (2)			7 (1)
<i>Pellorneum ruficeps</i>	18	8 (1)	12 (4)		8 (3)	2 (2)	1 (1)	1 (1)	25 (3)	1 (1)		2 (2)	
<i>Turdus cardis</i>	5	1 (1)		1 (1)						78 (3)			
<i>Turdus dissimilis</i>	2		12 (1)						2 (1)				
Total	46	50 (11)	25 (6)	1 (1)	8 (3)	3 (3)	1 (1)	1 (1)	28 (5)	137 (10)	1 (1)	2 (2)	8 (2)




FIGURE 1. *Leptotrombidium taiyuanense* (larva)—idiosoma dorsal. Scale bar: 100 μ m.




FIGURE 2. *Leptotrombidium taiyuanense* (larva)—idiosoma ventral. Scale bar: 100 μ m.

***Leptotrombidium hansenii* Traub and Lakshana, 1966**

Material examined. Location A: 7 larvae from *Pellorneum ruficeps* (Passeriformes: Pellorneidae), 1 February 2010; 1 larva from *Pellorneum ruficeps*, 4 February 2010.

Distribution and hosts. *L. hansenii* is reported to parasitize *Tupaia glis* (Diard) (Scandentia: Tupaiidae) in China (Stekolnikov 2013). We found *L. hansenii* in Vietnam for the first time. *Pellorneum ruficeps* is a new host for this chigger.




FIGURE 3. *Leptotrombidium taiyuanense* (larva)—scutum. Scale bar: 100 μ m.




FIGURE 4. *Leptotrombidium taiyuanense* (larva)—gnathosoma (left—ventral, right—dorsal). Scale bar: 100 μ m.




FIGURE 5. *Leptotrombidium taiyuanense* (larva)—legs I–III (dots—bases of setae; filled dots—upper side; empty dots—bottom side. Numbers—lengths of solenidia and famuli. Scale bar: 100 μ m.

***Leptotrombidium kunshui* Wen and Xiang, 1984**

Material examined. Location A: 1 larva from *Pellorneum ruficeps* (Passeriformes: Pellorneidae), 1 February 2010; 1 larva from *Pellorneum ruficeps*, 4 February 2010; 1 larva from *Larvivora sibilans* (Passeriformes: Muscicapidae), 4 February 2010.

Distribution and hosts. *L. kunshui* is reported from *Rattus tanezumi* Temminck (Rodentia: Muridae) in China (Wen & Xiang 1984). We found *L. kunshui* in Vietnam for the first time. Both *P. ruficeps* and *L. sibilans* are new hosts for this chigger.

***Leptotrombidium paradox* Vercammen-Grandjean and Langston, 1976**

Material examined. Location A: 1 larva from *Pellorneum ruficeps* (Passeriformes: Pellorneidae), 4 February 2010.

Distribution and hosts. *L. paradox* is reported to parasitize small rodents of the family Muridae (Rodentia) in India (Stan Fernandes & Kulkarni 2003) and in Russian Caucasus (Krasnodar Krai, Adygea, Karachai-Cherkessia, North Ossetia, Dagestan) (Stekolnikov 2004, 2013). We found *L. paradox* in Vietnam for the first time. *Pellorneum ruficeps* is a new host for this chigger.

***Leptotrombidium turdicola* Vercammen-Grandjean and Langston, 1976**

Material examined. Location A: 1 larva from *Pellorneum ruficeps* (Passeriformes: Pellorneidae), 4 February 2010.

Distribution and hosts. *Leptotrombidium turdicola* is known from *Turdus obscurus* Gmelin (Passeriformes: Turdidae) in Malaysia (Vercammen-Grandjean & Langston 1976) and from *Apodemus agrarius* (Rodentia: Muridae) in China (Stekolnikov 2013). We found *L. turdicola* in Vietnam for the first time. *Pellorneum ruficeps* is a new host for this chigger.

Genus *Neotrombicula* Hirst, 1925

***Neotrombicula elegans* Schluger, 1966**

Material examined. Location A: 24 larvae from *Pellorneum ruficeps* (Passeriformes: Pellorneidae), 1 February 2010; 1 larva from *Larvivora sibilans* (Passeriformes: Muscicapidae), 4 February 2010; larva from *Pellorneum ruficeps*; 4 February 2010; 2 larvae from *Turdus dissimilis* (Passeriformes: Turdidae), 4 February 2010.

Distribution and hosts. *N. elegans* infests small rodents *Apodemus agrarius*, *Apodemus flavicollis* (Melchior), *Apodemus sylvaticus* (Linnaeus) (Rodentia: Muridae), *Myodes glareolus* Schreber, and *Microtus arvalis* (Pallas) (Rodentia: Cricetidae) in Ukraine (Kudryashova 1998). Larvae found in Slovakia were isolated from soil samples (Kalúz & Vrabec 2014). We found *N. elegans* in Vietnam for the first time. *Pellorneum ruficeps*, *Larvivora sibilans* and *Turdus dissimilis* are new hosts for this chigger.

Remarks. Chiggers of this species from Vietnam are identical with *N. elegans* from Ukraine and Slovakia. The measurements of individuals of *N. elegans* are presented here: Vietnam (this paper): AW 66–74, PW 83–91, ASB 29–34, PSB 25–34, SD 57–68, SB 29–33, AP 28–34, H 51–56, AM 45–49, AL 45–49, PL 48–52, DS 34–36, VS 30–34, NDV 66–70, S 67–77, pa 278–314, pm 249–289, pp 289–316, Ip 841–902. Ukraine (Schluger 1966, Kudryashova 1998): AW 76, PW 90, ASB 34, PSB 31, SD 65, SB 32, AP 27, H 72, AM 49–59, AL 45–55, PL 62–72, DS 30–36, VS 36–42, NDV 73, S 85, pa 320, pm 286, pp 329, Ip 935. Slovakia (Kaluz & Vrabec 2014): AW 69–77, PW 83–91, ASB 31–36, PSB 23–29, SD 57–62, SB 31–36, AP 21–26, H 62–71, AM 55–67, AL 45–57, PL 62–69, DS 32–44, VS 30–38, NDV 64–80, S 77–101, pa 250–278, pm 232–270, pp 278–312, Ip 771–860.

Tribe Schoengastini Vercammen-Grandjean

Genus *Neoschoengastia* Ewing, 1929

***Neoschoengastia vietnamensis* Kaluz sp. nov.**
(Figs. 6–9)

Type material examined. Holotype: Location A: larva from *Turdus cardis* (Passeriformes: Turdidae), 4 February 2010. **Paratypes:** Location A: larva from *Niltava davidi* (Passeriformes: Muscicapidae), 1 February 2010; 14 larvae from *Larvivora sibilans* (Passeriformes: Muscicapidae), 5 February 2010; 2 larvae from *Turdus cardis* (Passeriformes: Turdidae), 1 February 2010; 12 larvae from *Turdus cardis*, 4 February 2010; 1 larva from *Pellorneum ruficeps* (Passeriformes: Pellorneidae), 1 February 2010.




FIGURE 6. *Neoschoengastia vietnamensis* sp. nov. (holotype)—idiosoma dorsal. Scale bar: 100 μ m.

Type depositions. Holotype and 10 paratypes are deposited in SNM; 10 paratypes in IEBR VAST; 10 paratypes in BMNH.

Etymology. The name of the new species is derived from the name of the country where we found type material of this chigger.

Description. Larva (n=12). SIF = 7B.S-B-3-3111.A000; fsp = 7-7-7; fPp = B.B.bBB; Ga = B; Pc = 3; fSc = AL > PL > AM; SB//PL; fCx = 1.1.1; fSt = 2.2; fD = 2H+8.6.6.4.4.2 = 32; DS = 32; VS = 28; NDV = 60; Ip = 869 (852–902).




FIGURE 7. *Neoschoengastia vietnamensis* sp. nov. (holotype)—idiosoma ventral. Scale bar: 100 μ m.

Idiosoma (Figs. 6, 7). Scutum nearly as wide as long, bearing AM, AL, PL setae and distally expanded (rounded) sensilla (20 long, 18 wide). Posterior part of scutum partially submerged beneath the cuticular striae of dorsum. Posterior striae create concentric pattern in each of lateral halves of scutum. Anterior margin of scutum sinuous with “shoulders”, lateral margins conical with small medial protrusion. Scutal puncta spread and cover anterior and lateral parts of scutum. Scutal ridge present connecting both bases of sensilla. Rounded eyes (2 + 2) in ocular plate, anterior eye larger than posterior. Humeral setae 57 (54–61) long, dorsal setae 46 (43–51) long,

arranged in regular rows, the posteriad rows varying in position of setae. Scutal measurements: AW 47 (46–49); PW 64 (63–67); SB 39 (36–42); ASB 19 (17–23); PSB 26 (20–29); SD 45 (37–51); P-PL 8 (6–10); AP 31 (29–34); AM 46 (43–51); AL 70 (62–78); PL 54 (51–59); S 20/18; fSc = AL > PL > AM. Sternal area with two pairs of sternal setae (fSt = 2.2), fCx = 1.1.1. Ventral side with 28 irregularly arranged ciliated setae. Lengths of ventral setae vary, increase from anterior 29 (26–32) to median 32 (29–36) and posterior 39 (36–43).

Gnathosoma (Fig. 8). Galeala barbed (Ga = B). Palps 71 (67–82) long, palpotibial claw 21 (18–23) long with 3 deeply indented prongs. Chelicera anteriorly with tricuspid cap.



FIGURE 8. *Neoschoengastia vietnamensis* sp. nov. (paratype)—gnathosoma (left—ventral, right—dorsal). Scale bar: 100 μ m.

Legs (Fig. 9). With pretarsus, claws and normal empodia. All setae on legs ciliated, slender. Specialized setae on leg segments—Leg I: Tarsus— S_1 17, f_1 3, nude pretarsala 11 ($PT' = N$), ciliated ST 32 and nude pST 20, 2 tibialae (anterior 20 and posterior 18), microtibiala 3, 3 genualae—anterior 17, median 23 and posterior 20, microgenuala 3; Leg II: Tarsus—nude pretarsala 11 ($PT'' = N$), S_2 15, f_2 3, 2 tibialae—anterior 16, posterior 16, genuala 17; Leg III: mastitarsala 56 (A-type; by Vercammen-Grandjean 1968) with 3–4 cilia, tibiala 17 and genuala 18. Number of barbed setae on leg segments (leg formula from coxa to tarsus): Leg I: 1-1-1-5-4-8-20; Leg II: 1-1-2-4-3-5-16; Leg III: 1-1-2-3-3-5-14. Length of tarsi I–III: I—82 (80–83), II—65 (63–69), III—81 (78–84). Width of tarsi I–III: I—18 (17–20); II—17 (16–18); III—17 (16–17). Leg length: pa = 308 (301–320); pm = 264 (258–274); pp = 297 (285–305); lp = 869 (848–902).




FIGURE 9. *Neoschoengastia vietnamensis* sp. nov. (paratype)—legs I–III (dots—bases of setae; filled dots—upper side; empty dots—bottom side. Numbers—lengths of solenidia and famuli. Scale bar: 100 μ m.

Differential diagnosis. The new species is very morphologically close to *Neoschoengastia posekany* Wharton and Hardcastle, 1946 but differs by having Ga = B; AW 47 (46–49); PW 64 (63–67); SB 39 (36–42). Moreover, tarsus III bears A-type (branched in base) mastisetæ and striae on the posterior part of the scutum are concentric. *N. posekany* differs from new species in having Ga = N; AW 75; PW 80; SB 47 and tarsus III with nude mastisetæ. In this species parallel striae on the scutum are posteriorly horizontal. Another species close to the newly described species is *Neoschoengastia heynemani* Nadchatram and Upham, 1966. *N. heynemani* differs from *Neoschoengastia vietnamensis* sp. nov. by SIF = 7B.S-N-3-3111.0000, Ga = N, fPp = B.N.NNB, clavate but narrow apical part of sensillum and by lacking mastisetæ on the tarsus III. Other differences are in the scutal formula (fSc: PL ≥ 2AM > AL) and length of leg III (pp. 263–270). On the contrary, *Neoschoengastia vietnamensis* sp. nov. has SIF = 7B.S-B-3-3111.A000; Ga = B, and fPp = B.B.NbB. The newly described species has also apically broad rounded sensilla, longer leg III (pp = 285–305) and fSc: PL > AM > AL, where PL is little longer than AM.

Neoschoengastia longitarsalis Schluger and Belskaya, 1966

Material examined: Location A: 1 larva from *Pellorneum ruficeps* (Passeriformes: Pellorneidae), 1 February 2010.

Distribution and hosts: *N. longitarsalis* is known from Turkmenistan from *Oenanthe picata* (Blyth) (Passeriformes: Muscicapidae), *Oenanthe finschii* (Heuglin) (Passeriformes: Muscicapidae), *Oenanthe isabellina* (Temminck) (Passeriformes: Muscicapidae), *Oenanthe hispanica* (Linnaeus) (Passeriformes: Muscicapidae), *Petronia petronia* Linnaeus (Passeriformes: Passeridae), *Coracias garullus* Linnaeus (Coraciiformes: Coraciidae), *Athene noctua* (Scopoli) (Strigiformes: Strigidae), and *Monticola saxatilis* (Linnaeus) (Passeriformes: Muscicapidae) (Kudryashova 1998). We found *N. longitarsalis* in Vietnam for the first time. *Pellorneum ruficeps* is a new host for this chigger.

Genus *Helenicula* Audy, 1954

Helenicula scanloni (Domrow and Nadchatram, 1964)

Material examined. Location A: 2 larvae from *Pellorneum ruficeps* (Passeriformes: Pellorneidae), 1 February 2010.

Distribution and hosts. In South Vietnam, *Helenicula scanloni* is reported to infest *Tupaia* sp. (Scandentia: Tapaiidae), *Menetes* sp. (Rodentia: Sciuridae), and *Rattus* sp. (Rodentia: Muridae); in Thailand, this chigger is reported to occur on birds *Gallus gallus* (Linnaeus) (Galliformes: Phasianidae), *Centropus sinensis* (Stephens) (Cuculiformes: Cuculidae), *Dicrurus hottentottus* (Linnaeus) (Passeriformes: Dicruridae), *Pellorneum ruficeps*, *Pomatorhinus hypoleucos* (Blyth) (Passeriformes: Timaliidae), and *Hydrornis oatesi* (Hume) (Passeriformes: Pittidae), as well as on mammalian hosts *Tupaia glis* (Scandentia: Tupaiidae), *Rattus rattus* (Linnaeus) (Rodentia: Muridae), and *Prionailurus bengalensis* (Kerr) (Carnivora: Felidae) (Nadchatram & Traub 1971).

Genus *Hypogastia* Vercammen-Grandjean, 1967

Hypogastia stekolnikovi Kaluz sp. nov.

(Figs. 10–14)

Material examined. Holotype: Location A: Larva from *Larviva sibilans* (Passeriformes: Muscicapidae); 1 February 2010. **Paratypes:** Location A: Seven larvae from *Niltava davidi* (Passeriformes: Muscicapidae); 3 February 2010.

Type depositions. Holotype and 2 paratypes will be deposited in SNM, 3 paratypes in IEBR VAST, and 2 paratypes in BMHN.

Etymology. The new species is named in honour of Dr. Alexandr A. Stekolnikov (Saint Petersburg, Russia) for his great contribution to taxonomy and ecology of chiggers.

Description. Larva (n=8). SIF = 7B.B.3.3111.3100; fsp = 7-7-7; fPp = B.B.BNB; Ga = B; Pc = 3; fSc = AL > PL > AM; SB//PL; fCx = 1.1.1; fSt = 2.2; fD = 2H+(8-9).6.6.4.4.2.2 = 34–35; DS = 34–35; VS = 36; NDV = 70; Ip = 914.




FIGURE 10. *Hypogastia stekolnikovi* sp. nov. (holotype)—idiosoma dorsal. Scale bar: 100 μ m.

Idiosoma (Figs. 10–12). Scutum nearly as wide as long, bearing AM, AL, PL setae and distally expanded (rounded) sensilla (20 long, 19 wide). Anterior margin of scutum sinuous with “shoulders”, lateral margins slightly conical. Scutal ridge present connecting both bases of sensilla. Posterior part of scutum partially submerged beneath the cuticular striae of dorsum. Posterior striae create concentric pattern in each of lateral halves of scutum, striae reach medially up to scutal ridge. Relatively large scutal puncta spread and cover nearly whole scutum in addition to small area posterior to bases of sensilla. Rounded eyes (2 + 2) in ocular plate, anterior eye larger than posterior. Humeral setae 50 (48–53) long, dorsal setae 43 (39–46) long, arranged in regular rows, posterior rows slightly

varying in position of setae. Scutal measurements: AW 56 (51–60); PW 72 (68–77); SB 45 (42–49); ASB 23 (20–26); PSB 31 (29–35); SD 54 (49–61); P-PL 9 (8–11); AP 38 (36–42); AM 42 (40–43); AL 53 (48–60); PL 49 (46–56); S 21/19; H 50 (48–53). Ciliated dorsal setae arranged in rows, the length of setae varying from anterior 44 (42–46) through median 40 (37–42) to posterior 37 (34–40). Sternal side with two pairs (fSt = 2.2) of sternal setae 33 (31–36), fCx = 1.1.1. Ventral side with 36 irregularly arranged ciliated setae. Lengths of ventral setae vary from anterior 32 (27–38) to median 31 (28–32) and posterior 33 (26–37).




FIGURE 11. *Hypogastia stekolnikovi* sp. nov. (holotype)—idiosoma ventral. Scale bar: 100 μ m.

Gnathosoma (Fig. 13). Galeala barbed (Ga = B). Palps 68 (66–71) long, palpotibial claw 17 (14–21) long with 3 deeply indented prongs. Chelicera anteriorly with tricuspid cap.

Legs (Fig. 14). With pretarsus, claws and normal empodia. All setae on legs ciliated, slender. Specialized setae

on leg segments—Leg I: Tarsus— S_1 21, f_1 2, nude pretarsala 10 ($PT' = N$), nude ST 21 and pST 18; 2 tibialae (anterior 16 and posterior 17), microtibiala 2, 3 genualae—anterior 16, median 16 and posterior 14, microgenuala 3; Leg II: Tarsus—nude pretarsala 12 ($PT'' = N$), S_2 18, f_2 2, 2 tibialae—anterior 14, posterior 14, genuala 15; Leg III: 2 nude mastitarsalae, anterior 43 and posterior 40, in some specimens moreover one type-A mastitarsala closed to posterior nude one; tibiala 15 (in some specimens, moreover, one type-A mastitibiala 36 long with 2–3 cilia in basal half) and genuala 17. Number of barbed setae on leg segments (leg formula from coxa to tarsus): Leg I: 1-1-(1+5)-4-8-20; Leg II: 1-1-(2+4)-3-5-16; Leg III: 1-1-(2+3)-3-5-14. Length of tarsi I–III: I—83 (77–85), II—67 (63–71), III—85 (80–89). Width of tarsi I–III: I—18 (18–20); II—17 (16–18); III—17 (17–18). Leg length: pa = 320 (297–336); pm = 268 (255–278); pp = 312 (301–324); Ip = 899 (868–938).

Differential diagnosis. The new species is very similar to *Hypogastia fullbergae* Brennan, 1948 but differs from it by fPp = B.B.BNB, DS = 34–35; VS = 36; NDV = 70 and by longer legs (Ip = 899 [868–938]); while *H. fullbergae* has fPp = B.B.NBB; DS = 30; VS = 28; NDV = 58; and shorter legs (Ip = 718–811). *Hypogastia stekolnikovi* sp. nov. also resembles the species *Hypogastia xiaguanensis* Yu, Hu and Zhang, 1983, but differs from it by fPp = B.B.BNB, Ga = B, fSc = AL > PL > AM and longer legs, while *H. xiaguanensis* has fPp = B.B.NNB, Ga = N; fSc = PL > AM > AL and shorter legs (Ip = 737).




FIGURE 12. *Hypogastia stekolnikovi* sp. nov. (holotype)—scutum. Scale bar: 100 μ m.

Discussion

We did not find trombiculids reported formerly from North Vietnam (Schluger *et al.* 1960), such as *Leptotrombidium horrida* Schluger, 1960, *Leptotrombidium arvina* Schluger, 1960, *Leptotrombidium gracipalpis* Schluger, 1960, *Leptotrombidium globosa* Schluger, 1960, *Leptotrombidium monstrosa* Schluger, 1960, *Leptotrombidium magna* Schluger, 1960, and *Leptotrombidium deliense* (Walch, 1922).

We did not find other schoengastinids, including chiggers of the genera *Hypogastia* and *Neoschoengastia*, parasitizing mainly birds. The species similar to *Hypogastia stekolnikovi* sp. nov. and *Neoschoengastia vietnamensis* sp. nov. are *H. xiaguanensis*, *N. posekany*, and *N. heyne*. *H. xiaguanensis* is reported to infest *Apodemus chevrieri* (Milne-Edwards) (Rodentia: Muridae) in China (Yu *et al.* 1983). *N. posekany* is reported to

infest hens in Hanoi, North Vietnam while in Malaysia it is reported to infest another birds *Streptopelia orientalis* (Latham) (Columbiformes: Columbidae), *Columba janthina* Temminck (Columbiformes: Columbidae), *Turdus celanops* Stejneger (Passeriformes: Turdidae), *Monticola solitarius* (Linnaeus) (Passeriformes: Muscicapidae), *Saxicola torquatus* (Linnaeus) (Passeriformes: Muscicapidae), and *Chloris sinica* (Linnaeus) (Passeriformes: Fringillidae) (Schluger *et al.* 1960b). *Neoschoengastia heynemani* is reported to infest kingfishers *Alcedo meninting* Horsfield (Coraciiformes: Alcedinidae) and *Alcedo euryzona* Temminck in Malaysia (Coraciiformes: Alcedinidae) (Nadchatram & Upham 1966). We did not find *Neoschoengastia gallinarum* (Hatori 1920) reported from chicken in North Vietnam (Schluger *et al.* 1960b) Although the newly described schoengastinid species from Vietnam were found on birds, in consideration of the findings of other, similar chiggers from that area, they may occur on a wider spectrum of hosts, including local mammals.



FIGURE 13. *Hypogastia stekolnikovi* sp. nov. (paratype)—gnathosoma (left—ventral, right—dorsal). Scale bar: 100 μ m.




FIGURE 14. *Hypogastia stekolnikovi* **sp. nov.** (paratype)—legs I–III (dots—bases of setae; filled dots—upper side; empty dots—bottom side. Numbers—lengths of solenidia and famuli. Scale bar: 100 μ m.

Acknowledgements

The study was supported in part by a grant No. LC06073 of the Ministry of Education, Youth and Sports of the Czech Republic and the Institutional Support of the IVB AS CR, v. v. i. (RVO: 68081766).

References

- Audy, J.R. (1954) Malaysian parasites, IX. Notes on the taxonomy of trombiculid mites with description of a new subgenus. *Studies of the Institute of Medical Research, Malayan*, 26, 123–170.
- Brennan, J.M. (1948) New North American chiggers (Acarina, Trombiculidae). *Journal of Parasitology*, 34, 465–478.
<http://dx.doi.org/10.2307/3273313>
- Brennan, J.M. (1951) Two new species of *Neoschöngastia* with a key to the species of the world (Acarina: Trombiculidae). *Journal of Parasitology*, 37, 577–582.
<http://dx.doi.org/10.2307/3273352>
- Brennan, J.M. (1962) Four new chiggers from Mexico. *Journal of Parasitology*, 48, 618–620.
<http://dx.doi.org/10.2307/3274924>
- Brennan, J.M. (1965) Two new species and other records of chiggers from Texas (Acarina: Trombiculidae). *Acarologia*, 7, 79–83.
- Brennan, J.M. & Amerson, A.B. (1971) Six new species and additional records of chiggers from the Central Pacific (Acarina: Trombiculidae). *Journal of Parasitology*, 57, 1311–1317.
<http://dx.doi.org/10.2307/3277989>
- Domrow, R. & Nadchatram, M. (1964) Four Thai chiggers with expanded sensillae (Acarina, Trombiculidae). *Acarologia*, 6, 476–483.
- Ewing, H.E. (1929) Four new species of chiggers (Acarina-Trombidiidae). *Entomological News*, 40, 294–297.
- Gill, F.B. & Donsker, D.B. (Eds.) (2015) IOC World Bird List (v. 5.2).
<http://dx.doi.org/10.14344/IOC.ML.5.2>
- Goff, M.L. (1983) Studies on Papua New Guinea chiggers (Acari: Trombiculidae) X. Two new species of Schoengastiini. *Journal of Medical Entomology*, 20, 40–43.
<http://dx.doi.org/10.1093/jmedent/20.1.40>
- Hatori, J. (1920) Tsutsugamushi disease in Formosa (5th communication). *Taiwan Igakkai Zasshi (Journal of the Medical Association of Formosa)*, 209, 317–352.
- Hirst, S. (1925) The adult form of the “harvest bug”. *Nature*, 116, 609.
- Kalúz, S. & Vrabec, M. (2014) The chigger *Neotrombicula elegans* (Acari: Trombiculidae) in Slovakia. *Folia Faunistica Slovaca*, 19, 99–102.
- Kudryashova, N.I. (1998) *Kleshchi krasnotelki (Acariformes, Trombiculidae) vostochnoy Palearktiki*. KMK Scientific Press, Moscow, 342 pp.
- Mammal Species of the World (2015) 3rd Edition (MSW3), *Wilson & Reeder's Mammal Species of the World*, American Society of Mammalogists, Johns Hopkins University Press. Available from: <http://vertebrates.si.edu/msw/mswCFApp/msw/index.cfm> (accessed 16 November 2015)
- Nadchatram, M. (1963) The larva and nymph of *Odontacarus audyi* (Radford) (Acarina: Trombiculidae). *Pacific Insects*, 5, 535–540.
- Nadchatram, M. & Traub, R. (1971) Chiggers of the genus *Helenicula* of the Old World including descriptions of 9 new species (Acarina: Prostigmata, Trombiculidae). *Journal of Medical Entomology*, 8, 562–597.
<http://dx.doi.org/10.1093/jmedent/8.5.562>
- Nadchatram, M. & Upham, R.W. Jr. (1966) Two new larval trombiculid mites (Acarina, Trombiculidae) collected from ground holes and birds in Malaysia. *Journal of Medical Entomology*, 3, 345–350.
- Nagayo, M., Miyagawa, Y., Mitamura, T. & Imamura, A. (1916) On the nymph and prosopon of the tsutsugamushi, *Leptotrombidium akamushi*, n. sp. (*Trombidium akamushi* Brumpt), carrier of the tsutsugamushi disease. *Journal of Experimental Medicine*, 25, 255–272.
<http://dx.doi.org/10.1093/jmedent/3.3-4.345>
- Radford, C.D. (1946) New species of larval mites (Acarina: Trombiculidae) from Manipur State, India. *Proceedings of the Zoological Society of London*, 116, 247–265.
<http://dx.doi.org/10.1111/j.1096-3642.1946.tb00122.x>
- Robson, C. (2005) *New Holland Field Guide to the Birds of South-East Asia*. New Holland Publishers, London, 304 pp.
- Schluger, E.G. (1966) Novye vidy kleshchey podsemejstva Trombiculidae Ewing (Acariformes, Trombiculidae). In: Mazurmovich, B.N. (Ed.), *Parazity, promezhutochnye khozyaeva i perenoshchiki*. Naukova Dumka, Kiev, pp. 208–215. [in Russian]
- Schluger, E.G. & Belskaya, G.S. (1966) Novye vidy kleshchey-krasnotelok (Acariformes, Trombiculidae) s ptic Turkmenii. *Izdavatelstvo Akademii Nauk Turkmensoy SSR, Seriya Biologicheskikh Nauk. Ashkhabad*, 1, 64–70. [in Russian]
- Schluger, E.G., Grokhovskaya, I.M., Dan, V.N., Nguyen, X.H. & Do, K.T. (1960a) The species of the subgenus *Leptotrombidium* (Acariformes, Trombiculidae) from North Vietnam. *Zoologicheskii Zhurnal*, 39, 1791–1801. [in Russian]
- Schluger, E.G., Grokhovskaya, I.M., Dan, V.N., Nguyen, X.H. & Do, K.T. (1960b) Fauna of the chiggers (Acariformes, Trombiculidae) of North Viet-Nam. *Parazitologicheskii Sbornik Zoologicheskogo Instituta Akademii Nauk SSSR*, XIX, 169–193. [in Russian with English summary]
- Schluger, E.G., Grokhovskaya, I.M., Dan, V.N., Nguyen, X.H. & Do, K.T. (1960c) The trombiculid mites of the genus

- Gahrlepiea* (Acariformes, Trombiculidae) from North Viet-Nam. *Revue d'Entomologie de l'URSS*, XXXIX, 2, 462–476. [in Russian]
- Schluger, E.G., Grokhovskaya, I.M., Dan, V.N., Nguyen, X.H. & Do, K.T. (1961) The trombiculid mites (chigger mites) of the genera *Doliosis* Oudemans, 1910, and *Traubacarus* Audy et Nadchatram, 1957 (Acariformes, Trombiculidae) from North Viet-Nam. *Revue d'Entomologie de l'URSS*, XL, 2, 448–453. [in Russian]
- Schluger, E.G., Grokhovskaya, I.M., Dan, V.N., Nguyen, X.H. & Do, K.T. (1963) Chigger mites of the genus *Trombicula* (Acariformes, Trombiculidae) from Democratic Republic Viet Nam. *Revue d'Entomologie de l'URSS*, XLII, 3, 691–701. [in Russian]
- Shatrov, A.B. & Kudryashova, N.I. (2008) Taxonomic ranking of major trombiculid subtaxa with remarks on the evolution of host-parasite relationship (Acariformes: Parasitengona: Trombiculidae). *Annales Zoologici*, 58, 279–287.
- Stan Fernandes, S.J. & Kulkarni, S.M. (2003) *Studies on the Trombiculid Mite Fauna of India*. Zoological Survey of India, Kolkata, 539 pp.
- Stekolnikov, A.A. (2004) Variability in *Leptotrombidium europaeum* and two new related chigger mite species (Acari: Trombiculidae) from Caucasus. *Parazitologiya*, 38, 388–405. [in Russian]
- Stekolnikov, A.A. (2013) *Leptotrombidium* (Acari: Trombiculidae) of the world. *Zootaxa*, 3728 (1), 1–173. <http://dx.doi.org/10.11646/zootaxa.3728.1.1>
- Tian, Q. & Wen, T. (1984) Two new species of the genus *Leptotrombidium* from Shanxi Province (Acariformes: Trombiculidae). In: Wen, T. (Ed.), *Sand mites of China (Acariformes: Trombiculidae and Leeuwenhoekidae)*. Hue Lin Publishing House, Peking, pp. 43–45. [in Chinese]
- Traub, R. & Lakshana, P. (1966) Some chiggers of the subgenus *Leptotrombidium* from Thailand, with descriptions of new species (Acarina, Trombiculidae). *Journal of Medical Entomology*, 3, 271–292. <http://dx.doi.org/10.1093/jmedent/3.3-4.271>
- Vercammen-Grandjean, P.H. (1967) Notes on the Trombiculidae. *Acarologia*, 9, 127–134.
- Vercammen-Grandjean, P.H. (1968) *The Chigger Mites of the Far East (Acarina: Trombiculidae & Leeuwenhoekidae)*. An Illustrated Key and a Synopsis; Some New Tribes, Genera and Subgenera. U. S. Army Medical Research and Development Command, Washington D.C., 135 pp.
- Vercammen-Grandjean, P.H. & Langston, R. (1976) *The Chigger Mites of the World (Acarina Trombiculidae)*. III. *Leptotrombidium complex*. George Williams Hooper Research Foundation, University of California, San Francisco, 1061 pp.
- Vercammen-Grandjean, P.H., Rohde, C.J. Jr. & Mesghali, H. (1970) Twenty larval Trombiculidae (Acarina) from Iran. *Journal of Parasitology*, 56, 773–806. <http://dx.doi.org/10.2307/3277727>
- Walch, E.W. (1922) Over *Trombicula deliensis* n. sp., vermoedelijke overbrengster der Pseudotyphus, en andere Trombiculae van Deli (Eerste Mededeeling). *Geneeskundig Tijdschrift voor Nederlandsch-Indië*, 62, 530–588.
- Wang, D.Q., Liao, H.R. & Lin, Z.H. (1981) Five new trombiculid mites of the genus *Leptotrombidium* (Acarina, Trombiculidae). *Acta Zootaxonomica Sinica*, 6, 44–52. [in Chinese]
- Wen, T. & Xiang, R. (1984) A new species and a new subspecies of avian *Leptotrombidium* from Yunnan Province (Acariformes: Trombiculidae). In: Wen, T. (Ed.), *Sand Mites of China (Acariformes: Trombiculidae and Leeuwenhoekidae)*. Hue Lin Publishing House, Peking, pp. 213–215. [in Chinese]
- Wharton, G.W. & Hardcastle, A.B. (1946) The genus *Neoschoengastia* (Acarinida: Trombiculidae) in the Western Pacific Area. *Journal of Parasitology*, 32, 286–322. <http://dx.doi.org/10.2307/3272682>
- Yu, Z.Z., Hu, G. & Zhang, L.Q. (1983) Seven species of chigger mites from China (Acarina: Trombiculidae). *Acta Zootaxonomica Sinica*, 8, 63–74. [in Chinese]