

Hotspots of mite new species discovery: Trombidiformes (2013–2015)

JIAN-FENG LIU¹ & ZHI-QIANG ZHANG^{1,2}

¹ School of Biological Sciences, the University of Auckland, Auckland, New Zealand

² Landcare Research, 231 Morrin Road, Auckland, New Zealand; corresponding author: email: ZhangZ@landcareresearch.co.nz

Abstract

This paper reveals the hotspots of new mite discovery through of a survey of type localities of new Trombidiformes species described in two journals (*Systematic & Applied Acarology* and *Zootaxa*) during the last three years (2013–2015). Taxonomically, the 491 new species of the Trombidiformes are highly unevenly distributed among 55 families with top 10 families accounting for over 66% of the total. The Eriophyidae is the top-ranked family. Geographically, these 491 new species are from 55 countries around the world and their distribution among the countries is highly uneven. The majority of these new species (69%) are from the top 10 countries and six of the top ten countries are also megadiversity countries. The top three countries are all from Asia (Iran, China and Malaysia) and they together accounted for over one third of all new species of the Trombidiformes described in the two journals during 2013–2015.

Key words: Mites, Trombidiformes, new species, hotspots, type locality, type depository

Introduction

Discoveries of new species around the world are unevenly distributed; some countries are hotspots for the discovery of new species because they are hotspots of biodiversity (Mittermeier 1988; Gaston 2000) with more undescribed species or a higher concentration of taxonomists (both local and overseas) interested in working on the biodiversity of these regions (or a combination of both). A recent survey of new mite species described during six years (2007–2012) showed that the distribution of 1379 new species are highly unevenly distributed among 92 countries, with the top 10 countries accounting for 62% of the total (Liu *et al.* 2013). This paper is a follow-up of Liu *et al.* (2013) and provides a survey of new species of the Trombidiformes described during the last three years (2013–2015). The methods of Liu *et al.* (2013) are followed here and the patterns of distribution of new species of the order Trombidiformes described during 2013–2015 are summarised. The bulk of this paper presents a list of the type localities and depositories of all new species of the Trombidiformes published in *Zootaxa* and *Systematic & Applied Acarology* (*SAA*) during the last three years (2013–2015) with the main purpose of exposing newly described species to a wider community as similar papers in the past (*e.g.* Fan 2006; Liu *et al.* 2013; Zhang 2014).

Abbreviations of zoological collections

ACIAU—Acarology Collection, Islamic Azad University, Fars Science and Research Branch, Marvdasht, Iran.

AFUMI—Acarological Collection, Department of Plant Protection, Faculty of Agriculture, University of Maragheh, Iran.

AMS—Australian Museum, Sydney, Australia.

AMU—Department of Animal Morphology, Adam Mickiewicz University, Poznań, Poland.

ARC-PPRI—Plant Protection Research Institute-Agricultural Research Center - Fruit Acarology Department, Dokki, Cairo, Egypt.

BASU—Collection of Acarology Laboratory, University of Bu-Ali Sina, Hamadan, Iran.

CAS—California Academy of Sciences, San Francisco, California, USA.

CALBS—Collection of the Acarology Laboratory, University of Bu Ali Sina, Hamedan, Iran.

- DiSSPA—Department of Soil, Plant and Food Sciences, section of Entomology and Zoology, University of Bari, Italy and at the University of Yüzüncü Yıl, Van, Turkey.
- ESALQ/USP—Collection of Departamento de Entomologia e Acarologia, ESALQ – Universidade de São Paulo, Piracicaba, São Paulo, Brazil.
- EUE—Acarology Laboratory of Erzincan University, Erzincan, Turkey.
- FFSDU—Fundamental Science Laboratory collection, Faculty of Fisheries, Süleyman Demirel University, Egirdir/Isparta, Turkey.
- FUE—Research collection of the Biology Department, Fırat University, Elazığ, Turkey.
- GUGC—Istitute of Entomology, Guizhou University, Guiyang, P.R. China.
- GUGX—Guangxi University, Guangxi Zhuang Autonomic Region, P.R. China.
- IARI—National Pusa Collection, Division of Entomology, Indian Agricultural Research Institute, New Delhi, India.
- IBIW—Research collection of the Institute for Biology of Inland Waters, Borok, Russia.
- IBSS—Research collection of the Institute of Biology and Soil Science, Vladivostok, Russia.
- IPHCMC—Institut Pasteur, Ho Chi Minh City, Vietnam.
- JAZM—Jalal Afshar Zoological Museum, College of Agriculture, University of Tehran, Iran.
- KUB—Insect museum of Department of Entomology, Faculty of Agriculture, Kasetsart University, Bangkok 10900, Thailand.
- KSU—Acarology Laboratory, Department of Plant Protection, College of Food and Agriculture Sciences, King Saud University, Saudi Arabia.
- MAGNT—Museum and Art Gallery of the Northern Territory, Bullocky Point, Fannie Bay, Northern Territory, 0820, Australia.
- MNCN—Museo Nacional de Ciencias Naturales, Madrid, Spain.
- MNHN—Muséum National d'Histoire naturelle, Paris, France.
- MNHP—Museum of the Natural History of Podgorica, Montenegro.
- MNHWU—Museum of Natural History, Wrocław University, Poland.
- MNZ—Museum of New Zealand, Wellington, New Zealand.
- MZLQ—Mite Reference Collection, Department of Entomology and Acarology, Escola Superior de Agricultura "Luiz de Queiroz", Brazil.
- MZUC—Museo de Zoología Universidad de Concepción, Concepción, Bío Bío Region, Chile.
- NBG—Collection of the Nikita Botanical Gardens—National Scientific Center, Yalta, Crimea, Ukraine.
- NCA-PPRI—National Collection of Arachnida, PPRI, Pretoria, South Africa.
- NIBR—National Institute of Biological Resources, South Korea.
- NHMB—Natural History Museum Belgrade, Serbia.
- NHMG—Natural History Museum of Geneva, Switzerland.
- NHMW—Naturhistorisches Museum, Wien, Austria.
- NJAU—Arthropod/Mite Collection of the Department of Entomology, Nanjing Agricultural University, Jiangsu Province, P.R. China.
- NMV—Museum Victoria, Melbourne, Australia.
- NSMT—Collection of the National Museum of Nature and Science, Tokyo, Japan.
- NTM—Museum and Art Gallery of the Northern Territory, Darwin, Australia.
- NZAC—New Zealand Arthropod Collection, Landcare Research, Auckland, New Zealand.
- NZMC—National Zoological Museum of China, Institute of Zoology, Chinese Academy of Sciences, Beijing, P.R. China.
- OSAL—Ohio State University Acarology Collection, Columbus, USA.
- PEU—Acari Collection of Departamento de Zoológia e Botânica (DZSJR), Universidade Estadual Paulista (UNESP), campus de São José do Rio Preto, São Paulo, Brazil.
- PPRP—ARC-Plant Protection Research Institute, Pretoria, South Africa.
- QM—Queensland Museum, Brisbane, Australia.
- RMNH—National Museum of Natural History, Leiden, The Netherlands.
- SBUC—Collection of the Acarology Laboratory, Shahid Bahonar University of Kerman, Kerman, Iran.
- SHZU—Department of Plant Protection, College of Agriculture, Shihezi University, Shihezi, Xinjiang 832003, P.R. China.
- SMF—Senckenberg-Museum Frankfurt, Germany.
- SNMB—Slovak National Museum, Bratislava, Slovakia.
- TMU—Acarological Collection, Department of Entomology, Faculty of Agriculture, Tarbiat Modares University, Tehran, Iran.
- TUMZ—the acarological collection of the Tyumen State University Museum of Zoology, Tyumen, Russia.
- UBS—Acarology Collection, Department of Entomology and Agricultural Zoology, Faculty of Agriculture, University of Belgrade, Serbia.

UFMG-AC—Acarological collection of the Federal University of Minas Gerais, Pampulha, Belo Horizonte—MG, Brazil
 UMMZ—University of Michigan Museum of Zoology, Ann Arbor, Michigan, USA.
 UNAM—Acarology Collection of the Universidad Autónoma de Mexico, Mexico City, Mexico.
 UNESP—Departamento de Zoologia e Botânica, Universidade Estadual Paulista, São José do Rio Preto, State of São Paulo, Brazil.
 ZAUMCN—Museu de Ciências Naturais, UNIVATES - Centro Universitário, Lajeado, Rio Grande do Sul State, Brazil.
 USNM—United States Natural History Museum, Smithsonian Institution, Washington, D.C., USA.
 UTI—Acarological Collection, Department of Plant Protection, Faculty of Agriculture, University of Tabriz, Tabriz, Iran.
 UWMO—Collection of the Department of Ecology and Environmental Protection, the University of Warmia and Mazury in Olsztyn, Poland.
 WAM—Western Australian Museum, Perth, Australia.
 ZISP—Zoological Institute, Russian Academy of Sciences, Saint-Petersburg, Russia.
 ZIZMUH—Biozentrum Grindel und Zoologisches Museum (formerly: Zoologisches Institut und Museum der Universität Hamburg), Hamburg, Germany.

Results and discussion

Trends in the number of new species

SAA published 95 new species of the Trombidiformes during 2013–2015 (Table 1). This is about 32 species per year, or about two and a half as many as the average during 2007–2012 (13 species per year). This increase is a reflection of the increase in the number of papers published in this journal during the last three years (Zhang 2014, 2015, 2016). In 2015, the frequency of *SAA* increased to eight issues from four issues (2012–2014) and this caused the sudden increase of new species in 2015. *Zootaxa* published 396 new species of the Trombidiformes during 2013–2015 (Table 1). This is an average of 132 species per year, which is only 8% more than the average during 2007–2012 (122). In comparison, the average number of new Scarcoptiformes species increased 56.9% (Li & Zhang 2016), whereas that of the Parasitiformes decreased 17.5% over the same period (Lam & Zhang 2016).

TABLE 1. The comparison of the numbers of new species of the order Trombidiformes described in *SAA* and *Zootaxa* between 2013–2015 and 2007–2012. Data of 2007–2012 from Liu *et al.* (2013).

Year	<i>SAA</i>	<i>ZOOTAXA</i>	Total
2007	12	70	82
2008	6	103	109
2009	9	165	174
2010	12	115	127
2011	21	134	155
2012	19	126	145
2013	19	114	133
2014	21	171	192
2015	55	111	166
Total	174	1109	1283

Taxonomic distribution of new species

The distribution of 491 new species of the Trombidiformes described during 2013–2015 is uneven at the infraorder level (Table 2). Overall, the proportion of new species of the Eupodina decreased to 23.01% and that of Eleutherengona increased to 38.09% during 2013–2015 when compared to those during 2007–2012 (Table 2). These patterns reflect mainly the increase in the number of some new Eleutherengona during 2013–2015 when four of top ten families are of this group: Harpirhynchidae, Tenuipalpidae, Syringophilidae and Stigmaeidae (Table 3A, B).

The 491 new species of Trombidiformes are unevenly distributed among 55 families (Table 3A). The top 10 families account for over 66% of the total number of species, whereas 26 families (47% of the total) have 1–3 species each. The Eriophyidae is the top-ranked family (Table 3) as it was during 2007–2012 (Liu *et al.* 2013).

TABLE 2. The comparison of taxonomic distribution of the new species of the Trombidiformes described in *SAA* and *Zootaxa* between 2013–2015 and 2007–2012. Data of 2007–2012 after Liu *et al.* (2013).

Year	Infraorder	SAA	ZOOTAXA	Total (% of the order Trombidiformes)
2007–2012	Eupodina	14	303	317 (39.97%)
	Anystina	35	265	300 (37.83%)
	Eleutherengona	30	146	176 (22.19%)
2013–2015	Eupodina	46	67	113 (23.01%)
	Anystina	17	174	191 (38.90%)
	Eleutherengona	32	155	187 (38.09%)

TABLE 3A. Taxonomic distribution new species of the Trombidiformes from *SAA* and *Zootaxa* among families (2013–2015); N_{sp}=number of new species within each paper, N_{fam.}=number of families with new species.

N _{sp}	N _{fam.}	Families of Order Trombidiformes
53	1	Eriophyidae
47	1	Harpirhynchidae
38	1	Hygrobatidae
34	1	Tenuipalpidae
32	1	Syringophilidae
30	1	Aturidae
27	1	Torrenticolidae
23	2	Diptilomiopidae, Trombiculidae
19	1	Stigmaeidae
12	1	Tydeidae
10	1	Unionicolidae
9	2	Cunaxidae, Tetranychidae
8	3	Arrenuridae, Podapolipidae, Pterygosomatidae,
7	1	Scutacaridae
6	2	Erythraeidae, Pygmephoridae
5	5	Bdellidae, Camerobiidae, Hydrachnidae, Microtrombidiidae, Phytoptidae
4	4	Halacaridae, Limnesiidae, Pseudocheylidae, Trombidiidae
3	4	Neopygmephoridae, Pionidae, Pontarachnidae, Tarsonemidae
2	7	Anisitsiellidae, Athienemanniidae, Calyptostomatidae, Hydryphantidae, Johnstonianidae, Oxidae, Sperchontidae
		Achaemenothrombiidae, Allochaetophoridae, Caeculidae, Caligonellidae, Cryptognathidae,
1	15	Dolichocybidae, Eupalopsellidae, Eupodidae, Frontipodopsidae, Iolinidae, Limnocharidae, Mideopsidae, Paratydeidae, Tarsocheylidae, Teneriffiidae

TABLE 3B. The comparison of the numbers of new species in the order Trombidiformes from *SAA* and *Zootaxa* between 2013–2015 and 2007–2012. The classification of the Trombidiformes follows that of Zhang *et al.* (2011). Data of 2007–2012 after Liu *et al.* (2013).

TAXA	2013–2015	2007–2012
Suborder Prostigmata		
Infraorder Eupodina		
Superfamily Bdelloidea		
Family Bdellidae	5	6
Family Cunaxidae	9	36
Superfamily Eriophyoidea		
Family Phytoptidae	5	5
Family Eriophyidae	53	204
Family Diptilomiopidae	23	48
Superfamily Eupodoidea		
Family Eupodidae	1	—
Family Penthalodidae	—	3
Family Rhagidiidae	—	8
Superfamily Halacaroidea		
Family Halacaridae	4	7
Superfamily Tydeoidea		
Family Tydeidae	12	—
Family Iolinidae	1	—
Infraorder Anystina		
Hypoder Anystae		
Superfamily Adamystoidea		
Family Adamystidae	—	5
Superfamily Anystoidea		
Family Pseudocheylidae	4	—
Family Teneriffiidae	1	—
Superfamily Caeculoidea		
Family Caeculidae	1	—
Superfamily Paratydeoidea		
Family Paratydeidae	1	—
Hypoder Parasitengona		
Superfamily Calyptostomatoidea		
Family Calyptostomatidae	2	1
Superfamily Erythraeoidea		
Family Erythraeidae	6	21
Superfamily Eylaoidea		
Family Limnocharidae	1	—
Superfamily Allotanaupodoidea		
Family Allotanaupodidae	—	5

.....continued on the next page

TABLE 3B. (Continued)

TAXA	2013–2015	2007–2012
Superfamily Chyzerioidea		
Family Chyzeriidae	—	1
Superfamily Trombiculoidae		
Family Johnstonianidae	2	
Family Leeuwenhoekiidae	—	1
Family Trombiculidae	23	19
Superfamily Trombidioidea		
Family Achaemenothrombiidae	1	1
Family Neothrombiidae	—	2
Family Microtrombidiidae	5	3
Family Trombidiidae	4	4
Superfamily Hydryphantoidea		
Family Hydryphantidae	2	3
Family Hydrodromidae	—	6
Family Teratothyadidae	—	1
Superfamily Hydrachnoidea		
Family Hydrachnidae	5	2
Superfamily Lebertioidea		
Family Sperchontidae	2	12
Family Teutonidae	—	1
Family Anisitsiellidae	2	26
Family Lebertiidae	—	1
Family Oxidae	2	
Family Torrenticolidae	27	32
Superfamily Hygrobatoidea		
Family Pontarachnidae	3	4
Family Limnesiidae	4	6
Family Frontipodopsidae	1	—
Family Hygrobatidae	38	39
Family Aturidae	30	44
Family Feltriidae	—	11
Family Unionicolidae	10	8
Family Pionidae	3	—
Superfamily Arrenuroidea		
Family Momoniidae	—	2
Family Mideopsidae	1	2
Family Athienemanniidae	2	1
Family Chappuisididae	—	1
Family Arrenuridae	8	35

.....continued on the next page

TABLE 3B. (Continued)

TAXA	2013–2015	2007–2012
Infraorder Eleutherengona		
Hypoder Raphignathina		
Superfamily Cheyletoidea		
Family Cheyletidae	—	8
Family Harpirhynchidae	47	—
Family Syringophilidae	32	68
Superfamily Myobioidea		
Family Myobiidae	—	3
Superfamily Pterygosomatoidea		
Family Pterygosomatidae	8	3
Superfamily Raphignathoidea		
Family Barbutiidae	—	1
Family Caligonellidae	1	1
Family Camerobiidae	5	7
Family Cryptognathidae	1	1
Family Eupalopsellidae	1	—
Family Raphignathidae	—	1
Family Stigmaeidae	19	20
Superfamily Tetranychoidae		
Family Allochaetophoridae	1	—
Family Linotetranidae	—	3
Family Tenuipalpidae	34	24
Family Tetranychidae	9	12
Hypoder Heterostigmata		
Superfamily Dolichocyboidea		
Family Dolichocybidae	1	—
Superfamily Pyemotoidea		
Family Caraboacaridae	—	1
Family Pyemotidae	—	2
Superfamily Pygmephoroidae		
Family Pygmephoridae	6	2
Family Scutacaridae	7	1
Family Neopygmephoridae	3	3
Superfamily Tarsocheyloidea		
Family Tarsocheylidiae	1	—
Superfamily Tarsonemoidea		
Family Podapolipidae	8	8
Family Tarsonemidae	3	5
Superfamily Trochometridioidea		
Family Trochometridiidae	—	1

Geographic distribution of new species

The 491 new species of the Trombidiformes described during 2013–2015 are from 55 countries around the world and the distribution of these species among the countries is highly uneven (Table 4). The majority of these new species (69%) are from the top 10 countries. It should be noted that (1) six of the top ten countries of new species discovery are also megadiversity countries (Mittermeier 1988); and (2) the top three countries are all from Asia (Iran, China and Malaysia) and they together accounted for over one third of all new species of the Trombidiformes described in *Zootaxa* and *SAA* during 2013–2015. Over 70% of the 55 countries have only 1–3 new species per country. In comparison, this level of patchiness is higher than that of new mite species described during 2007–2012: 62% of new species were from the top 10 countries and 40% of the countries had only 1–3 new species per country (Liu *et al.* 2013).

TABLE 4. Geographic distribution of the order Trombidiformes new species from *SAA* and *Zootaxa* (2013–2015) among countries (or regions); N_{sp} =number of new species within each country (or region), N_c =number of countries with new species.

N_{sp}	N_c	Name of countries (regions)
59	1	Iran
57	1	P.R. China*
54	1	Malaysia*
45	1	Australia*
30	1	Russia
29	1	USA*
24	1	Brazil*
15	1	Indonesia*
13	1	Chile
12	1	Turkey
11	1	Vietnam
10	1	Thailand
9	1	Hungary
8	1	Ghana
7	2	Kenya, Saudi Arabia
6	5	New Zealand, Laos, Mexico*, The Netherlands, Ukraine
5	2	Guyana, South Korea
4	3	Argentina, Comoros, Japan
3	8	Canada, Ethiopia, India*, Italy, Nigeria, Peru*, Serbia, Tanzania
2	4	Belarus, Gabon, Slovakia, North America (an unknown locality) Cameroon, Dominican Republic, Egypt, Guatemala, Greece, British West Indies, Cambodia, Madagascar*,
1	17	Macedonia, Mozambique, Oman, Papua New Guinea*, Portugal, South Africa*, Spain, Suriname, Trinidad and Tobago

*Megadiversity countries (Mittermeier 1988).

List of type localities and depositories of new species of the Order Trombidiformes in SAA 2013–2015

Order **Trombidiformes** (95 species)

Suborder **Prostigmata** (95 species)

Infraorder **Eupodina** (46 species)

Superfamily **Bdelloidea** (2 species)

Family **Bdellidae** (1 species)

Spinibdella tabarii Paktinat-Saejj & Bagheri, 2015 *in* Paktinat-Saejj, Bagheri, Saboori, Seilsepour & Ueckermann, 2015: 704—Holotype male (AFUMI), from the soil and rotten leaves under citrus, Saejj-Mahalleh village, Amol city, Mazandaran Province, Iran.

Family **Cunaxidae** (1 species)

Armascirus jini Liu & Guo, 2015 *in* Liu, Yi & Guo, 2015: 326—Holotype female (GUGC, GZU-GZ-CU-201308001), from Dushan, Guizhou, China (25°48'26"N, 107°30'47"E).

Superfamily **Eriophyoidea** (42 species)

Family **Eriophyidae** (26 species)

Aceria davidmansi Xue, Han & Zhang, 2015: 82—Holotype female (NZAC), from *Ulex europaeus*, Mahinapua Scenic Reserve, Ruatapu, Westland, New Zealand (42°47'34"S, 170°54'09"E).

Aceria heteropappi Xue & Hong, 2013 *in* Xue, Sadeghi, Hong & Sinaie, 2013: 44—Holotype female (NJAU, IRAN214.1), from *Heteropappus altaicus*, Ferdowsi University campus, Mashhad, Razavi Khorasan Province, Iran (36°18'00"N, 59°30'60"E).

Aculops coluteae Lotfollahi, de Lillo & Haddad, 2015 *in* Lotfollahi, de Lillo & Irani-Nejad, 2015: 416—Holotype female (UTI, CC-IEA-KN11L-1), from *Colutea cilicica*, Turkish bladder senna, Kandovan village, Osku, Iran (37°47'31"N, 46°14'57"E).

Aculops consoregalis Lotfollahi, de Lillo & Haddad, 2015 *in* Lotfollahi, de Lillo & Irani-Nejad, 2015: 413—Holotype female (UTI, CR-IEA-SD11L-1), from *Consolida regalis*, Larkspur, Royal Knight's-spur, Sardrood, Iran (38°02'02"N, 46°08'24"E).

Aculops menthaefoliae Lotfollahi, de Lillo & Haddad, 2015 *in* Lotfollahi, de Lillo & Irani-Nejad, 2015: 414—Holotype female (UTI, ML-IEA-MN11L-1), from *Mentha longifolia*, Mamaghan, Iran (37°50'45"N, 45°59'02"E).

Aculus portulacis Wang & El-halawany, 2014 *in* Wang, El-halawany, Xue & Hong, 2014: 426—Holotype female (NJAU, NJAUAcariEriEgypt7.1), from *Portulaca oleracea*, Al-Imam Mohamed Bin Saud Islamic University, Riyadh Province, Saudi Arabia (24°49'N, 46°42'E).

Cecidophyopsis rosmarinus Wang & El-halawany, 2014 *in* Wang, El-halawany, Xue & Hong, 2014: 413—Holotype female (NJAU, NJAUAcariEriEgypt6.1), from *Rosmarinus officinalis*, Dereiya, Riyadh Province, Saudi Arabia (24°42'N, 46°37'E).

Colopodacus cochinchinensis Zuo, Rajput, Xue & Hong, 2015: 290—Holotype female (NJAU, NJAUAcariEriGD27.1), from *Glycosmis cochinchinensis*, Nanling Mountain, Ruyuan Yao Autonomous County, Guangdong Province, P.R. China (24°54'43"N, 113°03'01"E).

Colopodacus formosus Zuo, Rajput, Xue & Hong, 2015: 288—Holotype female (NJAU, NJAUAcariEriGD2.1), from *Leycesteria formosa*, Nanling Mountain, Ruyuan Yao Autonomous County, Guangdong Province, P.R. China (24°54'43"N, 113°03'01"E).

Epitrimerus saudiarabis Wang & El-halawany, 2014 *in* Wang, El-halawany, Xue & Hong, 2014: 420—Holotype female (NJAU, NJAUAcariEriEgypt2.1), from *Phoenix dactylifera*, Al-Imam Mohamed Bin Saud Islamic University, Riyadh Province, Saudi Arabia (24°49'N, 46°42'E).

***Floracarus ruyuanicus* Zuo, Rajput, Xue & Hong, 2015: 293**—Holotype female (NJAU, NJAUAcariEriGD10A.1), from *Elaeocarpus dubius*, Nanling Mountain, Ruyuan Yao Autonomous County, Guangdong Province, P.R. China (24°54'43"N, 113°03'01"E).

***Tetra ampelopsis* Han, Rajput, Xue & Hong, 2015: 518**—Holotype female (NJAU, NJAUAcariEriHN175.1), from *Ampelopsis* sp., Yuelu Mountain, Changsha City, Hunan Province, P.R. China (28°11'39"N, 112°56'08"E).

***Tetra banqiaoensis* Han, Rajput, Xue & Hong, 2015: 511**—Holotype female (NJAU, NJAUAcariEriYN59.1), from *Populus* sp., Banqiao Town, Baoshan City, Yunnan Province, P.R. China (25°07'22"N, 99°09'08"E).

***Tetra heliotropii* Xue & Hong, 2013 in Xue, Sadeghi, Hong & Sinaie, 2013: 50**—Holotype female (NJAU, IRAN230.1), from *Heliotropium chorassanicum*, Ferdowsi University campus, Mashhad, Razavi Khorasan Province, Iran (36°18'00"N, 59°31'00"E).

***Tetra nitrariae* Li, Li, Zhang & Xue, 2014: 349**—Holotype female (SHZU, Tetra 5.1), from *Nitraria schoheri*, Tacheng city, Xinjiang Uygur Autonomous Region, P.R. China (44°32'21"N, 83°56'32"E).

***Tetra phragmites* Rajput, Zuo, Wang & Hong, 2015: 210**—Holotype female (NJAU, NJAUAcariEriGD33B.1), from *Phragmites australis*, Nanling mountain, Ruyuan Yao Autonomous County, Guangdong Province, P.R. China (24°53'50"N, 113°01'18"E).

***Tetra prunusis* Rajput, Zuo, Wang & Hong, 2015: 213**—Holotype female (NJAU, NJAUAcariEriZ166B.1), from *Prunus* sp., Tianmu mountain, Lin'an City, Zhejiang Province, P.R. China (30°20'37"N, 119°26'24"E).

***Tetra qingyuanensis* Rajput, Zuo, Wang & Hong, 2015: 206**—Holotype female (NJAU, NJAUAcariEriZ23B.1), from *Lespedeza* sp., Qingyuan county, Lishui City, Zhejiang Province, P.R. China (27°34'51"N, 119°4'98"E).

***Tetra sativae* Li, Li, Zhang & Xue, 2014: 343**—Holotype female (SHZU, Tetra 3.1), from *Medicago sativa*, Xinjiang Uygur Autonomous Region, Fukang county, P.R. China (43°54'04"N, 88°07'03"E).

***Tetra sojae* Rajput, Zuo, Wang & Hong, 2015: 215**—Holotype female (NJAU, NJAUAcariEriW75.1), from *Glycine soja*, Gunuijiang, Shitai County, Anhui Province, P.R. China (30°08'28"N, 117°47'17"E).

***Tetraspinus lucida* Han, Rajput, Xue & Hong, 2015: 519**—Holotype female (NJAU, NJAUAcariEriYN253.1), from *Ligustrum lucidum*, Fugong County, Nujiang Lisu autonomous prefecture, Yunnan Province, P.R. China (26°54'06"N, 98°51'31"E).

***Tetra tianchiensis* Li, Li, Zhang & Xue, 2014: 337**—Holotype female (SHZU, SHZUTetra 2.1), from *Sorbus tianschanica*, Xinjiang Uygur Autonomous Region, Fukang city, P.R. China (43°53'28"N, 88°07'11"E).

***Tetra tianschanicae* Li, Li, Zhang & Xue, 2014: 334**—Holotype female (SHZU, Tetra 1.1), from *Sorbus tianschanica*, Xinjiang Uygur Autonomous Region, Xinyuan county, P.R. China (43°15'26"N, 84°36'18"E).

***Tetra viciae* Li, Li, Zhang & Xue, 2014: 344**—Holotype female (SHZU, Tetra 4.1), from *Vicia ceacca*, Xinjiang Uygur Autonomous Region, Buerjin county, China (48°38'15"N, 87°02'43"E).

***Tetra virga* Han, Rajput, Xue & Hong, 2015: 512**—Holotype female (NJAU, NJAUAcariEriYN278.1), from *Rhododendron irroratum*, Fugong County, Nujiang Lisu autonomous prefecture, Yunnan Province, P.R. China (26°33'05"N, 98°55'08"E).

***Tetra vitexus* Rajput, Zuo, Wang & Hong, 2015: 217**—Holotype female (NJAU, NJAUAcariEriZ156C.1), from *Vitex negundo* var. *cannabifolia*, Tianmu mountain, Lin'an City, Zhejiang Province, P.R. China (30°20'37"N, 119°26'24"E).

Family Diptilomiopidae (16 species)

***Acarhis kleinhoviae* Tan, Yang, Lai & Wang, 2015: 771**—Holotype female (GUGX, GXUEDA1-1.1), from *Kleinhovia hospita*, Baisha County, Hainan Province, P.R. China (109°19'05"E, 18°62'14"N).

***Cheiracus costus* Tan, Yang, Lai & Wang, 2015: 778**—Holotype female (GUGX, GXUEDC2-3.1), from *Costus speciosus*, Baisha County, Hainan Province, P.R. China (109°20'11"E, 18°35'17"N).

***Diptacus elegantulum* Rajput, Han, Xue & Hong, 2014: 231**—Holotype female (NJAU, NJAUAcariEriZ232.1), from *Acer elegantulum*, Qingliang mountain, Lin'an City, Zhejiang Province, P.R. China (30°02'26"N, 118°56'04"E).

- Diptacus tianmuensis Rajput, Han, Xue & Hong, 2014:** 234—Holotype female (NJAU, NJAUAcariEriZ166A.1), from *Prunus* sp., Tianmu mountain, Lin'an City, Zhejiang Province, P.R. China (30°20'37"N, 119°26'24"E).
- Diptilomiopus engelhardter Xue, Cheng & Hong, 2013:** 183—Holotype female (NJAU, NJAUAcariErQ210.1), from *Engelhardtia roxburghiana*, Diaoluo Mountain, Hainan Province, P.R. China (18°43'30"N, 109°52'06"E).
- Diptilomiopus diaoluicus Xue, Cheng & Hong, 2013:** 178—Holotype female (NJAU, NJAUAcariErQ228.1), from *Eurya hainanensis*, Diaoluo Mountain, Hainan Province, P.R. China (18°43'30"N, 109°52'06"E).
- Diptilomiopus khatkhaii Konvipasruang & Chandrapatya, 2015:** 310—Holotype female (KUB, slide no. 1140), from *Oxyceros horridus*, Chaam District, Phetchaburi Province, Thailand (12°47'59"N, 99°58'1"E).
- Diptilomiopus mandongii Konvipasruang & Chandrapatya, 2015:** 307—Holotype female (KUB, slide no. 561), from *Cordia dichotoma*, Mueang Tak District, Tak Province, Thailand (16°52'54"N, 99°07'25"E).
- Khamroka tomentosi Konvipasruang & Chandrapatya, 2015:** 299—Holotype female (KUB, slide no. 1114), from *Ellipanthus tomentosus*, Cha-am District, Phetchaburi Province, Thailand (12°47'59"N, 99°58'1"E).
- Konola pingis Xue, Cheng & Hong, 2013:** 188—Holotype female (NJAU, NJAUAcariErQ198.1), from *Machilus pingii*, Diaoluo Mountain, Hainan Province, P.R. China (18°43'30"N, 109°52'06"E).
- Linacutus cathayensis Rajput, Han, Xue & Hong, 2014:** 228—Holotype female (NJAU, NJAUAcariEriZ214.1), from *Carya cathayensis*, Qingliang mountain, Lin'an City, Zhejiang Province, P.R. China (30°02'26"N, 118°56'04"E).
- Neorhynacus altingus Xue, Cheng & Hong, 2013:** 185—Holotype female (NJAU, NJAUAcariErQ213.1), from *Altingia obovata*, Diaoluo Mountain, Hainan Province, P.R. China (18°43'30"N, 109°52'06"E).
- Neorhynacus buettnerus Tan, Yang, Lai & Wang, 2015:** 773—Holotype female (GUGX, GXUEDN3-1.1), from *Buettneria aspera*, Baisha County, Hainan Province, P.R. China (109°19'05"E, 18°62'14"N).
- Rhynacus aceroides Flechtmann, 2014 in Flechtmann, Ballari & De Quinteros, 2014:** 76—Holotype female (ESALQ), from *Solanum lorentzii* leaves, Finca Experimental La Posta, Perico, Jujuy Province, Argentina (24°22'51"S, 65°08'22"W).
- Salaya takhronami Konvipasruang & Chandrapatya, 2015:** 304—Holotype female (KUB, slide no. 588), from *Sisyrolepis muricata*, Salaya Sub-district, Phutthamonthon District, Nakhon Pathom Province, Thailand (13°47'36"N, 100°19'21"E).
- Vimola scutellata Tan, Yang, Lai & Wang, 2015:** 776—Holotype female (GUGX, GXUEDV1-4.1), from *Memecylon scutellatum*, Bawangling Nature Reserve, Changjing County, Hainan Province, P.R. China (109°19'02"E, 19°06'12"N).

Superfamily Eupodoidea (1 species)

Family Eupodidae (1 species)

Pseudopenthaleus tauricus Khaustov, 2015b: 934—Holotype female (TUMZ, slide No. AK170394), from moss on soil on the meadow surrounded by Crimean Pines *Pinus pallasiana*, Yalta mountain-forest Nature Reserve, Crimea, Ukraine (44°31'01.5"N, 34°08'33.3"E).

Superfamily Tydeoidea (1 species)

Family Tydeidae (1 species)

Brachytydeus persiaensis Akbari, Haddad & Khanjani, 2015 in Akbari, Irani-Nejad, Khanjani, Arzanlou & Kaźmierski, 2015a: 426—Holotype female (UTI), from the soil under peach trees at vicinity of Shabestar, East Azerbaijan, Iran.

Infraorder Anystina (17 species)

Hypoder Parasitengona (17 species)

Superfamily Calyptostomatoidea (2 species)

Family Calyptostomatidae (2 species)

Calyptostoma giuliae Haitlinger & Šundić, 2015: 921—Holotype larva (MNHWU), from unidentified herbaceous plants, in Bantosey, 20 km north of Siem Reap, Cambodia.

Calyptostoma marantica Haitlinger & Šundić, 2015: 925—Holotype larva (MNHWU), from unidentified herbaceous plants, in Marante, Sulawesi, Indonesia.

Superfamily **Erythraeoidea** (5 species)

Family **Erythraeidae** (5 species)

Charletonia teriana **Hakimitabar, Saboori & Seiedy, 2013:** 169—Holotype larva (JAZM, ARS-20120710-1a), ectoparasitic on Phalangiidae, bark of trees in the botanical garden, Faculty of Agriculture, University of Tehran, Karaj, Alborz Province, Iran (35°48.189'N, 50°59.523'E).

Erythraeus (Zaracarus) hafezi **Saboori, Hakimitabar & Mahmoudi, 2014 in Mahmoudi, Saboori, Sari & Hakimitabar, 2014:** 80—Holotype larva (JAZM, ARS-20120208-1a), ectoparasitic on an unidentified Cicadidae, Shiraz city, Fars Province, Iran.

Moldoustium haitlingeri **Noei, Saboori & Šundić, 2013 in Noei, Saboori, Šundić, Hajizadeh & Pešić, 2013:** 269—Holotype larva (JAZM, ARS-20130126-1a), from soil, Loshan city, Guilan Province, Iran (36°35'N, 49°32'E).

Momorangia binaloudensis **Noei & Saboori, 2015 in Noei, Rabieh & Saboori, 2015:** 793—Holotype lava (JAZM, ARS-20150306-1a), ectoparasitic on a male of *Apamea impedita*, Shirbad, Binaloud Mountain, Khorasan Razavi Province, Iran (36°16'00"N, 59°04'18"E).

Nagoricanella salehi **Kamran & Alatawi, 2015:** 199—Holotype larva (KSU, accession no. ER: 14/30-35), from African foxtail grass, *Cenchrus ciliaris*, Rudat Al-Khuraim, Riyadh, Saudi Arabia (25°21.251'N, 47°17.837'E).

Superfamily **Trombidioidea** (6 species)

Family **Microtrombidiidae** (4 species)

Atractothrombium amirkabiri **Noei, Saboori & Hajizadeh, 2015b:** 658—Holotype larva (JAZM, ARS-20141105-1a), from soil in a forest close to a stream, Gijave village, Asalem city, Guilan Province, Iran (37°42.077'N, 48°54.077'E).

Ettmuelleria zahediae **Noei, Saboori & Hajizadeh, 2015b:** 662—Holotype larva (JAZM, ARS-20141105-2a), from soil in a forest close to a stream, Roudsar city, Guilan Province, Iran (37°01.647'N, 50°15.376'E).

Microtrombidium fumani **Noei, Saboori & Hajizadeh, 2015b:** 665—Holotype larva (JAZM, ARS-20141105-3a), from soil in a forest close to a stream, Masouleh city, Guilan Province, Iran (37°08.808'N, 48°59.254'E).

Milandanielia harsitensis **Adil & Sevsay, 2015 in Adil, Sevsay, Doğan & Dilkaraoğlu, 2015:** 651—Holotype adult (EUE), from harşit valley, Giresun Province, Turkey (40°46'45"N, 38°59'36"E).

Family **Trombidiidae** (2 species)

Paratrombium farnazae **Noei, Saboori & Hajizadeh, 2015a:** 316—Holotype lava (JAZM, ARS-20130517-1a), from soil in a forest, in Noghlebar village, Roodbar city, Iran (36°57.39'N, 49°32.65'E).

Podothrombium zlatarum **Saboori, Pešić & Šundić, 2015:** 122—Holotype larva (JAZM, ARS-20140731-1a), from herbaceous plants, in Zlatar Mountain, near Nova Varoš, Serbia.

Superfamily **Hydryphantoidea** (1 species)

Family **Hydryphantidae** (1 species)

Trichothyas anatolica **Boyaci & Güller, 2014:** 161—Holotype female (FFSDU), from a shallow spring with bottom completely covered by algae, Şarkıkaraağaç, Isparta, Turkey (38°7.281'N, 31°26.860'E).

Superfamily **Hygrobatoidea** (2 species)

Family **Hygrobatidae** (2 species)

Atractides asiaticus **Pešić & Smit, 2015a:** 784—Holotype male (RMNH, ZMA.ACA.P 32808), from stream near Ban Khoom, Doi Ang Kang, Thailand (19°56.905'N, 99°03.938'E).

Atractides inthanonensis **Pešić & Smit, 2015a:** 785—Holotype male (RMNH), from stream downstream of Siriphum Waterfall, Doi Inthanon NP, Thailand (18°31.532'N, 98°39.091'E).

Superfamily **Arrenuroidea** (1 species)

Family **Arrenuridae** (1 species)

Thoracophoracarus koreanus **Smit & Pešić, 2014:** 432—Holotype male (NIBR), from pool near stream, 1.7 km from

Jiksopolpol falls, 0.6 km from Naebyeansan Info Center (near Silsung temple), Naebyeansan National Park, Ne myeon Mt, CR10, South Korea (35°64'65.978"N, 126°58'62.285"E).

Infraorder **Eleutherengona** (32 species)

Hyporder **Raphignathina** (23 species)

Superfamily **Raphignathoidea** (11 species)

Family **Caligonellidae** (1 species)

Neognathus sinaei Khanjani & Samadpour, 2014 in Samadpour, Khanjani & Asalifayaz, 2014: 152—Holotype female (BASU), from soil and litter under Russian olive trees, *Elaeagnus angustifolia*, Mallayer vicinity, Hamedan Province, Iran.

Family **Cryptognathidae** (1 species)

Favognathus farshchiani Jafari, Bagheri & Saboori, 2014: 357—Holotype female (JAZM, ARS-20140613-1a), from soil under berry, apricot and elm trees in various orchards and green spaces around Isfahan, Isfahan Province, Iran.

Family **Eupalopsellidae** (1 species)

Eupalopsellus kermaniensis Mahdavi & Farzan, 2013 in Mahdavi, Asadi & Farzan, 2013: 191—Holotype female (SBUC), from *Artemisia* sp., Bongan, Baft, Kerman Province, Iran.

Family **Stigmaeidae** (8 species)

Cheylostigmaeus mahvashae Khanjani, Firoozfar & Mirmoayedi, 2013: 349—Holotype male (BASU), from soil under canopy of apricot trees, *Prunus armeniaca*, Rijab valley, Sarpolzohab, Kermanshah Province, Iran (34°9'–34°46"E, 45°53'–46°35"N).

Cheylostigmaeus variatus Doğan, Dilkaraoglu & Fan, 2015 in Doğan, Dilkaraoglu, Fan, Sevsay, Erman & Adil, 2015: 804—Holotype male (EUE), from hydrous grassy soil, Ekşisu Marsh, Turkey (39°43'52.9"N, 39°37'26.9"E).

Eryngiopus anatolicus Doğan, Dilkaraoglu & Fan, 2015 in Doğan, Dilkaraoglu, Fan, Erman, Sevsay & Adil, 2015: 438—Holotype female (EUE), from litter under *Astragalus* sp., Vauk highway crossing, Gümüşhane, Turkey (40°22'01"N 39°49'20"E).

Eustigmaeus ueckermannii Bagheri, 2013 in Bagheri & Beyzavi, 2013: 33—Holotype female (ACIAU), from the soil and humus under oak trees *Quercus brantii*, south of Kamfiruz region, Fars Province, Iran.

Paraneognathus iranicus Bagheri & Paktinat-Saejj, 2015 in Bagheri, Paktinat-Saejj, Saboori & Ahaniazad, 2015: 335—Holotype female (AFUMI), from the soiland humus under citrus trees, Bahnamir region, Babolsar city, Mazandaran Province, Iran.

Stigmaeus tolstikovi Khaustov, 2015c: 684—Holotype female (TUMZ, slide No. AK151114S), from rotten log of poplar (*Populus nigra*), Tyumen, Tyumen Province, Russia (57°09'52.3"N, 65°31'01.6"E).

Storchia elhamae Hassanzadeh, Khanjani, Safaralizadeh & Mirfakhraie, 2013: 354—Holotype female (BASU), from soil covered with grass, Mahabad vicinity, western Azerbaijan Province, Iran (36°51'12"N, 45°17'13"E).

Zetzellia camphorae Negm, Johann, Ferla & Amano, 2015: 964—Holotype female (KUM), from *Cinnamomum camphora*, Kitashirakawaoiwake-cho, Kyoto-shi, Kyoto prefecture, Japan (28°34'N, 83°98'E).

Superfamily **Tetranychoidae** (12 species)

Family **Tenuipalpidae** (8 species)

Acaricis urigersoni Xu & Zhang, 2013: 358—Holotype female (NZAC), from *Gahnia lacera*, Kepa Reserve, Auckland, New Zealand.

Aegyptobia kermaniensis Farzan, 2015 in Farzan & Asadi, 2015: 956—Holotype female (SBUC), from *Berberis vulgaris*, Kuhpayeh, Kerman-Kerman Province, Iran (28°58'N, 57°37'E).

Aegyptobia khanjanii Farzan & Asadi, 2013: 36—Holotype female (SBUC), from *Astragalus* sp., Jabalbarez, Jiroft town, Kerman Province, Iran (27°21'N, 57°42'E).

Dolichotetranychus riyadhensis Alatawi & Kamran, 2015 in Alatawi, Kamran & Negm, 2015: 826—Holotype female (KSU), from *Cynodon dactylon*, Raudat Khuraim, Riyadh, Saudi Arabia (25°21.251'N, 47°17.837'E).

Krugeria fuzhouensis Xu & Fan, 2015 *in* Xu, Fan & Huang, 2015: 88—Holotype female (NZMC), from *Pterospermum heterophyllum*, Shangxiadian Road, Fujian Agriculture and Forestry University, Cangshan district, Fuzhou city, Fujian Province, China (119°14'2.89"E, 26°5'14.98"N).

Phytopipalpus phoenicis Alatawi & Kamran, 2015 *in* Alatawi, Kamran & Negm, 2015: 817—Holotype female (KSU), from leaves and fiber of *Phoenix dactylifera*, Al-Imam Muhammad bin Saud Islamic University, Riyadh Province, Saudi Arabia (24°48'77"N, 46°42'73"E).

Tenuipalpus toropi Castro, Ramos & Feres, 2015 *in* De Castro, Ramos, Feres & Ochoa, 2015: 340—Holotype female (PEU, n. 9540), from *Sapium glandulatum*, Paulo de Faria, São Paulo, Brazil (19°55'S, 49°31'W).

Tenuipalpus zahiri Khanjani & Seeman, 2013 *in* Khanjani, Alvandy, Khanjani & Seeman, 2013: 392—Holotype female (CALBS), from hawthorn leaves, *Crataegus monogyna*, Simin, Hamedan Province, Iran (34°41'N, 48°35'E).

Family Tetranychidae (4 species)

Mononychellus cedrelae Flechtmann & Queiroz, 2015: 834—Holotype female (MZLQ), from the underside of bronzed leaves of *Cedrela fissilis*, Campus EMBRAPA, Colombo, Paraná, Brazil (25°16'06"S, 9°32'0"W).

Schizotetranychus iraniensis Mahdavi & Asadi, 2015a: 675—Holotype male (SBUC), from *Prunus cerasus*, Khabr-Kerman Province, Iran (28°48'N, 56°20'E).

Tetranychus iraniensis Mahdavi & Ueckermann, 2013 *in* Mahdavi, Asadi, Ueckermann & Farzan, 2013: 246—Holotype female (SBUC), from *Salsola* sp., Manujan-Kerman Province, Iran (27°19'N, 57°29'E).

Tetranychopsis iranensis Khanjani, Khanjani & Razmjou, 2015: 119—Holotype female (BASU), from hazelnut trees, *Corulus avellana*, Fondoghlo National Park Jungle, Ardabil, Ardabil Province, Iran (38°26'E, 48°34'N).

Hyporder Heterostigmata (9 species)

Superfamily Dolichocyboidea (1 species)

Family Dolichocybidae (1 species)

Dolichomotes sinuspersicus Mortazavi & Hajiqanbar, 2015 *in* Mortazavi, Hajiqanbar & Kamali, 2015: 446—Holotype female (TMU, AM20130314), from the elytra of a bostrichid beetle *Sinoxylon pugnax*, vicinity of Minab city, Hormozgan Province, Iran (27°19'09"N, 57°00'15"E).

Superfamily Tarsonemoidea (8 species)

Family Podapolipidae (8 species)

Cassidopolipus physonota Husband & OConnor, 2014: 440—Holotype female (UMMZ), from the wild olive tortoise beetle, *Physonota alutacea*, 2.41km SE El Cameron, 610 m Oaxaca, Mexico (15°53'34"N, 96°58'33"E).

Eutarsopolipus americanus Husband & Husband, 2013: 57—Holotype female (UMMZ, RWH200309-1), from *Chlaenius praesinus*, Silver Lake State Park, Oceana County, Michigan, USA.

Eutarsopolipus asiaticus Husband & Kurosa, 2013: 67—Holotype female (NSMT, Kurosa No. 21453-5), from *Chlaenius costiger*, Sato, Mikura-jima Is., Izu Iss., Tokyo, Japan.

Eutarsopolipus earnshawi Constantine & Seeman, 2014: 89—Holotype female (QM, # IQ-1000-D), from *Cratoferonia phylarchus*, southeast Queensland, Lamington National Park, rainforest, Australia (28°15'43"S, 153°10'12"E).

Eutarsopolipus lambkinae Constantine & Seeman, 2014: 96—Holotype female (QM), from *Notonomus angustibasis*, southeast Queensland, Lamington National Park, rainforest, Australia (28°14'17"S, 153°08'42"E).

Eutarsopolipus obrieni Husband & Husband, 2015: 972—Holotype female (UMMZ, RWH220310-10), from *Platynus darlingtoni*, vicinity of Cinchona, St. Andrew Parish, Jamaica, British West Indies.

Eutarsopolipus rutherfordae Constantine & Seeman, 2014: 104—Holotype female (QM), from *Trichosternus subvirens*, southeast Queensland, Australia.

Simalurapolipus hiraii Kurosa & Husband, 2013: 260—Holotype female (NSMT, 4940-8), from *Simaluracoerulea*, Hirayama, Aoi-ku, Shizuoka-shi, Shizuoka Prefecture, Japan.

List of type localities and depositories of new species of the Order Trombidiformes in ZOOTAXA 2013–2015

Superorder **Acariformes** (396 species)
Order **Trombidiformes** (396 species)
Suborder **Prostigmata** (396 species)
Infraorder **Eupodina** (67 species)
Superfamily **Bdelloidea** (12 species)
Family **Bdellidae** (4 species)

***Bdella farabii* Paktnat-Saej & Bagheri, 2015** in **Paktnat-Saej, Bagheri, Saboori & Ahaniazad, 2015: 526**—Holotype female (UMAC), from soil and rotten leaves under citrus trees, Saejj-Mahalleh village, Amol city, Mazandaran Province, Iran.

***Biscirus iranensis* Paktnat-Saej & Bagheri, 2015** in **Paktnat-Saej, Bagheri, Saboori & Ahaniazad, 2015: 523**—Holotype male (UMAC), from the soil and rotten leaves under citrus trees, Amreh village, Sari city, Mazandaran Province, Iran.

***Cyta kurdistanicus* Eghbalian, Khanjani & Ueckermann, 2014** in **Eghbalian, Khanjani, Safaralizadeh & Ueckermann, 2014: 575**—Holotype female (BASU), from soil covered with grass and litter under wild almond, *Amygdalus lycioides*, Sirvan river, Palangan village, Kurdistan Province, Iran (35°03.7'N, 46°35.97'E).

***Cyta leliae* Eghbalian, Khanjani & Ueckermann, 2014** in **Eghbalian, Khanjani, Safaralizadeh & Ueckermann, 2014: 571**—Holotype male (BASU), from soil and litter under grass and oak trees, *Quercus brantii*, “Sirvan river bank”, Palangan village, Kurdistan Province, Iran (35°03.7'N, 46°35.97'E).

Family **Cunaxidae** (8 species)

***Armascirus denheyeri* Kalúz, Ermilov & Vrabec, 2014: 243**—Holotype female (SNMB), from Hunli env., Arunachal Pradesh, India (28°19'32"N, 95°57'31"E).

***Armascirus fendai* Kalúz & Vrabec, 2013: 144**—Holotype female (SNMB), from soil with grass roots on sands, Nature Reserve Abrod—*Molinietum caeruleae*, Veľké Leváre village vicinity, Slovakia (48°31'57"N, 17°00'21"E).

***Armascirus masani* Kalúz & Vrabec, 2013: 150**—Holotype male (SNMB), from soil with grass roots, mountain meadow, Nová Polianka village vicinity, High Tatras Mts., Slovakia (49°07'17"N, 20°09'46"E).

***Armascirus orientalis* Kalúz, Ermilov & Vrabec, 2014: 248**—Holotype female (SNMB), from samples of bird's nest fungi *Cyathus* sp., lowland Lagerstroemia-dominated forest on dark loamy soil, Dong Nai Biosphere Reserve, Vietnam (11°25'N, 107°25'E).

***Bonzia flechtmanni* Rocha, Rodrigues & Ferla, 2015: 62**—Holotype female (ESALQ), from the leaf litter on Atlantic Araucaria forest, Centro de Pesquisas e Conservação da Natureza Pró-Mata, São Francisco de Paula, Rio Grande do Sul, Brazil (29°29'23.70"S, 50°12'16.30"W).

***Dactyloscirus multiscutus* Rocha, Rodrigues & Ferla, 2015: 68**—Holotype female (ESALQ), from the leaf litter on Atlantic rainforest, Centro de Pesquisas e Conservação da Natureza Pró-Mata, São Francisco de Paula, Rio Grande do Sul, Brazil (29°28'31.40"S, 50°9'58.30"W).

***Neocunaxoides promatae* Rocha, Rodrigues & Ferla, 2015: 60**—Holotype female (ESALQ), from the leaf litter on Atlantic Araucaria forest, Centro de Pesquisas e Conservação da Natureza Pró-Mata São Francisco de Paula, Rio Grande do Sul, Brazil (29°29'4.20"S, 50°13'15.16"W).

***Scutopalus tomentosus* Rocha, Skvarla & Ferla, 2013: 40**—Holotype female (ESALQ/USP), from *Plantago tomentosa*, Dois Lajeados, Rio Grande do Sul State, Brazil.

Superfamily **Eriophyoidea** (39 species)

Family **Phytopidae** (5 species)

***Mackiella reclinata* Chetverikov & Craemer, 2014** *in* Chetverikov, Craemer, Vishnyakov & Sukhareva, 2014: 271—
Holotype female (NCA-PPRI, slide #76-13), from *Phoenix reclinata*, National Botanical Garden, Pretoria, South Africa
(25°44'18.92"S, 28°28'16.98"E).

***Novophytoptus luzulis* Chetverikov, 2015: 493**—Holotype female (ZISP, #24-13), from *Luzula pilosa*, Plesetsk, Arkhangelsk Prov., Russia (62°43'51"N, 40°17'5"E).

***Novophytoptus maritimus* Chetverikov, 2015: 497**—Holotype female (ZISP, #u1/2-12), from *Juncus maritimus*, Kirillovka, Zaporozhye Prov., Ukraine (46°18'32"N, 35°18'30"E).

***Paraphytoptus intybi* Denizhan, Monfreda, Lillo & Cobanoğlu, 2015: 33**—Holotype female (DiSSPA, #3008), from *Cichorium intybus*, Eğmir lake, Ankara, Turkey (39°49'29"N 32°50'05"E).

***Phytoptus albae* Denizhan, Monfreda, Lillo & Cobanoğlu, 2015: 7**—Holotype female (DiSSPA, #2764), from *Populus alba*, Çamlıdere, Turkey (40°31'29"N, 32°29'24"E).

Family Eriophyidae (27 species)

***Aceria lobolinae* Lotfollahi & de Lillo, 2014** *in* Lotfollahi, Irani-Nejad & de Lillo, 2014: 81—Holotype female (UTI, EA-IEA-AR11L-1), from *Elaeagnus angustifolia*, Azarshahr, Iran (37°46'24"N, 45°57'20"E).

***Aceria tripuraensis* Menon, Joshi & Ramamurthy, 2014: 560**—Holotype female (IARI), from *Hibiscus macrophyllus*, Ishaan Chandranagar, Agartala, Tripura, India (23°45'55"N, 91°14'33"E).

***Aceria utzkae* Vidović, 2014: 489**—Holotype female (UBS and NHMB*, slide # 204/8), from *Carlina acanthifolia*, Goč, Serbia (43°33.519"N, 20°44.243"E).

***Aceria virgatae* Lotfollahi, Irani-Nejad, Khanjani, Moghadam & De Lillo, 2013: 354**—Holotype female (UTI, CV-IEA-MH10L-1), from *Centaurea virgata*, Maragheh countryside, Iran (37°25'35"N, 46°14'38"E).

***Aceria xeranthemis* Lotfollahi, Irani-Nejad, Khanjani, Moghadam & De Lillo, 2013: 354**—Holotype female (UTI, XS-IEA-TZ11L-1), from *Xeranthemum squarrosum*, Tabriz countryside, Iran (38°01'13"N, 46°25'09"E).

***Aculops huzhongensis* Xue, Guo & Hong, 2013: 77**—Holotype female (NJAU, NJUAcariEriH30-1), from *Salix* sp., Huzhong District, Heilongjiang Province, P.R. China (52°01'44"N, 123°35'44"E).

***Aculus castriferrei* Ripka, 2014: 265**—Holotype female (NFCSO, slide # 1315a), from *Agrimonia eupatoria*, Mesteri, Vas County, Hungary (47°14'6.155"N, 17°5'15.806"E).

***Aculus huzhongsalixus* Xue, Guo & Hong, 2013: 83**—Holotype female (NJAU, NJUAcariEriH30-2), from *Salix* sp., Huzhong District, Heilongjiang Province, P.R. China (52°01'44"N, 123°35'44"E).

***Calepitrimerus flexuosus* Xue, Guo & Hong, 2013: 37**—Holotype female (NJAU, NJUAcariEriH52), from *Spiraea flexuosa*, Yichun City, Heilongjiang Province, P.R. China (47°43'36"N, 128°55'12"E).

***Calepitrimerus maximowiczii* Xue, Guo & Hong, 2013: 40**—Holotype female (NJAU, NJUAcariEriH34), from *Crataegus maximowiczii*, Huzhong District, Heilongjiang Province, P.R. China (52°01'44"N, 123°35'44"E).

***Calepitrimerus pilosus* Xue, Guo & Hong, 2013: 42**—Holotype female (NJAU, NJUAcariEriH22), from *Agrimonia pilosa*, Jiagedaqi, Heilongjiang Province, P.R. China (50°24'27"N, 124°06'43"E).

***Calepitrimerus yichunensis* Xue, Guo & Hong, 2013: 49**—Holotype female (NJAU, NJUAcariEriH62), from *Sorbaria sorbifolia*, Harbin City, Heilongjiang Province, P.R. China.

***Cupacarus oxyphyllus* Xue, Guo & Hong, 2013: 50**—Holotype female (NJAU, NJUAcariEriH49), from *Euonymus oxyphyllus*, Yichun City, Heilongjiang Province, P.R. China (47°43'36"N, 128°55'12"E).

***Eriocaenus ramosissimi* Petanović & Amrine, 2015** *in* Petanović, Amrine, Chetverikov & Cvrković, 2015: 58—Holotype female (UBS, slide #1431/1), from *Equisetum ramosissimum*, Zemun, Altina, Serbia (44°50'26.3"N, 20°21'34.8"E).

***Epitrimerus sambucus* Xue, Guo & Hong, 2013: 54**—Holotype female (NJAU, NJUAcariEriH70), from *Sambucus williamsii*, Harbin City, Heilongjiang Province, P.R. China.

Epitrimerus wuyingensis Xue, Guo & Hong, 2013: 59—Holotype female (NJAU, NJAUAcariEriH47-2), from Acer sp., Wuying, Yichun City, Heilongjiang Province, P.R. China (47°43'36"N, 128°55'12"E).

Longisolenidionus amurensis Xue, Guo & Hong, 2013: 68—Holotype female (NJAU, NJAUAcariEriH54), from *Tilia amurensis*, Yichun City, Heilongjiang Province, P.R. China (47°43'36"N, 128°55'12"E).

Phyllocoptes jiagedaqiensis Xue, Guo & Hong, 2013: 73—Holotype female (NJAU, NJAUAcariEriH7), from *Cunninghamia* sp., Jiagedaqi, Heilongjiang Province, P.R. China (50°24'27"N, 124°06'43"E).

Phyllocoptruta juniperiana Han, Xue & Hong, 2013: 260—Holotype female (NJAU, NJAUAcariEriYN69.1), from *Juniperus chinensis*, Cangshan, Dali City, Yunnan Province, P.R. China (25°41'47"N, 100°08'28"E).

Proiectus granularpro Han, Xue & Hong, 2013: 253—Holotype female (NJAU, NJAUAcariEriYN61.1), from *Pinus tabuliformis*, Erhai Park, Dali City, Yunnan Province, P.R. China (25°41'47"N, 100°08'28"E).

Proiectus rodgeseta Han, Xue & Hong, 2013: 249—Holotype female (NJAU, NJAUAcariEriYN68.1), from *Pinus armandii*, Cangshan, Dali City, Yunnan Province, P.R. China (25°41'47"N, 100°08'28"E).

Shevtchenkella huzhongiensis Xue, Guo & Hong, 2013: 26—Holotype female (NJAU, NJAUAcariEriH33), from *Ulmus davidiana*, Huzhong District, Heilongjiang Province, P.R. China (52°01'44"N, 123°35'44"E).

Shevtchenkella jingboicus Xue, Guo & Hong, 2013: 30—Holotype female (NJAU, NJAUAcariEriH47-1), from Acer sp., Yichun City, Heilongjiang Province, P.R. China (47°43'36"N, 128°55'12"E).

Tetra angelica Xue, Guo & Hong, 2013: 94—Holotype female (NJAU, NJAUAcariEriH66), from *Angelica* sp., Jiagedaqi, Heilongjiang Province, P.R. China (50°24'27"N, 124°06'43"E).

Tetra jiagedaqia Xue, Guo & Hong, 2013: 100—Holotype female (NJAU, NJAUAcariEriH16), from *Lespedeza* sp., Jiagedaqi, Heilongjiang Province, P.R. China (50°24'27"N, 124°06'43"E).

Vittacus cannabis Xue, Guo & Hong, 2013: 107—Holotype female (NJAU, NJAUAcariEriJ7), from *Cannabis sativa*, Baicheng City, Jilin Province, P.R. China.

Vittacus mandshurica Xue, Guo & Hong, 2013: 106—Holotype female (NJAU, NJAUAcariEriH23), from *Corylus sieboldiana*, Jiagedaqi, Heilongjiang Province, P.R. China (50°24'27"N, 124°06'43"E).

Family Diptilomiopidae (7 species)

Diptacus coreanus Rajput, Li, Xue & Hong, 2015: 340—Holotype female (NJAU, NJAUAcariEriZ163.1), from *Litsea coreana*, from Tianmu mountain, Lin'an City, Zhejiang Province, P.R. China (30°20'37"N, 119°26'24"E).

Neorhynacus camphoratus Rajput, Li, Xue & Hong, 2015: 342—Holotype female (NJAU, NJAUAcariEriZ198A), from *Cinnamomum camphora*, from Tianmu mountain, Lin'an City, Zhejiang Province, P.R. China (30°22'07"N, 119°26'04"E).

Peralox dentatis Xue, Guo & Hong, 2013: 113—Holotype female (NJAU, NJAUAcariEriH63-1), from *Ulmus* sp., Harbin City, Heilongjiang Province, P.R. China.

Rhinophytoptus nemalobos Lotfollahi & de Lillo, 2014 in Lotfollahi, Irani-Nejad & de Lillo, 2014: 79—Holotype female (UTI, PD-IEA-KN11L-1), from *Prunus domestica*, Kandovan village, Osku, Iran (37°47'31"N, 46°14'57"E).

Rhinotergum boehmerius Rajput, Li, Xue & Hong, 2015: 345—Holotype female (NJAU, NJAUAcariEriZ185.1), from *Boehmeria gracilis*, from Tianmu mountain, Lin'an City, Zhejiang Province, P.R. China (30°22'07"N, 119°26'04"E).

Rhyncaphytoptus cathayensis Rajput, Li, Xue & Hong, 2015: 348—Holotype female (NJAU, NJAUAcariEriZ213A.1), from *Carya cathayensis*, from Qingliang mountain, Lin'an City, Zhejiang Province, P.R. China (30°02'26"N, 118°56'04"E).

Rhyncaphytoptus lanceolatus Rajput, Li, Xue & Hong, 2015: 352—Holotype female (NJAU, NJAUAcariEriZ148.1), from *Cunninghamia lanceolata*, from Tianmu mountain, Lin'an City, Zhejiang Province, P.R. China (30°2'37"N, 119°26'24"E).

Superfamily Halacaroidea (4 species)

Family Halacaridae (4 species)

Halacarus omului Pepato & Da Silveira, 2013: 592—Holotype female (UFMG-AC1200163), from *Sargassum cymosum*, São Paulo state, Lázaro Beach, Ubatuba, Brazil.

***Halacarus todaroi* Pepato & Da Silveira, 2013: 597**—Holotype female (UFMG-AC1200178), from shell debris, Itassucê Islet, Barequeçaba Beach, Brazil (23°50'S, 45°27'W).

***Rhombognathus abirus* Pepato & Da Silveira, 2015: 501**—Holotype female (UFMG-AC1200653), from algae associated to Rhizophora mangle, Fazenda River, municipality of Ubatuba, São Paulo State, Brazil (23°21'55.5"S, 44°50'18.6"W).

***Rhombognathus picinguabensis* Pepato & Da Silveira, 2015: 507**—Holotype female (UFMG-AC1200645), from algae associated to Rizophora mangle and Avicenia schaueriana, Fazenda River, municipality of Ubatuba, São Paulo State, Brazil (23°21'385"S, 44°50'385"W).

Superfamily **Tydeoidea** (12 species)

Family **Tydeidae** (11 species)

***Lorryia bathorii* Ripka & Kazmierski, 2013 in Ripka, Laniecka & Kazmierski, 2013: 24**—Holotype male (AMU, AK H-119), from *Tilia cordata*, Zala County, Keszthely, Hungary.

***Lorryia hungarica* Laniecka & Ripka, 2013 in Ripka, Laniecka & Kazmierski, 2013: 14**—Holotype female (AMU, AK H-027), from *Populus x canescens*, Óbuda, Budapest III, Hungary.

***Lorryia sanctikingai* Ripka & Laniecka, 2013 in Ripka, Laniecka & Kazmierski, 2013: 19**—Holotype female (AMU, AK H-066), from *Forsythia suspense*, Kispest, Budapest XIX, Hungary.

***Neopronematus solani* Laniecka & Kazmierski, 2013: 34**—Holotype female (AMU, AK H-126), from *Solanum dulcamara*, Törökvesz, Budapest II, Hungary.

***Prelorryia labrusca* Silva, Cunha & Ferla, 2014 in Silva, Cunha, Rocha, Panou & Ferla, 2014: 506**—Holotype female (ESALQ), from *Vitis labrusca*, Boqueirão do Leão County, State of Rio Grande do Sul, Brazil.

***Tydeus clavimaculatus* Kazmierski, 2013 in Ripka, Laniecka & Kazmierski, 2013: 10**—Holotype Tritonymph (AMU, AK H-295), from *Aesculus pavia*, Veszprém County, Zirc, Hungary.

***Tydeus manoi* Silva, Rocha & Ferla, 2014 in Silva, Cunha, Rocha, Panou & Ferla, 2014: 506**—Holotype female (ESALQ), from *Plantago tomentosa*, Candiota County, State of Rio Grande do Sul, Brazil.

***Tydeus martae* Kazmierski, 2013 in Ripka, Laniecka & Kazmierski, 2013: 5**—Holotype female (AMU, AK H-024), from *Hordeum vulgare*, Jász-Nagykun-Szolnok County, Szolnok, Hungary.

***Tydeus narolicatus* Kazmierski & Ripka, 2013 in Ripka, Laniecka & Kazmierski, 2013: 2**—Holotype female (AMU, AK H-337), from *Acer saccharinum*, Vas County, Mesteri, Hungary.

***Tydeus riopardensis* Silva, Cunha & Ferla, 2014 in Silva, Cunha, Rocha, Panou & Ferla, 2014: 503**—Holotype female (ZAUMCN), from *Vitis labrusca*, Bordeaux variety, Boqueirão do Leão County, State of Rio Grande do Sul, Brazil.

***Tydeus shabestariensis* Akbari, Haddad & Kaźmierski, 2015 in Akbari, Irani-Nejad, Khanjani, Arzanlou & Kaźmierski, 2015b: 265**—Holotype female (UTI), from the soil under apple orchards, vicinity of Shabestar city, East Azerbaijan, Iran.

Family **Iolinidae** (1 species)

***Neopronematus kamalii* Darbemamieh & Hajiqanbar, 2015 in Darbemamieh, Hajiqanbar, Khanjani & Kaźmierski, 2015: 236**—Holotype female (TMU), from apricot leaves, Sechek, Kermanshah, Iran (34°21'01.76"N, 47°18'01.38"E).

Infraorder **Anystina** (174 species)

Hypoder **Anystae** (7 species)

Superfamily **Anystoidea** (5 species)

Family **Pseudocheylidae** (4 species)

***Anoplocheylus kazemii* Bagheri, 2013 in Bagheri, Zarei, Ahaniazad, Gharekhany & Navaei-Bonab, 2013: 294**—Holotype female (PPRP), from the soil from apple tree orchards, Miandoab, West Azerbaijan Province, Iran.

***Anoplocheylus marivaniensis* Khanjani, Hoseini & Amini, 2014: 189**—Holotype female (BASU), from soil and rotten leaves of oak trees, *Quercus brantii*, Marivan vicinity, Kurdistan Province, Iran (35°26'N, 46°13'E).

Anoplocheylus qorvehiensis Khanjani, Hoseini & Amini, 2014: 192—Holotype female (BASU), from soil under Astragalus sp., Qorveh vicinity, Kurdistan Province, Iran (47°47'06.33"N, 35°09'03.62"E).

Anoplocheylus sinai Bagheri, 2013 in Bagheri, Zarei, Ahaniazad, Gharekhany & Navaei-Bonab, 2013: 293—Holotype female (PPRP), from soil in apple and blackcherry orchards, Miandoab and Azarshahr, East Azerbaijan Province, Iran.

Family **Teneriffiidae** (1 species)

Austroteneriffia shiraziensis Khanjani, Yazdanpanah & Fayaz, 2013: 49—Holotype female (BASU), from oak soil, Koohmara Sorkhi region, Fars Province, Iran (29°30'51"N, 52°11'43"E).

Superfamily **Caeculoidea** (1 species)

Family **Caeculidae** (1 species)

Neocaeculus orientalis Fuangarworn & Butcher, 2015: 262—Holotype female (CUMNH), from Lai Nan Sub-dist, Wiang Sa Dist., Nan Prov., Thailand (18°33'29.81"N, 100°47'48.67"E).

Superfamily **Paratydeoidea** (1 species)

Family **Paratydeidae** (1 species)

Tanytdeus beyzavii Khanjani, Nadri, Khanjani & Seeman, 2014: 175—Holotype female (BASU), from soil under gum bushes, Astragalus gossypinus, Alashtar, Kahman, Lorestan Province, Iran (33°53'09"N, 48°19'28"E).

Hypoder **Parasitengona** (167 species)

Superfamily **Erythraeoidea** (1 species)

Family **Erythraeidae** (1 species)

Charletonia cameroonensis Haitlinger & Kekeunou, 2014 in Haitlinger, Kekeunou & Lupicki, 2014: 41—Holotype larva (MNHWU), from Zonocerus variegatus, Zamakoe, Cameroon (3°34'44"N, 11°39'33"E).

Superfamily **Eylaoidea** (1 species)

Family **Limnocharidae** (1 species)

Limnochares spinosa Smit & Pešić, 2014: 5—Holotype male (RMNH), from little Lumotok stream, Borneo, Sayap, Mt Kinabalu, Malaysia (6°09.467"N, 116°34.027"E).

Superfamily **Trombiculidoidea** (25 species)

Family **Johnstonianidae** (2 species)

Diplothrombium ostovani Noei & Saboori, 2014 in Noei, Saboori & Hajizadeh, 2014: 248—Holotype larva (JAZM, ARS-20131003-2a), from soil (off host), forest close to river, Gijave village, Asalem city, Guilan Province, Iran (37°42.077"N, 48°54.077"E).

Diplothrombium sahragardi Noei & Saboori, 2014 in Noei, Saboori & Hajizadeh, 2014: 247—Holotype larva (JAZM, ARS-20131003-1a), from soil (off host), forest close to river, Gijave village, Asalem city, Guilan Province, Iran (37°42.077"N, 48°54.077"E).

Family **Trombiculidae** (23 species)

Diaguitacarus choapensis Stekolnikov & González-Acuña, 2015: 24—Holotype larva (ZISP, collection No. 8442, T-Tr.-72), from *Liolaemus pseudolemniscatus*, Las Chinchillas Natural Reserve, Coquimbo Region, Choapa Province, Chile.

Eutrombicula mistrali Stekolnikov & González-Acuña, 2015: 9—Holotype larva (ZISP, collection No. 9611, T-Tr.-67), from *Microlophus quadrivittatus*, Caleta Vitor, Arica and Parinacota Region, Arica Province, Chile (18°45"S, 70°20"W).

Eutrombicula nerudai Stekolnikov & González-Acuña, 2015: 3—Holotype larva (ZISP, collection No. 8492, T-Tr.-66), from *Liolaemus nitidus*, Llanos de Challe National Park, Atacama Region, Chile.

Eutrombicula picunche Stekolnikov & González-Acuña, 2015: 11—Holotype larva (ZISP, collection No. 8506, T-Tr.-68), from *Liolaemus fuscus*, La Campana National Park, Valparaíso Region, Quillota Province, Chile.

Leptotrombidium abramovi Stekolnikov, 2013: 34—Holotype larva (ZISP, T-Tr.-58), from *Hylomys megalotis*, Ban Doy village, 18 km N from Thakhek, Khammouane Province, Laos.

Leptotrombidium aenigmami Stekolnikov, 2013: 30—Holotype larva (ZISP, T-Tr.-57), from Laonastes aenigmamus, Ban Doy village, 18 km N from Thakhek, Khammouane Province, Laos.

Leptotrombidium bochkovi Stekolnikov, 2013: 60—Holotype larva (ZISP, L-316, T-Tr.-60), from Hylomys megalotis, Ban Doy village, 18 km N from Thakhek, Khammouane Province, Laos.

Leptotrombidium kinabalu Stekolnikov, 2013: 43—Holotype larva (IMR, L: 81572/b), from Dendrogale melanura, Kinabalu Mt, Sabah State, Malaysia.

Leptotrombidium laoense Stekolnikov, 2013: 92—Holotype larva (ZISP, T-Tr.-61), from Laonastes aenigmamus, Ban Doy village, 18 km N from Thakhek, Khammouane Province, Laos.

Leptotrombidium megabodense Stekolnikov, 2013: 42—Holotype larva (IMR, L: 82153/4)), from Tupaia montana, Kinabalu Mt, Sabah State, Malaysia.

Leptotrombidium megalangati Stekolnikov, 2013: 42—Holotype larva (IMR, L: 4159/1), from *Leopoldamys edwardsi*, Brinchang Mt, Pahang State, Malaysia.

Leptotrombidium megaloti Stekolnikov, 2013: 98—Holotype larva (ZISP, T-Tr.-62), from *Hylomys megalotis*, Ban Doy village, 18 km N from Thakhek, Khammouane Province, Laos.

Leptotrombidium minutui Stekolnikov, 2013: 44—Holotype larva (IMR, No 15924), from *Berylmys bowersi*, Ulu Langat, Selangor State, Malaysia.

Leptotrombidium saigoni Stekolnikov, 2013: 69—Holotype larva (IPHCMC, No 2004), from *Rattus rattus*, An Khe, Gia Lai Province, Vietnam.

Leptotrombidium tenompaki Stekolnikov, 2013: 45—Holotype larva (IMR, No19245), from Tupaia montana, Tenompak, Kinabalu Mt, Sabah State, Malaysia.

Leptotrombidium tikhonovi Stekolnikov, 2013: 50—Holotype larva (ZISP, T-Tr.-59), from *Hylomys megalotis*, Ban Doy village, 18 km N from Thakhek, Khammouane Province, Laos.

Leptotrombidium ului Stekolnikov, 2013: 44—Holotype larva (IMR, No 12247-1), from *Leopoldamys sabanus*, Ulu Langat, Selangor State, Malaysia.

Microtrombicula mapuche Stekolnikov & González-Acuña, 2015: 13—Holotype larva (ZISP, collection No. 7702, T-Tr.-69), from *Liolaemus chillanensis*, tourist camp Waldorf, Nevados de Chillán Mts, Biobío Región, Chile (36°51'41"S, 71°26'45"W).

Morelacarus camanchaca Stekolnikov & González-Acuña, 2015: 36—Holotype larva (ZISP, collection No. 8458, T-Tr.-75), from Liolaemus platei, Fray Jorge National Park, Coquimbo Region, Limarí Province, Chile.

Morelacarus jorgei Stekolnikov & González-Acuña, 2015: 34—Holotype larva (ZISP, collection No. 8451, T-Tr.-74), from Liolaemus platei, Fray Jorge National Park, Coquimbo Region, Limarí Province, Chile.

Parasacia molini Stekolnikov & González-Acuña, 2015: 16—Holotype larva (ZISP, collection No. 7707, T-Tr.-70), from Liolaemus chillanensis, Aguas Calientes, Nevados de Chillán Mts, Biobío Region, Chile (36°54'28"S, 71°22'14"W).

Paratrombicula philippii Stekolnikov & González-Acuña, 2015: 19—Holotype larva (ZISP, collection No. 8445, T-Tr.-71), from Liolaemus nigromaculatus, Fray Jorge National Park, Coquimbo Region, Limarí Province, Chile.

Proschoengastia antarctica Stekolnikov & González-Acuña, 2015: 30—Holotype larva (ZISP, collection No. 8483, T-Tr.-73), from *Neovison vison*, Navarino Island, Region of Magallanes and Antártica Chilena, Antártica Chilena Province, Chile.

Superfamily **Trombidioidea** (4 species)

Family **Achaemenothrombiidae** (1 species)

Achaemenothrombium dariusi Saboori, Wohltmann & Hakimitabar, 2013 *in* Saboori, Wohltmann, Hakimitabar & Shirvani, 2013: 144—Holotype larva (JAZM, ARS-20130205-1a), from *Euxoa fallax*, Kerman Province, Iran (30°11'8"N, 57°24'54"E).

Family **Microtrombidiidae** (1 species)

Empitrombium makolae Sevsay & Karakurt, 2013: 264—Holotype female (EUE), Köse mountain, Gümüşhane Province, Turkey (40°14'921N, 38°39'846E).

Family **Trombidiidae** (2 species)

Emitrombidium giocondi Mąkol & Sevsay, 2014: 96—Holotype adult (ZIZMUH), from pitfall traps, in olive groves, close to the seaside, Kırıkışlaçık village nr Muğla, Turkey (37°16'38.80"N, 27°33'47.97"E).

Paratrombium rovniense Konikiewicz & Mąkol, 2014: 588—Holotype larva (SIZK, K-8284-A), from “Pugach” mine, Klesov, Ukraine.

Superfamily **Hydryphantoidea** (1 species)

Family **Hydryphantidae** (1 species)

Protzia borneoensis Smit & Pešić, 2014: 7—Holotype female (RMNH), from torrential stream, Borneo, Alab-Kota Kinabalu road, Crocker Range, Malaysia (5°50.039'N, 116°20.156'E).

Superfamily **Hydrachnoidea** (6 species)

Family **Hydrachnidae** (5 species)

Hydrachna cordata Smit, 2014: 172—Holotype male (QM), Queensland, Lake Emma, Lakefield NP, Australia (15°17'47.43"S, 144°38'48.05"E).

Hydrachna curtiscutata Smit, 2014: 174—Holotype male (QM), Queensland, Temporary pond along Wyseby road, Australia (24°59.185'S, 148°24.416"E).

Hydrachna longiscutata Smit, 2014: 176—Holotype male (AMS), New South Wales, Artificial pool, Rockhole Loop, Mutawintji NP, Australia.

Hydrachna novae-hollandica Smit, 2014: 180—Holotype male (WAM), from Jackeroo's Waterhole, Western Australia, El Questro Station, The Kimberley, Australia.

Hydrachna queenslandica Smit, 2014: 182—Holotype male (QM), from Lake Emma, Queensland, Lakefield NP, Australia (15°17'47.43"S, 144°38'48.05"E).

Superfamily **Lebertioidea** (33 species)

Family **Sperchontidae** (2 species)

Sperchonopsis orientalis Smit & Pešić, 2014: 13—Holotype male (RMNH), from Mahua stream, Crocker Range, Borneo, Malaysia (5°45.225'N, 116°26.085'E).

Sperchon (Sperchon) kinabaluensis Smit & Pešić, 2014: 9—Holotype male (RMNH), from Kipungit River, Poring Hot Springs, Mt. Kinabalu, Malaysia (6°02.776'N, 116°41.432"E).

Family **Anisitsiellidae** (2 species)

Nilotonia (Dartonia) sketi Pešić, 2013a: 373—Holotype male (SMF), from pools of percolated water, Trung Trang Cave, Dao Cat Ba (= Cat Ba Island), Cat Ba National Park, W of Vinh Ha Long, Vietnam.

Hydrobaumia purpurea Smit, 2013: 207—Holotype female (RMNH), from Pandanus marsh, beach Base G, Jayapura, New Guinea, Papua Province, Indonesia (2°31.035'S, 140°44.360"E).

Family **Oxidae** (2 species)

Oxus (Oxus) fuscus Smit & Pešić, 2014: 15—Holotype female (RMNH), from Mahua stream, Borneo, Mahua, Crocker Range, Malaysia (5°47.838'N, 116°24.510"E).

Oxus (Oxus) manokwariensis Smit, 2013: 209—Holotype male (RMNH), from “Waterfall Creek”, Manokwari, West Papua Province, Indonesia (0°52.888'S, 134°02.628"E).

Family **Torrenticolidae** (27 species)

Monatractides epiales Pešić & Smit, 2014: 52—Holotype male (RMNH), from Mt Kinabalu, Poring Hot Springs, Kipungit River, Borneo, Malaysia (6°02.776'N, 116°41.432"E).

- Monatractides euterpe* Pešić & Smit, 2014: 42—Holotype male (RMNH), Atewa Hills, Unnamed creek upstream Sagyimase, Ghana ($6^{\circ}13.966'N$, $0^{\circ}33.114'W$).
- Monatractides heracles* Pešić & Smit, 2014: 60—Holotype male (RMNH), from Mt Kinabalu, Sayap, Little Lumotok stream, Borneo, Malaysia ($6^{\circ}09.467'N$, $116^{\circ}34.027'E$).
- Monatractides melpomene* Pešić & Smit, 2014: 44—Holotype male (RMNH), Amedzofe Falls, Ghana ($6^{\circ}50.656'N$, $0^{\circ}26.868'E$).
- Monatractides minuta* Pešić & Smit, 2014: 63—Holotype female (RMNH), from Mt Kinabalu, Great Lumotok stream, Borneo, Malaysia ($6^{\circ}09.336'N$, $116^{\circ}34.056'E$).
- Monatractides morpheus* Pešić & Smit, 2014: 54—Holotype male (RMNH), from Mt Kinabalu, small stream Layang Layang, Borneo, Malaysia ($6^{\circ}02.711'N$, $116^{\circ}33.627'E$).
- Monatractides phantasos* Pešić & Smit, 2014: 57—Holotype female (RMNH), from Mt Kinabalu, Sayap, stream Kemantis, Borneo, Malaysia ($6^{\circ}09.841'N$, $116^{\circ}33.936'E$).
- Monatractides phobetor* Pešić & Smit, 2014: 59—Holotype male (RMNH), from Mt Kinabalu, Sayap, Little Lumotok stream, Borneo, Malaysia ($6^{\circ}09.467'N$, $116^{\circ}34.027'E$).
- Monatractides polyhymnia* Pešić & Smit, 2014: 46—Holotype male (RMNH), Kyabobo NP, Kue River, Ghana ($8^{\circ}31.087'N$, $0^{\circ}36.049'E$).
- Monatractides terpsichore* Pešić & Smit, 2014: 51—Holotype female (RMNH), Agumatsa Wildlife Sanctuary, Plunge pool Agumatsa Falls, Ghana ($7^{\circ}06.350'N$, $0^{\circ}36.476'E$).
- Monatractides thalia* Pešić & Smit, 2014: 49—Holotype male (RMNH), Agumatsa Wildlife Sanctuary, Agumatsa River at first bridge, Ghana ($7^{\circ}06.830'N$, $0^{\circ}35.760'E$).
- Monatractides urania* Pešić & Smit, 2014: 54—Holotype male (RMNH), Kintampo Falls, Ghana ($8^{\circ}05.413'N$, $1^{\circ}41.881'W$).
- Neoatractides (Allotorrenticola) erato* Pešić & Smit, 2014: 24—Holotype male (RMNH), Atewa Hills, small rainforest stream upstream Sagyimase, Ghana ($6^{\circ}13.964'N$, $0^{\circ}33.116'W$).
- Neoatractides sundaensis* Pešić & Smit, 2014: 28—Holotype female (RMNH), Mt Kinabalu, Sayap, stream Kemantis, Borneo, Malaysia ($6^{\circ}09.841'N$, $116^{\circ}33.936'E$).
- Neoatractides uniscutatus* Pešić & Smit, 2014: 32—Holotype male (RMNH), Crocker Range, Inobong, unnamed stream Bansadon Trail, Borneo, Malaysia ($5^{\circ}51.456'N$, $116^{\circ}68.403'E$).
- Pseudotorrenticola borneoensis* Pešić & Smit, 2014: 32—Holotype male (RMNH), from Mt Kinabalu, Sayap, Little Lumotok stream, Borneo, Malaysia ($6^{\circ}09.497'N$, $116^{\circ}34.027'E$).
- Stygotorrenticola coniseta* Pešić & Gerecke, 2014: 373—Holotype male (SMF), from gours of dripping water- internal canyon, Grotta passante, Tam Son, Quan Ba, Ha Giang, QBTS 03, Vietnam ($23^{\circ}02'44.7"N$, $104^{\circ}59'23.9"E$).
- Torrenticola borneoensis* Pešić & Smit, 2014: 6—Holotype male (RMNH), from Mt Kinabalu, Sayap, stream Kemantis, Borneo, Malaysia ($6^{\circ}09.841'N$, $116^{\circ}33.936'E$).
- Torrenticola calliope* Pešić & Smit, 2014: 21—Holotype male (RMNH, ZMA.ACAR.32333), Roby River, Ethiopia ($9^{\circ}44.996'N$, $38^{\circ}59.743'E$).
- Torrenticola comorosensis* Pešić & Smit, 2015 in Pešić, Smit & Mary, 2015: 447—Holotype male (MNHN), from River Koualé tributary Mayotte, Comoros ($12^{\circ}47'57.3"S$, $45^{\circ}09'51.2"E$).
- Torrenticola episce* Pešić, Chatterjee, Das & Bordoloi, 2013: 455—Holotype male (MNHP), from the gut content of *Botia Dario*, Kakarikata beel Majuli Island, Jorhat district, Assam State, India.
- Torrenticola kinabaluensis* Pešić & Smit, 2014: 8—Holotype male (RMNH), from Crocker Range, downstream of national park entrance, Mahua stream, Borneo, Malaysia ($5^{\circ}46.491'N$, $116^{\circ}25.770'E$).
- Torrenticola (Megapalpis) anophthalma* Pešić & Gerecke, 2014: 368—Holotype female (SMF), from gours of dripping water-internal canyon, transition cave, Tam Son, Quan Ba, Ha Giang, QBTS 03, Vietnam ($23^{\circ}02'44.7"N$, $104^{\circ}59'23.9"E$).

Torrenticola neoindica Pešić & Smit, 2014: 20—Holotype male (RMNH), from Crocker Range, downstream of national park entrance, Mahua stream, Borneo, Malaysia (5°46.491'N, 116°25.770'E).

Torrenticola sabahensis Pešić & Smit, 2014: 11—Holotype male (RMNH), from Mt Kinabalu, Great Lumotok stream, Borneo, Malaysia (6°09.336'N, 116°08.417'E).

Torrenticola schilthuizeni Pešić & Smit, 2014: 25—Holotype male (RMNH), from Crocker Range, Kibamabangan River, Borneo, Malaysia (5°51.28'N, 116°08.417'E).

Torrenticola (Torrenticola) vietnamica Tuzovskij, 2013a: 548—Holotype female (IBIW, slide 9518), from Khe (Khé) River, the inflow of Cai River, Khanh Hoa Province, Vietnam (12°16.741'N, 108°54.669'E).

Superfamily **Hygrobatoidea** (87 species)

Family **Pontarachnidae** (3 species)

Litarachna gracilis Pešić, 2013b: 87—Holotype female (NIBR, 0000268890), from Gangneung beach, East Sea, South Korea (37°47.824'N, 128°55.085'E).

Litarachna lukai Pešić, 2013b: 85—Holotype male (NIBR, 0000268888), from beach, Wido Island, Yellow Sea, South Korea (35°35.089'N, 126°15.196'E).

Pontarachna ivonae Pešić, 2013b: 89—Holotype male (NIBR, 0000268892), from beach, Wido Island, South Korea (35°35.089'N, 126°15.196'E).

Family **Limnesiidae** (4 species)

Limnesia (Limnesia) multiscutata Smit, 2013: 210—Holotype female (RMNH), from stream Ilik, near Lake Gita, Anggi Lakes, New Guinea, West Papua Province, Indonesia (1°21.423'S, 133°58.553'E).

Limnesia (Limnesia) vietnamensis Tuzovskij, 2013a: 550—Holotype female (IBIW, slide 9694), from forest stream Dak Lua (Đák Lua), Cat Tien (Cát Tiên) National Park, Dong Nai (Đồng Nai) Province, Vietnam (11°29.844'N, 107°22.847'E).

Limnesia (Limnesia) vogelkop Smit, 2013: 212—Holotype male (RMNH), from stream ± 3 km S of Mokwam, West Papua Province, Indonesia (1°05.877'S, 133°54.223'E).

Raptorhydracarus tomasini Pešić & Gerecke, 2014: 376—Holotype male (SMF), from gours (rimstone pools) of dropping water end of the cave right arm, Grotta Ta Lu,Dong Van, Ha Giang, DV 08, Vietnam (23°15'37.8"N, 105°21'38.5"E).

Family **Frontipodopsidae** (1 species)

Frontipodopsis suturalis Smit & Pešić, 2014: 36—Holotype female (RMNH), from Mahua stream, Borneo, Mahua, Crocker Range, Malaysia (5°47.838'N, 116°24.510'E).

Family **Hygrobatidae** (36 species)

Atractides (Atractides) albaruthenicus Cichońka & Biesiadka, 2013: 254—Holotype male (UWMO), Suchaja Dolina, river Svisloč, Belarus (53°47'459"N, 23°97'78"E).

Atractides (Atractides) crockerensis Smit & Pešić, 2014: 29—Holotype male (RMNH), from Mahua river, upper course, Borneo, Malaysia (5°47.939'N, 116°24.317'E).

Atractides (Atractides) ghanaensis Pešić & Smit, 2015b: 344—Holotype male (RMNH), from small rainforest stream upstream Sagyimase, Atewa Hills, Ghana (6°13.964'N, 0°33.116'W).

Atractides (Atractides) neospatiosus Smit & Pešić, 2014: 24—Holotype male (RMNH), from first stream Minduk Sirung Trail, Borneo, Alab, Crocker Range, Malaysia (5°49.249'N, 116°19.862'E).

Atractides (Atractides) sabahensis Smit & Pešić, 2014: 28—Holotype male (RMNH), from Mahua stream at crossing with main road, Borneo, Crocker Range, Malaysia (5°45.225'N, 116°26.085'E).

Atractides (Atractides) svislocensis Cichońka & Biesiadka, 2013: 258—Holotype male (UWMO), Suchaja Dolina, river Svisloč, Belarus (53°47'459"N, 23°97'78"E).

Atractides curtisetus Smit & Pešić, 2014: 33—Holotype female (RMNH), from Little Lumotok stream, Borneo, Sayap, Mt Kinabalu, Malaysia (6°09.467'N, 116°34.027'E).

- Atractides reinhardi* Gütte, Gütte & Boyaci, 2015: 246—Holotype female (MAEU), from a pristine spring with dense emergent vegetation (*Nasturtium officinale*) with sandy and muddy bottom, Çavdır, Burdur, Turkey (37°9.44'N, 29°45.92'E).
- Atractides (Tympanomegapus) borneensis* Smit & Pešić, 2014: 35—Holotype male (RMNH), from Mahua stream, Borneo, Mahua, Crocker Range, Malaysia (5°47.838'N, 116°24.510'E).
- Australiobates elongatus* Smit, 2013: 215—Holotype female (RMNH), from Mokwam, New Guinea, West Papua Province, Indonesia (1°00.608'S, 133°53.863'E).
- Dropursides aberratus* Smit, 2013: 220—Holotype female (RMNH), from stream Wasai, SW of Manokwari, New Guinea, West Papua Province, Indonesia (0°55.563'S, 133°58.380'E).
- Hygrobates acutipalpis* Smit & Pešić, 2014: 15—Holotype male (RMNH), from Silau-Silau stream, park headquarters, Borneo, Mt Kinabalu, Malaysia (6°00.595'N, 116°32.497'E).
- Hygrobates hamatoides* Smit & Pešić, 2014: 18—Holotype male (RMNH), from small stream Panar Laban Hut, Mt Kinabalu, Malaysia (6°03.641'N, 116°33.968'E).
- Hygrobates (Hygrobates) gusakovi* Tuzovskij, 2013a: 554—Holotype female (IBIW, slide 9697), from forest stream Dak Lua (Đák Lua), Cat Tien (Cát Tiên) National Park, Dong Nai (Đồng Nai) Province, Vietnam (11°29.844'N, 107°22.847'E).
- Hygrobates (Rivobates) zawali* Pešić, 2015: 288—Holotype male (MNHP), from Lake Ohrid, Macedonia.
- Hygrobates striatus* Smit & Pešić, 2014: 19—Holotype female (RMNH), from Kipungit River, Poring Hot Springs, Mt Kinabalu, Malaysia (6°02.776'N, 116°41.432'E).
- Gondwanabates crassipalpis* Smit, 2015b: 570—Holotype male (QM), from Dalrymple Creek, Goomburra Forest Reserve, Queensland, Australia (27°58.781'S, 152°20.621'E).
- Gondwanabates flavus* Smit, 2015b: 572—Holotype female (QM), from aire River at crossing with Binns Road, Great Otway Ranges NP, Victoria, Australia (38°40.103'S, 143°34.805'E).
- Gondwanabates spatulifer* Smit, 2015b: 574—Holotype male (NMV), from Wannon River at crossing with Serra Road, Grampians NP, Victoria, Australia (37°20.918'S, 142°30.386'E).
- Procorticacarus curtipalpis* Smit, 2015c: 59—Holotype female (NTM), from Moline Bottom Rockhole, Kakadu NP, Northern Territory, Australia (13°34.418'S, 132°15.290'E).
- Procorticacarus kosciuszko* Smit, 2015c: 61—Holotype female (AMS), from Small unnamed stream below summit of Mt Kosciuszko, New South Wales, Mt Kosciuszko NP, Australia (36°27.322'S, 148°15.929'E).
- Procorticacarus nasutus* Smit, 2015c: 64—Holotype female (QM), from double Barrel Creek, Queensland, Babinda, Australia (17°20.852'S, 145°53.075'E).
- Procorticacarus neocaledonicus* Smit, 2015c: 66—Holotype male (MNHN), from Trou Bleu River, New Caledonia, Australia (22°20'12.3"S, 166°57'44.4"E).
- Procorticacarus neoprasadi* Smit, 2015c: 66—Holotype male (AMS), from Sawpit Creek at crossing with Kosciuszko Road, Mt Kosciuszko NP, New South Wales, Australia (36°20.984'S, 148°33.818'E).
- Procorticacarus paulae* Smit, 2015c: 68—Holotype male (WAM), from Carey Brook at crossing with Vasse Highway, Western Australia, Australia (34°24.876'S, 115°48.717'E).
- Procorticacarus quadriscutatus* Smit, 2015c: 69—Holotype female (AMS), from Carruthers Peak, Mt Kosciuszko NP, New South Wales, Australia (36°24.621'S, 148°18.338'E).
- Procorticacarus queenslandicus* Smit, 2015c: 71—Holotype female (RMNH), from Babinda River, Wooroonooran NP, Australia (17°20.484'S, 145°52.113'E).
- Procorticacarus septentrionalis* Smit, 2015c: 73—Holotype male (NTM), from Florence Creek downstream of Florence Falls, Litchfield NP, Northern Territory, Australia (13°05.885'S, 130°46.999'E).
- Procorticacarus wenlockensis* Smit, 2015c: 75—Holotype female (QM), from Wenlock River at crossing with Telegraph Road, Cape York Peninsula, Queensland, Australia (12°27'20.65"S, 142°38'27.87"E).

Protzia longiacetabulata Gütte & Boyaci, 2014: 585—Holotype female (FFSDU), from fast-flowing stream with gravel bottom, Dim Stream, Alanya, Turkey (36°35.321'N, 32°17.876'E).

Rhynchaustrobates (Rhynchaustrobates) sclerosus Smit, 2015b: 576—Holotype male (AMS), from Sugarloaf Creek, at crossing with Misty Mountain Road, Monga NP, New South Wales, Australia (35°33'44.70"S, 150°0'48.61"E).

Rhynchaustrobates (Rhynchaustrobates) truusae Smit, 2015b: 579—Holotype male (AMS), from little Wonder Creek, near end of Buckra—Bendinni Road, New South Wales, Australia (30°30.034"S, 152°39.387'E).

Rhynchaustrobates (Rhynchaustrobates) victorianus Smit, 2015b: 580—Holotype male (NMV), from scrubby Grampians NP, Valley Creek at Flat Rock Crossing, Australia (37°09.680"S, 142°26.680'E).

Rhynchaustrobates (Victoriabates) bipartitus Smit, 2015b: 580—Holotype female (QM), from Tully River at campground, Tully Gorge NP, Queensland, Australia (17°46.280"S, 145°39.073'E).

Rhynchaustrobates (Victoriabates) uniscutatus Smit, 2015b: 582—Holotype female (NTM), from Gubarra Creek, upstream of pool, Kakadu NP, Northern Territory, Australia (12°49.605"S, 132°52.708'E).

Sterkspruitia comorosensis Pešić, Smit & Mary, 2015: 456—Holotype male (MNHN), from River Djalimou, Mayotte, Comoros (12°57'13.8"S, 45°06'52.7"E).

Family Aturidae (30 species)

Albaxona gusevi Tuzovskij, 2013a: 557—Holotype female (IBIW, slide 9520), from Khé River near the inflow of Cai River, Khanh Hoa Province, Vietnam (12°16.741'N, 108°54.669'E).

Albaxona mahuaensis Smit & Pešić, 2014: 49—Holotype male (RMNH), from Mahua stream at crossing with main road, Borneo, Crocker Range, Malaysia (5°45.225'N, 116°26.085'E).

Albia (Albiella) ilikensis Smit, 2013: 228—Holotype female (RMNH), from stream Ilik, near lake Gita, Anggi Lakes, New Guinea, West Papua Province, Indonesia (1°21.423'S 133°58.553'E).

Albia (Albiella) papuaensis Smit, 2013: 229—Holotype female (RMNH), from unnamed creek, Arso-Senggi, New Guinea, Papua Province, Indonesia (3°03.020'S, 140°50.685'E).

Albia crocker Smit & Pešić, 2014: 62—Holotype male (RMNH), from small stream Waterfall Trial, Inobong, Borneo, Crocker Range, Malaysia (5°51.306'N, 116°08.292'E).

Aturus borneoensis Smit & Pešić, 2014: 64—Holotype male (RMNH), from Kipungit River, Poring Hot Springs, Mt Kinabalu, Malaysia (6°02.776'N, 116°41.432'E).

Axonopsella bisetosa Smit, 2013: 231—Holotype male (RMNH), from small stream upstream, Kobrey, near lake Gigi, Anggi Lakes, West Papua Province, Indonesia (1°23.328'S, 133°54.345'E).

Axonopsis caeca Smit & Pešić, 2014: 57—Holotype female (RMNH), from Mahua stream, Mahua, Borneo, Crocker Range, Malaysia (5°47.838'N, 116°24.510'E).

Axonopsis latipalpis Smit & Pešić, 2014: 53—Holotype male (RMNH), from Unnamed creek Bansadon Trial, Inobong, Borneo, Crocker Range, Malaysia (5°51.456'N, 116°08.403'E).

Axonopsis longigenitalis Smit & Pešić, 2014: 50—Holotype male (RMNH), from Kibamabangan River at waterfall, Inobong, Borneo, Crocker Range, Malaysia (5°51.280'N, 116°08.417'E).

Axonopsis rugosa Smit & Pešić, 2014: 52—Holotype female (RMNH), from Kipungit River, Poring Hot Springs, Mt Kinabalu, Malaysia (6°02.776'N, 116°41.432'E).

Axonopsis truncata Smit & Pešić, 2014: 54—Holotype male (RMNH), from Kipungit River, Poring Hot Springs, Borneo, Mt Kinabalu, Malaysia (6°02.776'N, 116°41.432'E).

Brachypoda (Hemibrachypoda) orientalis Pešić & Esen, 2013 in Esen, Pešić & Erman, 2013: 327—Holotype male (FUE), from Mezra stream, Pütürge, Malatya Province, Turkey (38°15.01'N, 38°38.37'E).

Brachypoda milicuae Pešić & Semenchenko, 2014: 336—Holotype male (NIBR), from stream near Naebyeansan Info Center, CR9 Ne myeon Mt, Naebyeansan NP, South Korea (35°38'25.623"N, 126°34'53.1438"E).

- Brachypoda rossica* Pešić & Semenchenko, 2014: 342—Holotype male (300-kas IBSS), from Levaya Burinda River, Magdagachinsky District, Amurskaya Province, Russia (53°27.731'N, 125°07.808'E).
- Brachypoda sokolowi* Pešić & Semenchenko, 2014: 339—Holotype male (299-kas IBSS), from Levaya Burinda River, Magdagachinsky District, Amurskaya Province, Russia (53°27.731'N, 125°07.808'E).
- Erebaxonopsis kipungitensis* Smit & Pešić, 2014: 58—Holotype male (RMNH), from Kipungit River, Poring Hot Springs, Mt Kinabalu, Malaysia (6°02.776'N, 116°41.432'E).
- Javalbia borneoensis* Smit & Pešić, 2014: 41—Holotype male (RMNH), from small stream Waterfall Trial, Borneo, Inobong, Crocker Range, Malaysia (5°51.306'N, 116°08.292'E).
- Javalbia kinabaluensis* Smit & Pešić, 2014: 44—Holotype male (RMNH), from Liwagu stream, at crossing with summit trail, Mt Kinabalu, Borneo, Malaysia (6°01.868'N, 116°32.912'E).
- Javalbia magniseta* Smit & Pešić, 2014: 44—Holotype female (RMNH), from Mahua stream at crossing with main road, Borneo, Crocker Range, Malaysia (5°45.225'N, 116°26.085'E).
- Javalbia montana* Smit & Pešić, 2014: 38—Holotype male (RMNH), from Liwagu stream, at crossing with summit trail, Borneo, Mt Kinabalu, Malaysia (6°01.868'N, 116°32.912'E).
- Javalbia reticulata* Smit & Pešić, 2014: 47—Holotype male (RMNH), from Small stream Layang Layang, Mt Kinabalu, Malaysia (6°02.711'N, 116°33.627'E).
- Javalbia solitaria* Smit & Pešić, 2014: 40—Holotype male (RMNH), from Kemantis stream, Borneo, Sayap, Mt Kinabalu, Malaysia (6°09.841'N, 116°33.936'E).
- Javalbia uncinata* Smit & Pešić, 2014: 47—Holotype male (RMNH), from Mahua stream, downstream of national park entrance, at crossing with national park road, Borneo, Crocker Range, Malaysia (5°46.491'N, 116°25.770'E).
- Kongsbergia albanorum* Gerecke, 2014: 29—Holotype male (NHMW), 620 m, UC 50 06, stream Lumi i Varfrit, Sicily (PA) Piana degli Albanesi, I 495 Int, Sicily, Italy.
- Kongsbergia jaentschi* Gerecke, 2014: 32—Holotype male (NHMW), 620 m, UC 50 06, stream Lumi i Varfrit, P. d. Albanesi (PA), I 495 Int, Sicily, Italy.
- Ljania inconspicua* Smit & Pešić, 2014: 61—Holotype female (RMNH), from Mahua stream at crossing with main road, Borneo, Crocker Range, Malaysia (5°45.225'N, 116°26.085'E).
- Ljania obliteratea* Smit & Pešić, 2014: 62—Holotype female (RMNH), from Kemantis stream, Sayap, Borneo, Mt Kinabalu, Malaysia (6°09.841'N, 116°33.936'E).
- Woolastookia basilicalabrifica* Gerecke, 2014: 38—Holotype male (NHMW), spring brook, Fonte Cannello, Saracena Pellegrino (CS), Calabria, Italy.
- Zelandaturus magnipes* Smit, 2015a: 444—Holotype male (MNZ), from Camping Gully stream near Rakaia Gorge, hyporheic, South Island, New Zealand (43°31.608'S, 171°40.345'E).

Family **Unionicolidae** (10 species)

- Koenikea (Notomideopsis) sentaniensis* Smit, 2013: 226—Holotype male (RMNH), from Lake Sentani, , New Guinea, Papua Province, Indonesia (2°33.576'S, 140°25.438'E).
- Neumania (Neumania) silvestris* Smit, 2013: 223—Holotype male (RMNH), from unnamed creek, UNCEN-forest, Abepura, New Guinea, Papua Province, Indonesia (2°34.925'S, 140°39.173'E).
- Neumania (Soarella) comorosensis* Pešić & Smit, 2015 in Pešić, Smit & Mary, 2015: 590—Holotype female (MNHN), from River Djalimou, upstream, Mayotte, Comoros (12°56'58.4"S, 45°07'23.7"E).
- Recifella (Eorecifella) fluvialis* Smit, 2013: 223—Holotype male (RMNH), from River Kamp Walker, Abepura, New Guinea, Papua Province, Indonesia (2°34.202'S, 140°38.886'E).
- Unionicola aculeatella* Tuzovskij & Semenchenko, 2015: 424—Holotype male (IBIW-6908), from Gusinoe Lake, Park “Samarskaya Luka”, National natural, Stavropol District, Samara Province, Russia.

Unionicola primoryensis Tuzovskij & Semenchenko, 2015: 445—Holotype male (IBSS-kas-325), from mantle cavity of *Lanceolaria chankensis*, Astrakhansky Bay, Khankaysky District, Khanka Lake, Primory Territory, Russia.

Unionicola rossica Tuzovskij & Semenchenko, 2015: 408—Holotype female (IBIW-2819), from small lakes near settlement Markovo, Anadyr District, Magadan Province, Russia.

Unionicola samaraensis Tuzovskij & Semenchenko, 2015: 434—Holotype female (IBIW-7453), from Kondurcha River (right inflow of the Sok River), Krasnoyarsk District, Samara Province, Russia.

Unionicola setipella Tuzovskij & Semenchenko, 2015: 436—Holotype male (IBIW-3750), from Rybinsk reservoir near village Pogorelka, Nekouz District, Yaroslavl Province, Russia.

Unionicola (Unionicola) novaeguineae Smit, 2013: 220—Holotype male (RMNH), from Lake Sentani, Papua Province, Indonesia (2°33.576'S, 140°25.438'E).

Family **Pionidae** (3 species)

Piona anadyrensis Tuzovskij, 2013c: 327—Holotype male (IBIW, slide 2857), from small thermocarstic lake, settlement Markovo, Anadyr District, Chukotka, Russia.

Piona interrupta interruptella Tuzovskij, 2013c: 331—Holotype female (IBIW, slide 2305), from Majorskoe Lake, settlement Markovo, Anadyr District, Chukotka, Russia.

Piona magadanensis Tuzovskij, 2013c: 322—Holotype male (IBIW, slide 2430), from small forest pond, settlement Ola, Ola District, Magadan Province, Russia.

Superfamily **Arrenuroidea** (10 species)

Family **Mideopsidae** (1 species)

Xystonotus (Xystonotus) phytotelmaticola Pešić, 2015 in Pešić, Piccoli, Araújo, Rezende & Gonçalves: 149—Holotype male (PEU), from *Vrisea procera*, Ilha do Cardoso, Cananéia, São Paulo, Brazil (25°04'16"S, 47°55'23"W).

Family **Athienemanniidae** (2 species)

Africasia acuticoxalis Smit & Pešić, 2014: 67—Holotype female (RMNH), from unnamed creek Bansadon Trail, Borneo, Inobong, Crocker Range, Malaysia (5°51.456'N, 116°08.403'E).

Africasia vietnamitica Pešić & Gerecke, 2014: 376—Holotype female (SMF), from small lake - siphon in right arm, Dragon Cave, Sang Tung, Dong Van, Ha Giang, DV 09, Vietnam (23°12'41.8"N, 105°14'12.9"E).

Family **Arrenuridae** (7 species)

Arrenurus (Megaluracarus) anitahoffmannae Ramírez-Sánchez & Rivas, 2013: 318—Holotype male (UNAM), Del Negro lake, Villahermosa, Tabasco, Mexico (18°00'31"N, 92°55'08"W).

Arrenurus (Megaluracarus) maya Ramírez-Sánchez & Rivas, 2013: 326—Holotype male (UNAM), Chac-Ha cenote, 3.5 km northeast of Calotmul, Yucatan, Mexico (21°04'40"N, 88°08'27"W).

Arrenurus (Megaluracarus) olmeca Ramírez-Sánchez & Rivas, 2013: 321—Holotype male (UNAM), Loma del Caballo lake, route 180 km 7.5, Centro, Tabasco, Mexico (17°59'01"N, 93°00'04"W).

Arrenurus (Megaluracarus) urbanus Ramírez-Sánchez & Rivas, 2013: 328—Holotype male (UNAM), Cuemanco canal, Xochimilco ecological park, Mexico City, Mexico (19°17'20"N, 99°06'06"W).

Arrenurus (Rhinophoracarus) comorosensis Pešić & Smit, 2015 in Pešić, Smit & Mary, 2015: 593—Holotype female (MNHN), from River Koualé, upstream, Mayotte, Comoros (12°48'46. 5"S, 45°09'40. 9"E).

Arrenurus ovatipetiolatus Esen & Erman, 2013 in Esen, Erman & Dilkaraoglu, 2013: 74—Holotype male (FUE), Lake Bağlama, Pazarcık, Kahramanmaraş Province, Turkey (37°16.52'N, 37°07.30'E).

Thoracophoracarus uniacetabulatus Smit & Pešić, 2014: 67—Holotype female (RMNH), from unnamed creek Bansadon Trail, Borneo, Inobong, Crocker Range, Malaysia (5°51.456'N, 116°08.403'E).

Infraorder **Eleutherengona** (155 species)

Hypoder **Raphignathina** (135 species)

Superfamily **Cheyletoidea** (79 species)

Family **Harpirhynchidae** (47 species)

Anharpyrhynchus apodus Bochkov & Klompen, 2014a: 320—Holotype female (OSAL, 0083789), from *Lichmera indistincta*, Beagle Bay, Western Australia, Australia ($16^{\circ}53'24"S, 122^{\circ}31'12"E$).

Anharpyrhynchus elizae Bochkov & Klompen, 2014a: 317—Holotype female (OSAL, 0083656), from *Cyanocitta cristata*, O'Leno State Park, Columbia Co., Florida, USA ($29^{\circ}55'12"N, 82^{\circ}34'48"W$).

Anharpyrhynchus lukoschusi Bochkov & Klompen, 2014a: 322—Holotype female (OSAL, 0083777), from *Manorina flavigula*, Mount Hart, Western Australia, Australia ($17^{\circ}01'12"S, 125^{\circ}04'12"E$).

Crassacarus alexfaini Bochkov, OConnor & Klompen, 2015: 68—Holotype female (UMMZ, BMOC 09-1222-004, #1), from *Cardinalis cardinalis*, Livingston Co., Michigan, USA.

Crassacarus fritschi Bochkov, OConnor & Klompen, 2015: 72—Holotype female (UMMZ, BMOC 08-1110-002, #1), from *Bombycilla cedrorum*, Muskegon State game Area, along US route 31, Muskegon Co., Michigan, USA ($43^{\circ}17'46"N, 86^{\circ}04'01"W$).

Crassacarus melanerpes Bochkov, OConnor & Klompen, 2015: 76—Holotype female (OSAL, 0077574), from *Melanerpes formicivorus*, Craig, Johnson Co., Kansas, USA ($38^{\circ}58'12"N, 94^{\circ}49'12"W$).

Crassacarus sialia Bochkov, OConnor & Klompen, 2015: 73—Holotype female (OSAL, 0083648), from *Sialia currucoides*, Gillette, 19.3 km E of, Campbell Co., Wyoming, USA ($44^{\circ}17'24"N, 105^{\circ}30'00"W$).

Crassacarus tinae Bochkov, OConnor & Klompen, 2015: 70—Holotype female (OSAL, 0083653), from *Carduelis tristis*, Gillette, Campbell Co., Wyoming, USA ($44^{\circ}17'24"N, 105^{\circ}30'00"W$).

Fainharpirhynchus legatus Bochkov & Klompen, 2014a: 308—Holotype female (OSAL, 0083623), from *Legatus leucophaius*, 1.6 km E on Cumuto Road, Cumuto, Sangre Grande, Trinidad and Tobago ($10^{\circ}34'48"N, 61^{\circ}12'00"W$).

Fainharpirhynchus mossi Bochkov & Klompen, 2014a: 311—Holotype female (OSAL, 0083893), from *Myiarchus crinitus*, North America, no other data.

Harpypalpoides hirundinoides Bochkov & Klompen, 2014b: 469—Holotype female (OSAL, 0082920), from *Hirundo rustica*, Schaijk, Noord Brabant, The Netherlands ($51^{\circ}45'00"N, 5^{\circ}37'48"E$).

Harpypalpoides johnstoni Bochkov & Klompen, 2014b: 470—Holotype female (OSAL, 0082946), from *Fringilla coelebs*, Arnhem, Gelderland, The Netherlands ($51^{\circ}58'48"N, 5^{\circ}55'12"E$).

Harpypalpoides regulus Bochkov & Klompen, 2014b: 472—Holotype female (OSAL, 0082833), from *Regulus regulus*, Schaijk, Noord Brabant, The Netherlands ($51^{\circ}45'00"N, 5^{\circ}37'48"E$).

Harpypalpoides sita Bochkov & Klompen, 2014b: 474—Holotype female (OSAL, 0082882), from *Sitta pygmaea*, Monterey Co., California, USA ($35^{\circ}48'36"N, 121^{\circ}21'36"W$).

Harpypalpoides sylvia Bochkov & Klompen, 2014b: 473—Holotype female (OSAL, 0082874), from *Sylvia atricapilla*, Amsterdam, Noord-Holland, The Netherlands ($52^{\circ}22'12"N, 4^{\circ}53'24"E$).

Harypalpus lonchura Bochkov & Klompen, 2014b: 461—Holotype female (OSAL, 0082504), from *Lonchura castaneothorax*, Mount Hart, Western Australia, Australia ($17^{\circ}01'12"S, 125^{\circ}04'12"E$).

Harypalpus pyrrhula Bochkov & Klompen, 2014b: 463—Holotype female (OSAL, 0082535), from *Pyrrhula pyrrhula*, Nijmegen, Gelderland, The Netherlands ($51^{\circ}49'60"N, 5^{\circ}52'00"E$).

Harypalpus sturnus Bochkov & Klompen, 2014b: 464—Holotype female (OSAL, 0082907), from *Sturnus vulgaris*, Nijmegen, Gelderland, The Netherlands ($51^{\circ}49'60"N, 5^{\circ}52'00"E$).

Harypalpus taeniopygia Bochkov & Klompen, 2014b: 462—Holotype female (OSAL, 0082531), from *Taeniopygia guttata*, Brooking Springs, Western Australia, Australia ($18^{\circ}07'12"S, 125^{\circ}34'48"E$).

Harpyrhynchoides actitis Bochkov, OConnor & Klompen, 2015: 23—Holotype female (OSAL, 0077502), from *Actitis macularius*, Vancouver, British Columbia, Canada ($49^{\circ}09'36"N, 122^{\circ}51'36"W$).

- Harpyrhynchoides athene* Bochkov, OConnor & Klompen, 2015: 57**—Holotype female (OSAL, 0082592), from *Athene canicularia*, Lubbock, Lubbock Co., Texas, USA ($33^{\circ}34'48''N$, $101^{\circ}51'36''W$).
- Harpyrhynchoides bubulcus* Bochkov, OConnor & Klompen, 2015: 29**—Holotype female (UMMZ, BMOC 89-0315-001, #1), from *Bubulcus ibis*, Lake Adaire, Orlando, Orange Co., Florida, USA ($28^{\circ}33'60''N$, $81^{\circ}15'39''W$).
- Harpyrhynchoides calidris* Bochkov, OConnor & Klompen, 2015: 20**—Holotype female (OSAL, 0079988), from *Calidris minutilla*, Lawrence, Douglas Co., Kansas, USA ($38^{\circ}58'12''N$, $95^{\circ}14'24''W$).
- Harpyrhynchoides charadrius* Bochkov, OConnor & Klompen, 2015: 23**—Holotype female (OSAL, 0077525), from *Charadrius vociferus*, Lubbock Co., Texas, USA.
- Harpyrhynchoides coccyzus* Bochkov, OConnor & Klompen, 2015: 46**—Holotype female (UMMZ, BMOC 06-0518-001, #1 5), from *Coccyzus americanus*, UM Campus, Ann Arbor, Washtenaw Co., Michigan, USA ($42^{\circ}16'46''N$, $83^{\circ}44'19''W$).
- Harpyrhynchoides crotophaga* Bochkov, OConnor & Klompen, 2015: 63**—Holotype female (OSAL, 0082454), from *Crotophaga ani*, Weg naar Zee, Paramaribo, Suriname ($05^{\circ}49'12''N$, $55^{\circ}10'12''W$).
- Harpyrhynchoides gallowayi* Bochkov, OConnor & Klompen, 2015: 52**—Holotype female (ZISP, AVB 13-0325-004, #1), from *Columba livia*, Winnipeg, Manitoba, Canada.
- Harpyrhynchoides ixobrychus* Bochkov, OConnor & Klompen, 2015: 32**—Holotype female (UMMZ, BMOC 81-0605-006), from *Ixobrychus exilis*, Oakland Co., Michigan, USA.
- Harpyrhynchoides megascops* Bochkov, OConnor & Klompen, 2015: 42**—Holotype female (UMMZ, BMOC 09-0207-001, #1), from *Megascops asio*, Marble Ct., Commerce Twp., Oakland Co., Michigan, USA ($42^{\circ}33'52''N$, $83^{\circ}27'49''W$).
- Harpyrhynchoides pluvialis* Bochkov, OConnor & Klompen, 2015: 26**—Holotype female (OSAL, 0069439), from *Pluvialis dominica*, Wooster, Wayne Co., Ohio, USA ($40^{\circ}49'48''N$, $81^{\circ}52'48''W$).
- Harpyrhynchoides puffinus* Mertins, 2015 in Bochkov, OConnor & Klompen, 2015: 34**—Holotype female (UMMZ, BMOC 15-0302-001, #1), from *Puffinus gravis*, Atlantic Ocean adjacent to east cost of Martin Co., Florida, USA.
- Harpyrhynchoides setophaga* Bochkov& Klompen, 2014c: 406**—Holotype female (ZISP H-T-9; ZISP AVB 14-0725-001), from *Setophaga ruticilla*, Manitoba, Winnipeg, Canada.
- Harpyrhynchoides spizella* Bochkov& Klompen, 2014c: 415**—Holotype female (OSAL, 0077831), from *Spizella passerina*, Hopland Field Station, Mendocino Co., California, USA ($39^{\circ}00'00''N$, $123^{\circ}04'48''W$).
- Harpyrhynchoides xanthocephalus* Bochkov& Klompen, 2014c: 414**—Holotype female (OSAL, 0083369), from *Xanthocephalus xanthocephalus*, Valentine, Cherry Co., Nebraska, USA ($42^{\circ}52'12''N$, $100^{\circ}33'00''W$).
- Harpyrhynchoides zenaida* Bochkov, OConnor & Klompen, 2015: 44**—Holotype female (UMMZ, BMOC 07-0503-005, #1), from *Zenaida macroura*, UM Campus, Ann Arbor, Washtenaw Co., Michigan, USA ($42^{\circ}16'46''N$, $83^{\circ}44'19''W$).
- Neharpyrhynchus agelaius* Bochkov, OConnor & Klompen, 2015: 103**—Holotype female (UMMZ, BMOC 09-0701-022, #1), from *Agelaius phoeniceus*, Oakland Co., Michigan, USA.
- Neharpyrhynchus bombycilla* Bochkov, OConnor & Klompen, 2015: 98**—Holotype female (UMMZ, BMOC 08-01110-002, #1), from *Bombycilla cedrorum*, US route 31, Muskegon State game Area, Muskegon Co., Michigan, USA ($43^{\circ}17'46''N$, $86^{\circ}04'01''W$).
- Neharpyrhynchus campylorhynchus* Bochkov, OConnor & Klompen, 2015: 88**—Holotype female (OSAL, 0083261), from *Campylorhynchus brunneicapillus*, USA.
- Neharpyrhynchus picidarum* Bochkov, OConnor & Klompen, 2015: 90**—Holotype female (UMMZ, BMOC 07-0409-003, #1), from *Colaptes auratus*, Pontiac, Oakland Co., Michigan, USA ($42^{\circ}36'40''N$, $83^{\circ}17'16''W$).
- Neharpyrhynchus quiscula* Bochkov, OConnor & Klompen, 2015: 99**—Holotype female (UMMZ, BMOC 09-0910-002, #1), from *Quiscalus quiscula*, 1.3 mi E Big Rapids, Mecosta Co., Michigan, USA ($43^{\circ}41'55''N$, $85^{\circ}27'27''W$).
- Neharpyrhynchus spizella* Bochkov, OConnor & Klompen, 2015: 106**—Holotype female (UMMZ, BMOC 11-0317-002, #1), from *Spizella passerine*, Whitefish Point Bird Observatory, Chippewa Co., Michigan, USA ($46^{\circ}39'04''N$, $85^{\circ}02'20''W$).

Neharpyrhynchus turdus Bochkov, OConnor & Klompen, 2015: 93—Holotype female (UMMZ, BMOC 10-0316-001, #1), from *Turdus migratorius*, 4.4 mi ESE Saline, Washtenaw Co., Michigan, USA (42°09'22"N, 83°43'05"W).

Neharpyrhynchus vireo Bochkov, OConnor & Klompen, 2015: 104—Holotype female (OSAL, 0079454), from *Vireo olivaceus*, Leon Co., Florida, USA (30°29'24"N, 84°12'36"W).

Perharpyrhynchus caprimulgus Bochkov & Klompen, 2014a: 313—Holotype female (OSAL, 0083703), from *Caprimulgus fossii*, Mozambique (19°52'48"S, 34°36'00"E).

Perharpyrhynchus charadrius Bochkov, OConnor & Klompen, 2015: 81—Holotype female (UMMZ, BMOC 09-0408-001, #1), from *Charadrius vociferous*, UM campus, Ann Arbor, Washtenaw Co., Michigan, USA (42°16'46"N, 83°44'19"W).

Perhapyrhynchus elseyornis Bochkov & Klompen, 2014a: 315—Holotype female (OSAL, 0083216), from *Elseyornis melanops*, Brooking Springs, Western Australia, Australia (18°07'12"S, 125°34'48"E).

Trichorhynchiella myiarchus Bochkov & Klompen, 2014a: 305—Holotype female (OSAL, 0083898), from *Myiarchus crinitus*, an unknown locality, North America.

Family Syringophilidae (32 species)

Apodisyringiana hirundapi Skorack, Kaszewska & Kavetska, 2015: 584—Holotype female (AMU), from *Hirundapus caudacutus*, Yokohama, Japan.

Aulonastus pteroptochos Sikora, Kaszewska & Skoracki, 2014: 425—Holotype female (AMU), from *Pteroptochos tarnii*, Valdivia, Chile (39°49'S, 73°14'W).

Gunabopicobia masalaje Kaszewska, Kavetska & Skoracki, 2014: 294—Holotype female (AMU, Reg. No. AMU-SYR.489), from quill of body feathers of *Ptilinopus iozonus*, Papua New Guinea.

Krantziaulonastus dubininii Skoracki & Sikora, 2014: 198—Holotype female (ZISP, Reg. No. ZISP-AVB-011-2908-069), from *Passer domesticus*, Suvorov, Tula district, Russia.

Megasyringophilus cacatua Glowska & Laniecka, 2013: 390—Holotype female (ZIZMUH), from *Cacatua galerita*, Mount Hart, Australia.

Neoaulonastus apalis Klimovičová, Mikula, Kahure & Hromada, 2014: 573—Holotype female (AMU, Reg. No. AMU-SYR.474), from *Apalis porphyrolaema*, Nyandarua, Kenya (0°44'58.84"S, 36°40'56.68"E).

Neoaulonastus granatina Skoracki, Hromada & Unsoeld, 2013: 376—Holotype female (AMU, Reg. No. AMU-SYR.383), from quill of body feathers of *Granatina ianthinogaster*, Simangiro, Tanzania.

Neoaulonastus quelea Skoracki, Hromada & Unsoeld, 2013: 372—Holotype female (AMU, Reg. No. AMU-SYR.386), from quills of body feathers of *Quelea quelea*, Usa-River, Arusha, Tanzania.

Neoaulonastus tanzanicus Skoracki, Hromada & Unsoeld, 2013: 369—Holotype female (AMU, Reg. No. AMU-SYR.385), from quills of body feathers of *Euplectes axillaris*, Usa-River, Arusha, Tanzania.

Neosyringophilopsis tybidus Skoracki, Klimovičová, Muchai & Hromada, 2014: 184—Holotype female (AMU, Reg. No. AMU-SYR.444), from quill of wing-covert feathers of *Pogoniulus bilineatus*, W. Naro Moru, Ewaso Ngiro River, Kenya.

Peristerophila lature Kaszewska, Kavetska & Skoracki, 2014: 297—Holotype female (AMU, Reg. No. AMU-SYR.503), from *Ducula luctuosa*, Sulawesi Island, Indonesia.

Peristerophila upupi Klimovičová, Mikula, Kahure & Hromada, 2014: 575—Holotype female (AMU, Reg. No. AMU-SYR.476), from *Upupa epops*, Nyahururu, Kenya (0°0'37.69"S, 36°22'48.89"E).

Phipicobia pygiptilae Glowska & Schmidt, 2014b: 196—Holotype female (USNM), from quills of body feathers of *Pygiptila stellaris*, Upper Rewa River, Upper Takutu-Upper Essequibo, Guyana.

Picineoaulonastus pogoniulus Skoracki, Klimovičová, Muchai & Hromada, 2014: 180—Holotype female (AMU, Reg. No. AMU-SYR.442), from quill of wing covert of *Pogoniulus bilineatus*, North Nandi Forest, Kenya.

Picobia myiopagi Glowska & Milensky, 2014: 381—Holotype female (USNM), from quills of body feathers of *Myiopagis gaimardi*, Upper Rewa River, Upper Takutu-Upper Essequibo, Guyana.

Picobia ochoi Glowska & Milensky, 2014: 374—Holotype female (USNM), from quills of body feathers of *Tolmomyias sulphurescens*, ca. 19 km N, Mt. Holitipu, Paruima, Cuyuni-Mazaruni, Guyana.

Picobia passeri Skoracki & Sikora, 2014: 195—Holotype female (ZISP, Reg. No. ZISP-AVB-011-2908-068), from *Passer domesticus*, Suvorov, Tula district, Russia.

Picobia sayornis Glowska, 2014: 140—Holotype female (USNM), from quill of body feathers of the black phoebe *Sayornis nigricans*, El Caicho Biological Station, Tumbes, Peru.

Picobia schmidti Glowska & Milensky, 2014: 378—Holotype female (USNM), from quills of body feathers of *Lophotriccus pileatus*, El Caicho Biological Station, Tumbes, Peru.

Pipicobia terpsiphoni Glowska & Schmidt, 2014a: 59—Holotype female (USNM), from quill of body feathers of black-headed paradise-flycatcher *Terpsiphone rufiventer*, NW corner of N'dogo Lagoon, Moukalaba-Doudou National Park, Ogooué, Maritime Province, Gabon.

Rafapicobia milenskyi Glowska & Schmidt, 2014b: 200—Holotype female (USNM), from quill of body feathers of *Conopophaga aurite*, Upper Rewa River, Upper Takutu-Upper Essequibo, Guyana.

Rafapicobia thamnophili Glowska & Schmidt, 2014b: 198—Holotype female (USNM), from quills of body feathers of *Thamnophilus insignis*, Paruima, Cuyuni-Mazaruni, Guyana.

Syringophiloidus apus Skorack, Kaszewska & Kavetska, 2015: 585—Holotype female (AMU), from *Apus melba*, Valdivia, Chile.

Syringophiloidus furthi Glowska & Schmidt, 2014a: 61—Holotype female (USNM), from coverts of black-headed paradise-flycatcher *Terpsiphone rufiventer*, Ogooué Maritime Province, NW corner of N'dogo Lagoon, Moukalaba-Doudou National Park, Gabon.

Syringophilopsis pari Skoracki & Mironov, 2013: 559—Holotype female (ZISP–AVB 011–2908–051), from secondary of *Poecile palustris*, Novolitovsk, Partizanskii District, Primorsky Krai, Russia (42°51'40"N, 132°53'5.5"E).

Syringophilopsis philemonis Glowska & Laniecka, 2013: 386—Holotype female (ZIZMUH), from *Philemon citreogularis*, Beagle Bay, Australia.

Syringophiloidus picidus Skoracki, Klimovičová, Muchai & Hromada, 2014: 184—Holotype female (AMU, Reg. No. AMU-SYR.445), from quill of secondary feather of *Dendropicos fuscescens*, Mutanda, Kenya.

Syringophiloidus quetzali Skoracki, Mironov & Unsoeld, 2013: 292—Holotype female (AMU), from secondary covert quill of *Pharomachrus mocinno*, Guatemala.

Syringophilopsis szeffleri Glowska, 2014: 142—Holotype female (USNM), from the scale-crested pygmy-tyrant *Lophotriccus pileatus*, El Caicho Biological Station, Tumbes, Peru.

Syringophiloidus teledromas Sikora, Kaszewska & Skoracki, 2014: 420—Holotype female (AMU), from *Teledromas fuscus*, Neuquén, Cipolletti, Rio Negro, Argentina (38°56'S, 67°59'W).

Syringophilopsis trogoni Skoracki, Mironov & Unsoeld, 2013: 295—Holotype female (UNAM), from *Trogon citreolus*, Chamela Biological Station of UNAM, Jalisco, Mexico (19°29'52"N, 105°2'33"W).

Torotroglia volgini Skoracki & Mironov, 2013: 562—Holotype female (ZISP–AVB 011–2908–052), from secondary of *Emberiza elegans*, Novolitovsk, Partizanskii District, Primorsky Krai, Russia (42°51'40"N, 132°53'5.5"E).

Superfamily Pterygosomatoidea (8 species)

Family Pterygosomatidae (8 species)

Callopistiella atacamensis Silva-de la Fuente, Casanueva & Moreno, 2015 in Fuente, Paredes-León, Casanueva, Escobar-Huerta & Salas, 2015: 73—Holotype female (MZUC, 40360), from *Callopistes maculatus*, Atacama Region, Caldera, Copiapó Province, Chile (26°56'S, 70°44'W).

Pterygosoma aqabensis Fajfer & Melnikov, 2014: 404—Holotype female (ZISP), from *Pseudotrapelus aqabensis*, Mecca, Saudi Arabia (21°22'N, 39°37'E).

Pterygosoma dhofarensis Fajfer & Melnikov, 2014: 409—Holotype female (ZISP), from *Pseudotrapelus dhofarensis*, Ayoon, Dhofar, Oman (17°15'N, 53°54'E).

Pterygosoma engai Fajfer, 2013: 272—Holotype female (ZMUC, R36435), from *Agama sankaranica*, Idanre, Ondo State, Nigeria.

Pterygosoma fragilis Fajfer, 2013: 260—Holotype female (ZMUC, R3669), from *Agama agama*, Bura Tana, Garissa County, Kenya.

Pterygosoma garissi Fajfer, 2013: 260—Holotype female (ZMUC, R3619), from *Agama agama*, Bura Tana, Garissa County, Kenya.

Pterygosoma indare, 2013: 278—Holotype female (ZMUC, R36424), from *Agama sankaranica*, Idanre, Ondo State, Nigeria.

Pterygosoma olape, 2013: 280—Holotype female (ZMUC, R36473), from *Agama sankaranica*, Idanre, Ondo State, Nigeria.

Superfamily **Raphignathoidea** (16 species)

Family **Camerobiidae** (5 species)

Neophyllobius dogani Khanjani & Hoseini, 2013: 522—Holotype female (BASU), from soil under *Astragalus* sp., Chaghlvandy, Lorrestan Province, Iran (33°46'09.79"N, 48°39'26.21"E).

Neophyllobius edwardi Khanjani & Hoseini, 2013: 514—Holotype female (BASU), from soil and rotten leaves of oak trees *Quercus brantii*, Sorkhi region, Koohmare, Fars Province, Iran (29°31'25"N, 52°12'14"E).

Neophyllobius lorestanicus Khanjani, Hoseini, Yazdanpanah & Masoudian, 2014: 450—Holotype female (BASU), from soil under of *Prunus domestica*, Sarcheshmahe village, Mahallate, Markazi Province, Iran (33°54'39.94"N, 50°27'10.15"E).

Neophyllobius ostovani Khanjani & Ahmad-Hoseini, 2014 in Khanjani, Hoseini, Yazdanpanah & Masoudian, 2014: 446—Holotype female (BASU), from soil and rotten leaves of oak trees, Sorkhi region, Koohmare, Fars Province, Iran (29°29'26"N, 52°11'1"E).

Neophyllobius seemani Khanjani & Hoseini, 2013: 518—Holotype female (BASU), from litter under oak trees, Nahavad, Hamedan Province, Iran (29°32'08"N, 52°13'02"E).

Family **Stigmeidae** (11 species)

Agistemus macrosetosus Stathakis & Kapaxidi, 2014 in Stathakis, Kapaxidi & Papadoulis, 2014: 168—Holotype female (AUA), from *Rubus* sp., Kaiafas Lake, Co. Elia, Greece.

Agistemus riograndensis Johann & Ferla, 2013 in Johann, Carvalho, Majolo & Ferla, 2013: 248—Holotype female (MCN, CAS-03), from *Vitis vinifera*, Alfrocheiro cultivar, Candiota, state of Rio Grande do Sul, Brazil.

Cheylostigmaeus tarae Khanjani, 2014 in Khanjani, Nasrollahi, Zamani & Fayaz, 2014: 371—Holotype male (BASU), from soil and litter under pear trees, *Pyrus communis*, Yaghob Abad village, Baneh vicinity, Kurdistan Province, Iran (36°02'N, 45°44"E).

Eustigmaeus tjumeniensis Khaustov & Tolstikov, 2014: 539—Holotype female (TUMZ, NO.A310514), from *Plagiomnium cuspidatum* and *Brachythecium salebrosum*, vicinities of lake Kuchak, Tyumen Province, Russia (57°21'18"N, 66°03'01"E).

Mediolata ronaldi Laniecka, 2013 in Ripka, Laniecka & Kazmierski, 2013: 42—Holotype deutonymph (AMU, AK H-128), from *Tilia tomentosa*, Borsod-Abaúj-Zemplén County, Széphalom, Hungary.

Paravillersia jamaliensis Khaustov, 2014: 67—Holotype female (TUMZ, EB030814), from mosses on fen, vicinity of settlement Katravozh, Priuralskiy Region, Yamalo-Nenets Autonomous Okrug, Russia (66°19'40"N, 66°05'60"E).

Stigmaeus delaramae Khanjani, 2014 in Khanjani, Nasrollahi, Zamani & Fayaz, 2014: 374—Holotype female (BASU), from soil and litter under pear trees, *Pyrus communis*, Baneh vicinity, Yaghob Abad village, Kurdistan Province, Iran (36°02'N, 45°44'E).

Stigmaeus kuznetsovi Khaustov, 2014: 241—Holotype female (TUMZ, AK090495), from the nest of *Microtus socialis*, Yalta mountain pasture, Crimea, Ukraine.

***Stigmaeus mitrofanovi* Khaustov, 2014: 241**—Holotype female (TUMZ, AK190897), from the galleries of bark beetle *Pityogenes bistridentatus*, under the bark of *Pinus pallasiana*, Yalta mountain pasture, Crimea, Ukraine.

***Stigmaeus silvestris* Khaustov, 2014: 246**—Holotype female (TUMZ, AK110201), from rotten log of pine, *Pinus pallasiana*, Nikita mountain pasture, Crimea, Ukraine.

***Zetzellia ampelae* Johann & Ferla, 2013 in Johann, Carvalho, Majolo & Ferla, 2013: 253**—Holotype female (MCN, CAS-12), from *Vitis vinifera*, Bordeaux cultivar, Boqueirão do Leão, state of Rio Grande do Sul, Brazil.

Superfamily **Tetranychoidea** (32 species)

Family **Allochaetophoridae** (1 species)

***Allochaetophora iranica* Mahdavi & Asadi, 2015b: 363**—Holotype female (SBUK), from *Phragmites australis*, Manujan (Rozeye Eram)-Kerman Province, Iran (27°19'N, 57°30'E).

Family **Tenuipalpidae** (26 species)

***Amblypalpus iraniensis* Farzan, Asadi & Ueckermann, 2013 in Farzan, Asadi, Ueckermann, Seeman & Beard, 2013: 56**—Holotype female (SBUK), from *Amygdalus scoparia*, Dalfard, Jiroft-Kerman Province, Iran (28°58'N, 57°37'E).

***Amblypalpus thymus* Farzan, Asadi & Ueckermann, 2013 in Farzan, Asadi, Ueckermann, Seeman & Beard, 2013: 62**—Holotype female (SBUK), from *Thymus vulgaris*, Kuhpayeh, Kerman Province, Iran.

***Brevipalpus azores* Beard & Ochoa, 2015 in Beard, Ochoa, Braswell & Bauchan, 2015: 7**—Holotype female (USNM, #75-8887), from lemon, Azores, Portugal.

***Brevipalpus feresi* Ochoa & Beard, 2015 in Beard, Ochoa, Braswell & Bauchan, 2015: 16**—Holotype female (PEU), from *Croton floribundus*, São Paulo, Rio Petro, Brazil.

***Brevipalpus ferraguti* Ochoa & Beard, 2015 in Beard, Ochoa, Braswell & Bauchan, 2015: 21**—Holotype female (MNCN), from “Ngaio”, *Myoporum laetum*, Valencia, Spain.

***Brevipalpus noranae* Halawa & Fawzy, 2014: 88**—Holotype female (ARC-PPRI), from *Malus domestica* and *Citrus aurantium* as mixed plantations, in Moshtohor village, Tukh district, Qalyubia Province, Egypt (30°21'18"N, 31°13'30"E).

***Brevipalpus tucuman* Beard & Ochoa, 2015 in Beard, Ochoa, Braswell & Bauchan, 2015: 43**—Holotype female (USNM), from *Citrus sinensis* and *C. latifolia*, Argentina.

***Chaudhripalpus costacola* Beard & Seeman, 2014 in Beard, Seeman & Bauchan, 2014: 15**—Holotype female (QM), from *Casuarina cristata*, Newell Highway, approx. 15 km N Moree, 86 km S Goondiwindi, New South Wales, Australia (29°21'20"S, 150°00'24"E).

***Crossipalpus gersoni* Beard & Seeman, 2014 in Beard, Seeman & Bauchan, 2014: 36**—Holotype female (QM), from stems of *Allocasuarina luehmannii*, old Marlborough-Sarina Road, Mount Slopeaway, Queensland, Australia (22°52'16"S, 149°50'28"E).

***Crossipalpus raveni* Beard & Seeman, 2014 in Beard, Seeman & Bauchan, 2014: 44**—Holotype female (QM), from *Allocasuarina inophloia*, 90 km W Moonie, Moonie Highway, Queensland, Australia (27°56'57"S, 149°31'19"E).

***Magdalenapalpus caperatus* Beard & Seeman, 2014 in Beard, Seeman & Bauchan, 2014: 56**—Holotype female (MAGNT), from *Casuarina equisetifolia*, Nightcliff Boat Ramp, Darwin, Northern Territory, Australia (12°22'46"S, 130°50'34"E).

***Magdalenapalpus forsteri* Seeman & Beard, 2014 in Beard, Seeman & Bauchan, 2014: 62**—Holotype female (QM), from *Allocasuarina torulosa*, “Toondahra”, base of Mt Lorna, SSW Mundubbera, Australia (25°59'15"S, 151°21'26"E).

***Palpipalpus hesperius* Beard & Seeman, 2014 in Beard, Seeman & Bauchan, 2014: 81**—Holotype female (WAM), from *Allocasuarina* sp., Walpole Yacht Club, Coalmine Beach, Nornalup Inlet, Walpole, Western Australia, Australia (34°59'27"S, 116°44'22"E).

***Pentamerismus hicklingorum* Seeman & Beard, 2014 in Beard, Seeman & Bauchan, 2014: 104**—Holotype female (QM), from *Casuarina equisetifolia*, Bayside Drive, Beachmere, Southeast Queensland, Australia (27°05'52"S, 153°05'21"E).

Pentamerismus sititoris Beard & Seeman, 2014 *in* Beard, Seeman & Bauchan, 2014: 89—Holotype female (QM), from *Casuarina cristata*, approx. 15 km N Moree, 86 km S Goondiwindi, Newell Highway, New South Wales, Australia (29°21'20"S, 150°00'24"E).

Pentamerismus wardo Seeman & Beard, 2014 *in* Beard, Seeman & Bauchan, 2014: 97—Holotype female (WAM, BRI voucher, BRI [AQ814925]), from *Allocasuarina scleroclada*, approx. 3 km N Coomberdale, along Moora-Watheroo Road, Western Australia, Australia (30°24'17"S, 116°02'29"E).

Philippipalpus belah Beard & Seeman, 2014 *in* Beard, Seeman & Bauchan, 2014: 121—Holotype female (QM), from *Casuarina cristata*, approx. 15 km N Moree, 86 km S Goondiwindi, Newell Highway, New South Wales, Australia (29°21'20"S, 150°00'24"E).

Philippipalpus flumaquercus Beard & Seeman, 2014 *in* Beard, Seeman & Bauchan, 2014: 116—Holotype female (QM), from *Casuarina cunninghamiana*, Didcot Creek, 3 km ENE Didcot, nr Biggenden, Queensland, Australia (25°27'54"S, 151°53'47"E).

Philippipalpus nigraquercus Seeman & Beard, 2014 *in* Beard, Seeman & Bauchan, 2014: 124—Holotype female (QM), from *Allocasuarina littoralis*, Mugga Lane, Canberra, Symonston, Australian Capital Territory, Australia (35°23'05"S, 149°08'10"E).

Prolixus meyerae Xu & Zhang, 2014: 2—Holotype female (NZAC), from Alice Eaves Reserve, Orewa, Auckland, New Zealand.

Tenuipalpus apichai Castro & Feres, 2013: 483—Holotype female (PEU, NO. 8942), from *Celtis iguanae*, São José do Rio Preto, São Paulo State, Brazil (20°46"S, 49°15"W).

Tenuipalpus bacuri Flechtmann & Noronha, 2013: 496—Holotype female (ESALQ), from *Platonia insignis*, Embrapa-Amazonia Oriental, Belém, Pará, Brazil.

Tenuipalpus nambii Castro & Feres, 2013: 485—Holotype female (PEU, NO. 8947), from *Sterculia striata*, São João de Iracema, São Paulo State, Brazil (20°28"S, 50°17"W).

Terminalichus sanya Xu & Fan, 2014 *in* Xu, Fan & Huang, 2014: 235—Holotype female (NZMC), from *Terminalia catappa*, Egret Park, Fenghuang Road, Sanya city, Hainan Province, China (18°14'54.12"N, 109°30'57.82"E).

Tenuipalpus tapiae Castro & Feres, 2013: 483—Holotype female (PEU, NO. 8935), from *Alchornea glandulosa*, Estação Ecológica de Paulo de Faria, Paulo de Faria, São Paulo State, Brazil (19°55"S, 49°31"W).

Ultratenuipalpus avarua Xu, Fan & Zhang, 2013: 224—Holotype female (NZAC), from *Cocos* sp., Avarua, Rarotonga, Cook Islands, New Zealand.

Family Tetranychidae (5 species)

Petrobia (Tetranychina) hispaniola Sánchez & Flechtmann, 2014 *in* Martínez, Flechtmann & De Moraes, 2014: 554—Holotype female (ESALQ/USP), from *Citrus sinensis*, Río Seco, La Vega, Dominican Republic (Caribbean).

Schizotetranychus krungthepensis Naing & Auger, 2014 *in* Naing, Chandrapatya, Navajas & Auger, 2014: 258—Holotype male (KUB), from *Saccharum officinarum*, Ram Intra, Bangkok, Thailand.

Sonotetranychus menziesii Yi & Zhang, 2013: 335—Holotype female (NZAC, 03-359 Z-A), from *Nothofagus menziesii*, Golden Bay, Mt Stevens Takaka, New Zealand.

Tetranychus (Tetranychus) occultaspina Naing & Auger, 2014 *in* Naing, Chandrapatya, Navajas & Auger, 2014: 263—Holotype male (KUB), from *Ipomoea aquatic*, Kasetsart University campus, Khet Bangkhen, Bangkok, Thailand.

Tetranychus (Tetranychus) truncatissimus Naing & Auger, 2014 *in* Naing, Chandrapatya, Navajas & Auger, 2014: 269—Holotype male (KUB), from *Bambusa multiplex*, Khet Chatuchak, Bangkok, Thailand.

Hypoder Heterostigmata (20 species)

Superfamily Pygmephorooidea (16 species)

Family Pygmephoridae (6 species)

Pediculaster ermilovi Khaustov, 2015g: 419—Holotype female (TUMZ, AK151114), from the rotten log of poplar (*Populus nigra*), Tyumen, Tyumen Province, Russia (57°09'52.3"N, 65°31'01.6"E).

Pediculaster lignarius Khaustov, 2015g: 425—Holotype female (TUMZ, AK170514), from the rotten log of birch (*Betula pendula*), vicinity of Tyumen, Tyumen Province, Russia (57°09'50.5"N, 65°27'24.2"E).

Pseudopygmephorellus mazandaranicus Katlav & Hajiqanbar, 2015 in Katlav, Hajiqanbar & Talebi, 2015: 106—Holotype female (TUMZ, AK-20130318-H), from beetle *Onthophagus* sp., cow-pads in Dohezar Forest, Tonekabon town, west of Mazandaran Province, Iran (36°39'N, 50°40'E).

Pseudopygmephorellus troxi Khaustov & Trach, 2014: 92—Holotype female (NBG, VT220890/P), from *Trox cadaverinus*, Bamburovo vill., ca. 50 km SWW Vladivostok, ca 10 km NW Slavyanka, Primorye, Far East, Russia.

Zambedania argentiniana Camerik & Magowsk, 2014: 89—Holotype female (ZIZMUH), from *Stenoterommata platensis*, Parque Urugua-I, Prov. Misiones 16, Argentina.

Zambedania madagascariana Camerik & Magowsk, 2014: 88—Holotype female (NHMG), from Sainte Marie, Madagascar.

Family Scutacaridae (7 species)

Scutacarus crinitus Khaustov, 2015d: 274—Holotype female (TUMZ, AK170914), from ants *Lasius niger*, vicinity of Tyumen, Tyumen Province, Russia (57°13'43.6"N, 65°28'28.4"E).

Scutacarus heterotrichus Khaustov, 2015a: 355—Holotype female (TUMZ, AK150515), from the nest of *Lasius flavus*, vicinity of lake Kuchak, Tyumen Province, Russia (57°20'13.9"N, 66°03'30.4"E).

Scutacarus insolitus Khaustov, 2015a: 352—Holotype female (TUMZ, AK240815/1), from the nest of *Lasius flavus*, vicinity of Tyumen, Tyumen District, Tyumen Province, Russia (57°09'50.4"N, 65°27'26.7"E).

Scutacarus lasiophilus Khaustov, 2015d: 267—Holotype female (TUMZ, VS300714), from ants *Lasius niger*, vicinity of settlement Narimanovo, Tyumen region, Tyumen Province, Russia (57°21'56"N, 65°08'21"E).

Scutacarus moseri Khaustov, 2015a: 350—Holotype female (TUMZ, AK240815), from the nest of *Lasius flavus*, vicinity of Tyumen, Tyumen District, Tyumen Province, Russia (57°09'50.4"N, 65°27'26.7"E).

Scutacarus myrmicinus Khaustov, 2015d: 272—Holotype female (TUMZ, VS300714A), from the nest of *Myrmica ruginodis*, vicinity of settlement Narimanovo, Tyumen region, Tyumen Province, Russia (57°21'56"N, 65°08'21"E).

Scutacarus sibiricensis Khaustov, 2015a: 358—Holotype female (TUMZ, VS030814), from the nest of *Lasius flavus*, vicinity of settlement Kazanskoye, Kazanskiy District, Tyumen Province, Russia (55°37'21.1"N, 69°22' 17.7"E).

Family Neopygmephoridae (3 species)

Petalomium crossi Khaustov, 2015f: 554—Holotype female (TUMZ, NO. AP051214), from ant *Myrmicaria distincta*, coast of lake Tana, territory of the Bahir Dar Fisheries and other Aquatic Life Research Center, Bahir Dar city, Amhara Region, Ethiopia (11°36.70'N, 037°22.36'E).

Petalomium reductus Khaustov, 2015f: 555—Holotype female (TUMZ, NO. AP051214/1), from ants *Myrmicaria distincta*, coast of lake Tana, territory of the Bahir Dar Fisheries and other Aquatic Life Research Center, Bahir-Dar city, Amhara Region, Ethiopia (11°36.70'N, 037°22.36'E).

Troxodania magnifica Khaustov & Trach, 2014: 89—Holotype female (NBG, NO.VT220890), from *Trox cadaverinus*, Bamburovo vill., ca. 50 km SWW Vladivostok, ca. 10 km NW Slavyanka, Primorye, Far East, Russia.

Superfamily Tarsocheyloidea (1 species)

Family Tarsocheylidae (1 species)

Hoplocheylus magnificus Khaustov, 2015e: 286—Holotype female (ZISP, AY051014), from the nest of the ant *Formica rufa*, spruce forest, vicinity of settlement Shatrovo, Shatrovskiy district, Kurgan Province, Russia (56°30'1.95"N, 64°38'53.76"E).

Superfamily Tarsonemoidea (3 species)

Family Tarsonemidae (3 species)

Daidalotarsonemus annonae Sousa, Lofego & Gondim, 2014: 430—Holotype female (UNESP, NO. 9226), from *Annona coriacea*, Goiana, State of Pernambuco, Brazil (7°38'16.08"S, 34°57'9.41"W).

Fungitarsonemus setillus Sousa, Lofego & Gondim, 2014: 438—Holotype female (UNESP, NO. 9228), from *Annona muricata*, Goiana, State of Pernambuco, Brazil (7°38'16.08"S, 34°57'9.41"W).

Kaliszewskia ochoai Lofego, Demite & de Moraes, 2015: 567—Holotype female (PEU, NO. 9515), from *Plinia* sp., Serra do Conduru, Uruçuca, Bahia State, Brazil.

Acknowledgments

We thank Wendy Lam (University of Auckland) and Editor Dr Qing-Hai Fan (Ministry for Primary Industries, Auckland, New Zealand) for reviewing this manuscript and providing comments. Z.-Q. Zhang's research on New Zealand mites was supported mainly by Core Funding for Crown Research Institutes from the Ministry of Business, Innovation and Employment's Science and Innovation Group.

References

- Adil, S., Sevsay, S., Dogan, S. & Dilkaraoglu, S. (2015) *Milandanielia harsitensis* (Acari: Microtrombidiidae), a new species from Turkey. *Systematic & Applied Acarology*, 20, 647–654.
<http://dx.doi.org/10.11158/saa.20.6.7>
- Akbari, A., Irani-Nejad, K.H., Khanjani, M., Arzanlou, M. & Kazmierski, A. (2015a) A new tydeid species (Acari: Tydeidae) with a key to *Brachytydeus* species from East Azerbaijan Province, Iran. *Systematic & Applied Acarology*, 20, 423–430.
<http://dx.doi.org/10.11158/saa.20.4.7>
- Akbari, A., Irani-Nejad, K.H., Khanjani, M., Arzanlou, M. & kažmierski, A. (2015b) *Tydeus shabestariensis* sp. nov. and description of the male of *Neopronematus sepasgosariani* (Acari: Tydeoidea), with a key to the Iranian species of *Tydeus*. *Zootaxa*, 4032 (3), 264–276.
<http://dx.doi.org/10.11646/zootaxa.4032.3.2>
- Alatawi, F.J., Kamran, M. & Negm, M.W. (2015) False spider mites (Acari: Tenuipalpidae) associated with date palm orchards in Saudi Arabia: description of two new species and new records. *Systematic & Applied Acarology*, 20, 809–830.
<http://dx.doi.org/10.11158/saa.20.7.9>
- Bagheri, M. & Beyzavi, G. (2013) *Eustigmaeus ueckermannii*, a new species of the genus *Eustigmaeus* Berlese (Acari: Stigmaeidae) from central Iran. *Systematic & Applied Acarology*, 18, 30–34.
<http://dx.doi.org/10.11158/saa.18.1.3>
- Bagheri, M., Paktnat-Saejj, S., Saboori, A. & Ahaniazad, M. (2015) A new species of *Paraneognathus* from Iran and re-description of *Eustigmaeus plumifer* (Acari: Trombidiformes: Caligonellidae, Stigmaeidae). *Systematic & Applied Acarology*, 20, 329–338.
<http://dx.doi.org/10.11158/saa.20.3.10>
- Bagheri, M., Zarei, E., Ahaniazad, M., Gharekhany, G. & Navaei-Bonab, R. (2013) Two new species of the genus *Anoplochelylus* Berlese, 1910 (Acari: Trombidiformes: Pseudochelyidae) from Iran. *Zootaxa*, 3599 (3), 291–297.
<http://dx.doi.org/10.11646/zootaxa.3599.3.6>
- Beard, J.J., Ochoa, R., Braswell, W.E. & Bauchan, G.R. (2014) *Brevipalpus phoenicis* (Geijskes) species complex (Acari: Tenuipalpidae)—a closer look. *Zootaxa*, 3944 (1), 1–67.
<http://dx.doi.org/10.11646/zootaxa.3944.1.1>
- Beard, J.J., Seeman, O.D. & Bauchan, G.R. (2014) Tenuipalpidae (Acari: Trombidiformes) from Casuarinaceae (Fagales). *Zootaxa*, 3778 (1), 1–157.
<http://dx.doi.org/10.11646/zootaxa.3778.1.1>
- Bochkov, A.V. & Klompen, H. (2014a) New *Harpircynchinae* Dubinin (Acariformes: Harpirhynchidae)—intracutaneous and feather-base parasites of birds. *Zootaxa*, 3860 (4), 301–324.
<http://dx.doi.org/10.11646/zootaxa.3860.4.1>
- Bochkov, A.V. & Klompen, H. (2014b) A review of the subfamily *Harpypalpinae* Fain, 1972 (Acariformes: Harpirhynchidae)—parasites of passerine birds. *Zootaxa*, 3857 (4), 451–477.
<http://dx.doi.org/10.11646/zootaxa.3857.4.1>
- Bochkov, A.V. & Klompen, H. (2014c) A review of the *zumpti* species group of the genus *Harpyrhynchoides* (Acariformes: Harpirhynchidae)—ectoparasites of passerines. *Zootaxa*, 3884 (5), 401–418.
<http://dx.doi.org/10.11646/zootaxa.3884.5.1>

- Bochkov, A.V., OConnor, B.M. & Klompen, H. (2015) A review of the mite subfamily Harpirhynchinae (Acariformes: Harpirhynchidae)—parasites of New World birds (Aves: Neognathae). *Zootaxa*, 4023 (1), 1–130.
<http://dx.doi.org/10.11646/zootaxa.4023.1.1>
- Boyaci, Y.Ö. & Gölle, P. (2014) New records of the water mite family Hydryphantidae (Acari: Hydrachnidia) from Turkey, with the description of a new species. *Systematic & Applied Acarology*, 19, 160–165.
<http://dx.doi.org/10.11158/saa.19.2.6>
- Camerik, A.M. & Magowski, W.Ł. (2014) The genus *Zambedania* Mahunka 1972 (Acari: Heterostigmatina: Pygmephoridae)—redescription of the type species *Z. africana* and descriptions of two new species from Africa and South America. *Zootaxa*, 3793 (1), 71–98.
<http://dx.doi.org/10.11646/zootaxa.3793.1.3>
- Castro, E.B.D., Ramos, F. A., Feres, R.J. & Ochoa, R. (2015) A new species of *Tenuipalpus* Donnadieu (Acari: Tenuipalpidae) from Brazil, with ontogeny of chaetotaxy. *Systematic & Applied Acarology*, 20, 339–356.
<http://dx.doi.org/10.11158/saa.20.3.11>
- Chetverikov, P.E. (2015) Hidden diversity of endoparasitic eriophyoid mites: two new *Novophytoptus* Roivainen, 1947 (Acari: Eriophyoidea: Phytoptidae) species from the parenchymatous tissues of rushes (Juncaceae). *Zootaxa*, 4006 (3), 481–505.
<http://dx.doi.org/10.11646/zootaxa.4006.3.4>
- Chetverikov, P.E., Craemer, C., Vishnyakov, A.E. & Sukhareva, S.I. (2014) CLSM anatomy of internal genitalia of *Mackiella reclinata* n. sp. and systematic remarks on eriophyoid mites from the tribe Mackiellini Keifer, 1946 (Eriophyoidea, Phytoptidae). *Zootaxa*, 3860 (3), 261–279.
<http://dx.doi.org/10.11646/zootaxa.3860.3.5>
- Cichocka, M. & Biesiadka, E. (2013) Two new species of the genus *Atractides* Koch, 1837 (Acari: Hydrachnidia, Hygrobatidae) from Belarus. *Zootaxa*, 3608 (4), 253–263.
<http://dx.doi.org/10.11646/zootaxa.3608.4.3>
- Constantine, R.A. & Seeman, O.D. (2014) Three new species of *Eutarsopolipus* (Acari: Podapolipidae) from Australian carabid beetles (Coleoptera: Carabidae). *Systematic & Applied Acarology*, 19, 87–109.
<http://dx.doi.org/10.11158/saa.19.1.8>
- Darbemamieh, M., Hajiqanbar, H., Khanjani, M. & Kaźmierski, A. (2015) New species and records of *Neopronematus* (Acari: Iolinidae) from Iran with a key to world species. *Zootaxa*, 3990 (2), 235–246.
<http://dx.doi.org/10.11646/zootaxa.3990.2.4>
- De Castro, E.B. & Feres, R. (2013) New species of *Tenuipalpus* (Acari: Tenuipalpidae) from Semideciduous Forest remnants in the State of São Paulo, Brazil. *Zootaxa*, 3716 (3), 475–493.
<http://dx.doi.org/10.11646/zootaxa.3716.3.9>
- Denizhan, E., Monfreda, R., De Lillo, E. & Çobanoğlu, S. (2015) Eriophyoid mite fauna (Acari: Trombidiformes: Eriophyoidea) of Turkey: new species, new distribution reports and an updated catalogue. *Zootaxa*, 3991 (1), 1–63.
<http://dx.doi.org/10.11646/zootaxa.3991.1.1>
- Doğan, S., Dilkaraoglu, S., Fan, Q.-H., Erman, O., Sevsay, S. & Adil, S. (2015) Description of a new species of the genus *Eryngiopus* Summers (Acari: Stigmaeidae) from Turkey. *Systematic & Applied Acarology*, 20, 431–440.
<http://dx.doi.org/10.11158/saa.20.4.8>
- Dogan, S., Dilkaraoglu, S., Fan, Q.-H., Sevsay, S., Erman, O. & Adil, S. (2015) Description of a species of the genus *Cheylostigmaeus* Willmann (Acari: Stigmaeidae) from Eksisu Marsh, Turkey. *Systematic & Applied Acarology*, 20, 797–808.
<http://dx.doi.org/10.11158/saa.20.7.8>
- Eghbalian, A.H., Khanjani, M., Safaralizadeh, M.H. & Ueckermann, E.A. (2014) Two new species of *Cyta* (Acari: Prostigmata: Bdellidae) from Western Iran. *Zootaxa*, 3847 (4), 567–575.
<http://dx.doi.org/10.11646/zootaxa.3847.4.6>
- Esen, Y., Erman, O. & Dilkaraoglu, S. (2013) Contribution to the study of arrenuroid water mites (Acari: Hydrachnidia) from Turkey. *Zootaxa*, 3666 (1), 73–83.
<http://dx.doi.org/10.11646/zootaxa.3666.1.7>
- Esen, Y., Pešić, V. & Erman, O. (2013) Water mites of the genus *Brachypoda* (Acari: Hydrachnidia: Aturidae) in Turkey. *Zootaxa*, 3686 (3), 326–334.
<http://dx.doi.org/10.11646/zootaxa.3686.3.2>
- Fajfer, M. (2013) Three new species of scale mites (Acari: Pterygosomatidae) parasitizing *Agama sankaranica* (Sauria: Agamidae). *Zootaxa*, 3700 (2), 271–282.
<http://dx.doi.org/10.11646/zootaxa.3700.2.5>
- Fajfer, M. (2013) Two new species of scale mites (Acari: Pterygosomatidae) parasitizing *Agama agama* (Sauria: Agamidae) from Kenya. *Zootaxa*, 3745 (2), 257–262.
<http://dx.doi.org/10.11646/zootaxa.3745.2.5>
- Fajfer, M. & Melnikov, D. (2014) New species and records of scale mites (Acari: Pterygosomatidae) from Arabian agamid lizards (Squamata: Agamidae). *Zootaxa*, 3764 (4), 401–417.
<http://dx.doi.org/10.11646/zootaxa.3764.4.1>

- Fan, Q.H. (2006) Bibliographic analysis of acarological papers published in Zootaxa from 2001 to 2005, with a catalogue of described new taxa. *Zootaxa*, 1385, 53–66.
- Farzan, S. & Asadi, M. (2013) A new species of *Aegyptobia* Sayed, 1950 (Acari: Trombidiformes: Tenuipalpidae) from Iran. *Systematic & Applied Acarology*, 18, 35–40.
<http://dx.doi.org/10.11158/saa.18.1.4>
- Farzan, S. & Asadi, M. (2015) A new species of *Aegyptobia* Sayed, 1950 (Acari: Tenuipalpidae) from Iran. *Systematic & Applied Acarology*, 20, 955–960.
<http://dx.doi.org/10.11158/saa.20.8.10>
- Farzan, S., Asadi, M., Ueckermann, E.A., Seeman, O.D. & Beard, J.J. (2013) A review of *Amblypalpus* and *Priscapalpus* (Acari: Trombidiformes: Tenuipalpidae), including two new species of *Amblypalpus* from Iran. *Zootaxa*, 3716 (1), 53–64.
<http://dx.doi.org/10.11646/zootaxa.3716.1.4>
- Flechtmann, C.H., Ballari, M.C. & Quinteros, S.L.Q.D. (2014) Eriophyoidea (Acari) on Solanaceous plants from Argentina, with description of a new species of *Rhynacus* (Diptilomiopidae) and a key to species. *Systematic & Applied Acarology*, 19, 73–78.
<http://dx.doi.org/10.11158/saa.19.1.6>
- Flechtmann, C.H. & Noronha, A.C. (2013) A new species of the genus *Tenuipalpus* (Prostigmata: Tenuipalpidae) with remarks on a conceivable ovipositor in flat mites. *Zootaxa*, 3681 (4), 493–499.
<http://dx.doi.org/10.11646/zootaxa.3681.4.10>
- Flechtmann, C.H. & Queiroz, D.L.D. (2015) *Mononychellus* Wainstein, 1971 (Acari, Prostigmata, Tetranychidae): description of a new species from Brazil and key to species. *Systematic & Applied Acarology*, 20, 831–838.
<http://dx.doi.org/10.11158/saa.20.7.10>
- Fuangworn, M. & Butcher, B.A. (2015) *Neocaeculus orientalis* sp. nov. (Acari, Trombidiformes, Caeculidae) from Thailand. *Zootaxa*, 4048 (2), 251–268.
<http://dx.doi.org/10.11646/zootaxa.4048.2.6>
- Gaston, K.J. (2000) Global patterns in biodiversity. *Nature*, 405, 220–227.
- Gerecke, R. (2014) Studies on European species of the water mite family Aturidae Thor (Acari: Hydrachnidia). *Zootaxa*, 3841 (1), 1–46.
<http://dx.doi.org/10.11646/zootaxa.3841.1.1>
- Glowska, E. (2014) New quill mites (Cheyletoidea: Syringophilidae) parasitizing tyrannid birds (Passeriformes: Tyrannidae) in Peru. *Zootaxa*, 3814 (1), 139–145.
<http://dx.doi.org/10.11646/zootaxa.3814.1.9>
- Glowska, E. & Laniecka, I. (2013) Two new quill mite species (Prostigmata: Syringophilidae) parasitizing Australian birds. *Zootaxa*, 3670 (3), 385–390.
<http://dx.doi.org/10.11646/zootaxa.3670.3.9>
- Glowska, E. & Milensky, C.M. (2014) New species of the genus *Picobia* (Cheyletoidea: Syringophilidae) parasitizing tyrannid birds (Passeriformes: Tyrannidae). *Zootaxa*, 3821 (3), 373–383.
<http://dx.doi.org/10.11646/zootaxa.3821.3.6>
- Glowska, E. & Schmidt, B.K. (2014a) New quill mites (Cheyletoidea: Syringophilidae) parasitizing the black-headed paradise-flycatcher *Terpsiphone rufiventer* (Passeriformes: Monarchidae) in Gabon. *Zootaxa*, 3786 (1), 57–64.
<http://dx.doi.org/10.11646/zootaxa.3786.1.3>
- Glowska, E. & Schmidt, B.K. (2014b) New taxa of the subfamily Picobiinae (Cheyletoidea: Syringophilidae) parasitizing antbirds and gnatcatchers (Passeriformes: Thamnophilidae, Conopophagidae) in Guyana. *Zootaxa*, 3861 (2), 193–200.
<http://dx.doi.org/10.11646/zootaxa.3861.2.7>
- Gulle, P. & Boyaci, Y.Ö. (2014) A new *Protzia* species and the first description of the female *Atractides (Polymegapus) persicus* Pešić & Asadi 2010 from Turkey. *Zootaxa*, 3852 (5), 585–590.
<http://dx.doi.org/10.11646/zootaxa.3852.5.7>
- Gulle, P., Gulle, İ. & Boyaci, Y.Ö. (2015) A new *Atractides* Koch (Acari: Hydrachnidia, Hygrobatidae) species from southwestern Turkey. *Zootaxa*, 3957 (2), 246–248.
<http://dx.doi.org/10.11646/zootaxa.3957.2.11>
- Haitlinger, R. (2014) *Charletonia cameroonensis* Haitlinger & Kekeunou sp. nov. and the first record of *C. justyna* Haitlinger, 1987 (Acari: Erythraeidae) from Cameroon with redescription of the species. *Zootaxa*, 3760 (1), 39–53.
<http://dx.doi.org/10.11646/zootaxa.3760.1.2>
- Haitlinger, R. & Šundić, M. (2015) Two new species of *Calyptostoma cambridgei*, 1875 (Acari: Prostigmata: Calyptostomatidae) from Cambodia and Sulawesi, Indonesia. *Systematic & Applied Acarology*, 20, 919–926.
<http://dx.doi.org/10.11158/saa.20.8.7>
- Hakimitabar, M., Saboori, A. & Seiedy, M. (2013) A new species of larval *Charletonia* (Acari: Erythraeidae) parasitic on Arachnida from Iran. *Systematic & Applied Acarology*, 18, 163–176.
<http://dx.doi.org/10.11158/saa.18.2.9>
- Halawa, A.M. & Fawzy, M.M. (2014) A new species of *Brevipalpus* Donnadieu (Acari: Tenuipalpidae) and key to the Egyptian species. *Zootaxa*, 3755 (1), 87–95.

- http://dx.doi.org/10.11646/zootaxa.3755.1.4
- Han, X., Rajput, S., Xue, X.-F. & Hong, X.-Y. (2015) Four new species of *Tetra* and *Tetraspinus* (Acari: Eriophyidae) from China. *Systematic & Applied Acarology*, 20, 507–522.
<http://dx.doi.org/10.11158/saa.20.5.7>
- Han, X., Xue, X.-F. & Hong, X.-Y. (2013) Three new eriophyoid mite species in the tribe Phyllocoptini from Yunnan Province, southwestern China (Acari: Eriophyidae: Phyllocoptinae). *Zootaxa*, 3702 (3), 247–261.
<http://dx.doi.org/10.11646/zootaxa.3702.3.3>
- Hassanzadeh, M., Khanjani, M., Safaralizadeh, M.H. & Mirfakhraie, S. (2013) A new species of the genus *Storchia* Oudemans (Acari: Stigmeidae) from northwest Iran. *Systematic & Applied Acarology*, 18, 351–356.
<http://dx.doi.org/10.11158/saa.18.4.5>
- Husband, R.W. & Husband, D.O. (2013) *Eutarsopolipus americanus* sp. nov. (Acari: Podapolipidae), subelytral parasite of *Chlaenius praesinus* Dejean (Coleoptera: Carabidae) from Michigan and Missouri, USA. *Systematic & Applied Acarology*, 18, 53–60.
<http://dx.doi.org/10.11158/saa.18.1.6>
- Husband, R.W. & Husband, D.O. (2015) *Eutarsopolipus obrieni* n. sp. (Acari: Podapolipidae), ectoparasite of *Platynus darlingtoni* (Coleoptera: Carabidae) in Jamaica, a second species in the *brettae* group of *Eutarsopolipus*. *Systematic & Applied Acarology*, 20, 967–976.
<http://dx.doi.org/10.11158/saa.20.8.12>
- Husband, R.W. & Kurosa, K. (2013) *Eutarsopolipus asiaticus* sp. nov. (Acari: Podapolipidae), subelytral parasite of *Chlaenius costiger* Chaudoir (Coleoptera: Carabidae) from Japan. *Systematic & Applied Acarology*, 18, 61–70.
<http://dx.doi.org/10.11158/saa.18.1.7>
- Husband, R.W. & OConnor, B.M. (2014) A new genus and species of Podapolipidae (Acari: Heterostigmata) parasitic on *Physonota alutacea* (Boheman)(Coleoptera: Chrysomelidae; Cassidinae) in Mexico and Central America. *Systematic & Applied Acarology*, 19, 435–446.
<http://dx.doi.org/10.11158/saa.19.4.7>
- Jafari, S., Bagheri, M. & Saboori, A. (2014) A new species of the genus *Favognathus* from Iran and re-description of *Eryngiopus affinis* Barilo (Acari: Cryptognathidae, Stigmeidae). *Systematic & Applied Acarology*, 19, 353–362.
<http://dx.doi.org/10.11158/saa.19.3.8>
- Johann, L., Carvalho, G.S., Majolo, F. & Ferla, N.J. (2013) Stigmaeid mites (Acari: Stigmeidae) from vineyards in the state of Rio Grande do Sul, Brazil. *Zootaxa*, 3701 (2), 238–256.
<http://dx.doi.org/10.11646/zootaxa.3701.2.6>
- Kaluz, S., Ermilov, S.G. & Vrabec, M. (2014) Two new species of the genus *Armascirus* (Acari: Prostigmata: Cunaxidae) from India and Vietnam, with a description of the preimaginal stage of *Armascirus fendai*. *Zootaxa*, 3835 (2), 237–250.
<http://dx.doi.org/10.11646/zootaxa.3835.2.4>
- Kaluz, S. & Vrabec, M. (2013) Two new species of *Armascirus* (Acari: Prostigmata: Cunaxidae) from Slovakia. *Zootaxa*, 3734 (2), 141–155.
<http://dx.doi.org/10.11646/zootaxa.3734.2.3>
- Kamran, M. & Alatawi, F.J. (2015) New species and new records of subfamily Abrolophinae (Acari: Erythraeidae) from Saudi Arabia. *Systematic & Applied Acarology*, 20, 195–202.
<http://dx.doi.org/10.11158/saa.20.2.6>
- Kaszewska, K., Kavetska, K. & Skoracki, M. (2014) Two new species of quill mites of the family Syringophilidae (Acariformes: Cheyletoidea) associated with treronine doves (Columbiformes: Columbidae: Treroninae). *Zootaxa*, 3846 (2), 293–300.
<http://dx.doi.org/10.11646/zootaxa.3846.2.9>
- Katlav, A., Hajiqanbar, H. & Talebi, A.A. (2015) *Pseudopygmephorellus mazandaranicus* sp. nov. (Acari: Heterostigmata: Pygmephoridae), phoretic on scarabaeid dung beetles (Coleoptera: Scarabaeidae) from Iran. *Zootaxa*, 3919 (1), 100–110.
<http://dx.doi.org/10.11646/zootaxa.3919.1.5>
- Khanjani, M., Alvandy, S., Khanjani, M. & Seeman, O.D. (2013) A new species of *Tenuipalpus* Donnadiue (Acari: Tenuipalpidae) from Iran. *Systematic & Applied Acarology*, 18, 389–395.
<http://dx.doi.org/10.11158/saa.18.4.7>
- Khanjani, M., Firoozfar, A. & Mirmoayed, A. (2013) *Cheylostigmaeus mahvashae* sp. nov., a new species of the family Stigmeidae (Acari) from Kermanshah Province, Iran. *Systematic & Applied Acarology*, 18, 345–350.
<http://dx.doi.org/10.11158/saa.18.4.4>
- Khanjani, M. & Hoseini, M.A. (2013) Three new species of the genus *Neophylllobius* Berlese (Acari: Camerobiidae) from southern and southwestern Iran. *Zootaxa*, 3666 (4), 510–522.
<http://dx.doi.org/10.11646/zootaxa.3666.4.6>
- Khanjani, M., Hoseini, M.A. & Amini, F. (2014) Two new *Anoplocheylus* species (Acari: Trombidiformes: Pseudocheylelidae) from Kurdistan province of Iran. *Zootaxa*, 3861 (2), 185–192.
<http://dx.doi.org/10.11646/zootaxa.3861.2.6>
- Khanjani, M., Hoseini, M.A., Yazdanpanah, S. & Masoudian, F. (2014) *Neophylllobius lorestanicus* sp. nov. and *N. ostovani* sp.

- nov. (Acari: Camerobiidae) from Iran. *Zootaxa*, 3764 (4), 441–454.
<http://dx.doi.org/10.11646/zootaxa.3764.4.4>
- Khanjani, M., Khanjani, M. & Razmjou, J. (2015) A new species of the genus *Tetranychopsis* (Acari: Tetranychidae) from Ardabil Province, Iran. *Systematic & Applied Acarology*, 20, 110–120.
<http://dx.doi.org/10.11158/saa.20.1.10>
- Khanjani, M., Nadri, A.R., Khanjani, M. & Seeman, O.D. (2014) Post larval stages of *Tanytdeus beyzavii* sp. nov. (Acari: Paratydeidae) from Iran. *Zootaxa*, 3895 (2), 170–182.
<http://dx.doi.org/10.11646/zootaxa.3895.2.2>
- Khanjani, M., Nasrollahi, S., Zamani, A.S. & Fayaz, B.A. (2014) *Cheylostigmaeus tarae* sp. nov. and *Stigmaeus delaramae* sp. nov. (Acari: Stigmeidae) from Kurdistan, Iran. *Zootaxa*, 3841 (3), 364–378.
<http://dx.doi.org/10.11646/zootaxa.3841.3.3>
- Khanjani, M., Yazdanpanah, S. & Fayaz, B.A. (2013) *Austroteneriffia shiraziensis* sp. nov. (Acari: Teneriffiidae) from southwestern Iran, with description of male and immature stages. *Zootaxa*, 3683 (1), 35–50.
<http://dx.doi.org/10.11646/zootaxa.3683.1.2>
- Khaustov, A.A. (2014a) New species and new records of mites of the genus *Stigmaeus* (Acari: Prostigmata: Stigmeidae) from Crimea. *Zootaxa*, 3794, 237–253.
<http://dx.doi.org/10.11646/zootaxa.3794.2.3>
- Khaustov, A.A. (2014b) A new species of the genus *Paravillersia* (Acari: Prostigmata: Stigmeidae) from Western Siberia, with supplementary description of *Paravillersia grata* Kuznetsov, 1978. *Zootaxa*, 3873 (1), 62–72.
<http://dx.doi.org/10.11646/zootaxa.3873.1.5>
- Khaustov, A.A. (2015a) Myrmecophilous pygmephoroid mites (Acari: Pygmephoridae) associated with *Lasius flavus* (Hymenoptera: Formicidae) in Russia. *Zootaxa*, 4044 (3), 345–370.
<http://dx.doi.org/10.11646/zootaxa.4044.3.2>
- Khaustov, A.A. (2015b) A new genus of the family Eupodidae (Acari: Eupodoidea) with the description of a new species from Crimea. *Systematic & Applied Acarology*, 20, 927–936.
<http://dx.doi.org/10.11158/saa.20.8.8>
- Khaustov, A.A. (2015c) New species and new records of mites of the genus *Stigmaeus* (Acari: Prostigmata: Stigmeidae) from Western Siberia with redescription of *S. livschitzi* Kuznetsov, 1977. *Systematic & Applied Acarology*, 20, 681–692.
<http://dx.doi.org/10.11158/saa.20.6.10>
- Khaustov, A.A. (2015d) Three new species of myrmecophilous scutacarid mites (Acari: Scutacaridae) from Western Siberia, Russia. *Zootaxa*, 4013 (2), 265–279.
<http://dx.doi.org/10.11646/zootaxa.4013.2.7>
- Khaustov, A.A. (2015e) To systematics of the mite genus *Hoplocheylus* (Acariformes: Tarsocheylidae). *Zootaxa*, 3957 (3), 277–299.
<http://dx.doi.org/10.11646/zootaxa.3957.3.2>
- Khaustov, A.A. (2015f) Two new myrmecophilous species of the genus *Petalonium* (Acari: Pygmephoridae: Neopygmephoridae). *Zootaxa*, 3999 (4), 549–559.
<http://dx.doi.org/10.11646/zootaxa.3999.4.5>
- Khaustov, A.A. (2015g) Two new species of the genus *Pediculaster* (Acari: Pygmephoridae) from Western Siberia, Russia. *Zootaxa*, 3926 (3), 413–429.
<http://dx.doi.org/10.11646/zootaxa.3926.3.6>
- Khaustov, A.A. & Tolstikov, A.V. (2014) A new species and new records of the genus *Eustigmaeus* (Acari: Prostigmata: Stigmeidae) from Western Siberia. *Zootaxa*, 3861 (6), 531–553.
<http://dx.doi.org/10.11646/zootaxa.3861.6.2>
- Khaustov, A.A. & Trach, V.A. (2014) Mites of the superfamily Pygmephoridae (Acari: Heterostigmata: Neopygmephoridae, Pygmephoridae) associated with *Trox cadaverinus* (Coleoptera: Trogidae) from the Far East of Russia, with description of a new genus and two new species. *Zootaxa*, 3754 (1), 86–96.
<http://dx.doi.org/10.11646/zootaxa.3754.1.4>
- Klimovičová, M., Mikula, P., Kahure, N. & Hromada, M. (2014) A review of quill mites (Acari: Syringophilidae) parasitising Kenyan birds. *Zootaxa*, 3857 (4), 571–580.
<http://dx.doi.org/10.11646/zootaxa.3857.4.6>
- Konikiewicz, M. & Mąkol, J. (2014) A fossil Paratrombiinae mite (Actinotrichida: Trombidioidea) from the Rovno amber, Ukraine. *Zootaxa*, 3847 (4), 583–589.
<http://dx.doi.org/10.11646/zootaxa.3847.4.8>
- Konvipasruang, P. & Chandrapatya, A. (2015) Two new genera and four new species of Diptilomiopidae from Thailand. *Systematic & Applied Acarology*, 20, 297–312.
<http://dx.doi.org/10.11158/saa.20.3.7>
- Kurosa, K. & Husband, R.W. (2013) A new genus and species, *Simalurapolipus hiraii* (Acari: Heterostigmatina: Podapolipidae) parasitic on *Simalura coerulea* (Coleoptera: Tenebrionidae) in Japan. *Systematic & Applied Acarology*, 18, 252–262.
<http://dx.doi.org/10.11158/saa.18.3.8>

- Lam, W. & Zhang, Z.-Q. (2016) Hotspots of mite new species discovery: Parasitiformes (2013–2015). *Systematic & Applied Acarology*, 21 (12), 1693–1709.
<http://doi.org/10.11158/saa.21.12.10>
- Li, G.-Y. & Zhang, Z.-Q. (2016) Hotspots of mite new species discovery: Scarcoptiformes (2013–2015). *Zootaxa*, 4208 (2), 101–126.
<http://doi.org/10.11646/zootaxa.4208.2.1>
- Li, J.-W., Li, G.-Y., Zhang, J.-P. & Xue, X.-F. (2014) Five new species of the genus *Tetra* Keifer (Acari: Eriophyidae: Phyllocoptinae) from Xinjiang Uygur Autonomous Region, China. *Systematic & Applied Acarology*, 19, 329–352.
<http://dx.doi.org/10.11158/saa.19.3.7>
- Liu, D., Yi, T.-C., Xu, Y. & Zhang, Z.-Q. (2013) Hotspots of new species discovery: new mite species described during 2007 to 2012. *Zootaxa*, 3663 (1), 1–102.
<http://dx.doi.org/10.11646/zootaxa.3663.1.1>
- Liu, Z.G., Yi, T.C. & Guo, J.J. (2015) A new species of *Armascirus* (Acari, Prostigmata, Cunaxidae) from China. *Systematic & Applied Acarology*, 20, 322–328.
<http://dx.doi.org/10.11158/saa.20.3.9>
- Lofego, A.C., Demite, P.R. & Moraes, G.D. (2015) A new genus and species of Tarsonemidae (Acari: Heterostigmata) from the Atlantic Forest, Brazil. *Zootaxa*, 3986 (5), 561–568.
<http://dx.doi.org/10.11646/zootaxa.3986.5.3>
- Lotfollahi, P., Irani-Nejad, K.H. & De Lillo, E. (2014) Eriophyoid mites (Acari: Trombidiformes: Eriophyoidea) of Rosales trees in Iran: two new species and three new records. *Zootaxa*, 3861 (1), 76–85.
<http://dx.doi.org/10.11646/zootaxa.3861.1.4>
- Lotfollahi, P., Irani-Nejad, K.H., Khanjani, M., Moghadam, M. & De Lillo, E. (2013) Eriophyoid mites (Acari: Prostigmata: Eriophyidae) associated with Compositae in Iran. *Zootaxa*, 3664 (3), 349–360.
<http://dx.doi.org/10.11646/zootaxa.3664.3.5>
- Lotfollahi, P., Lillo, E.D. & Irani-Nejad, K.H. (2015) Three new *Aculops* species (Acari: Trombidiformes: Eriophyidae) from Iran. *Systematic & Applied Acarology*, 20, 410–422.
<http://dx.doi.org/10.11158/saa.20.4.6>
- Mahdavi, S.M. & Asadi, M. (2015a) A new species of *Schizotetranychus* (Acari, Tetranychidae) from Iran. *Systematic & Applied Acarology*, 20, 674–680.
<http://dx.doi.org/10.11158/saa.20.6.9>
- Mahdavi, S.M. & Asadi, M. (2015b) Third report of *Allochaetophoridae* Reck, 1959 (Acari: Trombidiformes) based on the description of a new species from Iran. *Zootaxa*, 4012 (2), 361–368.
<http://dx.doi.org/10.11646/zootaxa.4012.2.6>
- Mahdavi, S.M., Asadi, M. & Farzan, S. (2013) A new species of *Eupalopsellus* Sellnick, 1950 (Acari: Eupalopsellidae) from the south of Iran. *Systematic & Applied Acarology*, 18, 190–196.
<http://dx.doi.org/10.11158/saa.18.2.11>
- Mahdavi, S.M., Asadi, M., Ueckermann, E.A. & Farzan, S. (2013) A new species of *Tetranychus* Dufour, 1832 (Acari, Trombidiformes: Tetranychidae) from Iran. *Systematic & Applied Acarology*, 18, 245–251.
<http://dx.doi.org/10.11158/saa.18.3.7>
- Mahmoudi, F., Saboori, A., Sari, A. & Hakimitabar, M. (2014) A new species of larval *Erythraeus* (*Zaracarus*) (Acari: Trombidiformes: Erythraeidae) from Iran, with a key to the world species of the subgenus. *Systematic & Applied Acarology*, 19, 79–86.
<http://dx.doi.org/10.11158/saa.19.1.7>
- Makol, J. & Sevsay, S. (2014) The genus *Emitrombidium* Lombardini, 1949 (Actinotrichida: Trombidiidae) resurrected. *Zootaxa*, 3786 (1), 91–98.
<http://dx.doi.org/10.11646/zootaxa.3786.1.8>
- Martinez, L.S., Flechtmann, C.H. & De Moraes, G.J. (2014) Plant mites of the Dominican Republic, with a description of a new species of *Petrobia* (*Tetranychina*) Wainstein, 1960 (Acari, Prostigmata, Tetranychidae) and a key to the species of this subgenus. *Zootaxa*, 3846 (4), 547–560.
<http://dx.doi.org/10.11646/zootaxa.3846.4.3>
- Menon, P., Joshi, S. & Ramamurthy, V.V. (2014) A new species of eriophyoid mite, *Aceria tripuraensis* sp.n. (Acari: Eriophyoidea), on *Hibiscus macrophyllus* from India. *Zootaxa*, 3760 (4), 553–562.
<http://dx.doi.org/10.11646/zootaxa.3760.4.4>
- Mortazavi, A., Hajiqanbar, H. & Kamali, K. (2015) A new species of the family Dolichocybidae Mahunka, 1970 (Acari: Heterostigmata) associated with *Sinoxylon pugnax* Lesne (Insecta: Coleoptera: Bostrichidae) from Iran. *Systematic & Applied Acarology*, 20, 441–448.
<http://dx.doi.org/10.11158/saa.20.4.9>
- Naing, H.H., Chandrapatya, A., Navajas, M. & Auger, P. (2014) New species and new records of Tetranychidae (Acarina, Prostigmata) from Thailand. *Zootaxa*, 3802 (2), 257–275.
<http://dx.doi.org/10.11646/zootaxa.3802.2.7>
- Negm, M.W., Johann, L., Ferla, N.J. & Amano, H. (2015) A new species of *Zetzellia* Oudemans (Acari, Trombidiformes,

- Raphignathoidea, Stigmeidae) and a key to Stigmeidae of Japan. *Systematic & Applied Acarology*, 20, 961–966.
<http://dx.doi.org/10.11158/saa.20.8.11>
- Noei, J., Rabieh, M.M. & Saboori, A. (2015) First record of *Momorangia* (Acari: Erythraeidae) from Asia with description of a new species. *Systematic & Applied Acarology*, 20, 789–798.
<http://dx.doi.org/10.11158/saa.20.7.7>
- Noei, J., Saboori, A. & Hajizadeh, J. (2014) Two new species and one new record of larvae of the family Johnstonianidae (Acari: Prostigmata) from Iran with a key to species of the genus *Diplothrombium*. *Zootaxa*, 3785 (2), 241–257.
<http://dx.doi.org/10.11158/zootaxa.3785.2.7>
- Noei, J., Saboori, A. & Hajizadeh, J. (2015a) A new species and first record of larval *Paratrombium* (Acari: Trombidiidae) from Iran, with a key to world species. *Systematic & Applied Acarology*, 20, 313–321.
<http://dx.doi.org/10.11158/saa.20.3.8>
- Noei, J., Saboori, A. & Hajizadeh, J. (2015b) Three new species and two new records of the family Microtrombidiidae (Acari: Prostigmata) from Iran. *Systematic & Applied Acarology*, 20, 655–673.
<http://dx.doi.org/10.11158/saa.20.6.8>
- Noei, J., Saboori, A., Šundić, M., Hajizadeh, J. & Pešić, V. (2013) A new species and two new records of larval mites (Acari: Prostigmata; Erythraeidae, Smarididae) from northern Iran and Montenegro. *Systematic & Applied Acarology*, 18, 263–272.
<http://dx.doi.org/10.11158/saa.18.3.9>
- Paktinat-Saejj, S., Bagheri, M., Saboori, A. & Ahaniazad, M. (2015) Two new Bdellidae (Trombidiformes: Bdelloidea) from Iran and the status of *Neobiscirus* Gomelauri, 1963. *Zootaxa*, 4013 (4), 519–530.
<http://dx.doi.org/10.11158/zootaxa.4013.4.3>
- Paktinat-Saejj, S., Bagheri, M., Saboori, A., Seilsepour, N. & Ueckermann, E.A. (2015) A new snout mite, *Spinibdella tabarii* sp. nov. (Trombidiformes: Bdellidae) from Iran, with a summary of *Spinibdella* distributions worldwide. *Systematic & Applied Acarology*, 20, 693–706.
<http://dx.doi.org/10.11158/saa.20.6.11>
- Pepato, A.R. & Da Silveira, P.S.A. (2013) Two new species of *Halacarus* (Acari, Prostigmata) from Brazil. *Zootaxa*, 3670 (4), 591–600.
<http://dx.doi.org/10.11158/zootaxa.3670.4.10>
- Pepato, A.R. & Da Silveira, P.S.A. (2015) Two new species of *Rhombognathus* (Halacaridae, Trombidiformes) from a Mangrove in the northern littoral zone of São Paulo State (Brazil). *Zootaxa*, 3905 (4), 500–510.
<http://dx.doi.org/10.11158/zootaxa.3905.4.4>
- Pešić, V. (2013a) A remarkable new *Nilotonia* species (Acari, Hydrachnidia, Anisitsiellidae) from percolating water of a cave in Cat Ba island in Halong Bay, Vietnam. *Zootaxa*, 3710 (4), 372–380.
<http://dx.doi.org/10.11158/zootaxa.3710.4.6>
- Pesic, V. (2013b) Pontarachnid mites from marine interstitial, with a description of three new species from South Korea (Acari: Hydrachnidia: Pontarachnidae). *Zootaxa*, 3701 (1), 83–92.
<http://dx.doi.org/10.11158/zootaxa.3701.1.7>
- Pešić, V. (2015) A new species of the water mite genus *Hygrobates* Koch, 1837 (Acari: Hydrachnidia: Hygrobatidae) from the ancient Lake Ohrid. *Zootaxa*, 3926 (2), 287–295.
<http://dx.doi.org/10.11158/zootaxa.3926.2.9>
- Pešić, V., Chatterjee, T., Das, M.K. & Bordoloi, S. (2013) A new species of water mite (Acari, Hydrachnidia) from Assam, India, found in the gut contents of the fish *Botia dario* (Botiidae). *Zootaxa*, 3746 (3), 454–462.
<http://dx.doi.org/10.11158/zootaxa.3746.3.4>
- Pešić, V. & Gerecke, R. (2014) Water mites from caves of the Ha Giang province, northern Vietnam (Acari: Hydrachnidia). *Zootaxa*, 3774 (4), 367–380.
<http://dx.doi.org/10.11158/zootaxa.3774.4.5>
- Pešić, V., Piccoli, G., Araújo, M., Rezende, J.M. & Gonçalves, A.Z. (2015) A new species of *Xystonotus* Wolcott, 1900 (Acari, Hydrachnidia, Mideopsidae) from bromeliad phytotelmata in Brazilian Atlantic rainforest. *Zootaxa*, 3981 (1), 147–150
<http://dx.doi.org/10.11158/zootaxa.3981.1.10>
- Pešić, V. & Semenchenko, K. (2014) Water mites of the genus *Brachypoda* Lebert, 1879 (Acari: Hydrachnidia: Aturidae) from South Korea and the Russian Far East. *Zootaxa*, 3753 (4), 335–346.
<http://dx.doi.org/10.11158/zootaxa.3753.4.3>
- Pešić, V. & Smit, H. (2014a) Torrenticoloid water mites (Acari: Hydrachnidia: Torrenticolidae) from Ghana. *Zootaxa*, 3820 (1), 1–80.
<http://dx.doi.org/10.11158/zootaxa.3820.1.1>
- Pešić, V. & Smit, H. (2014b) Torrenticoloid water mites (Acari: Hydrachnidia: Torrenticolidae) from Malaysian Borneo. *Zootaxa*, 3840 (1), 1–72.
<http://dx.doi.org/10.11158/zootaxa.3840.1.1>
- Pešić, V. & Smit, H. (2015a) Two new species of the genus *Atractides* Koch, 1837 (Acari: Hydrachnidia: Hygrobatidae) with an updated checklist of the water mites of Thailand. *Systematic & Applied Acarology*, 20, 782–788.

- http://dx.doi.org/10.11158/saa.20.7.6
- Pešić, V. & Smit, H. (2015b) Water mites of the genus *Atractides* Koch, 1837 (Acari: Hydrachnidia: Hygrobatiidae) from Ghana. *Zootaxa*, 3911 (3), 343–356.
<http://dx.doi.org/10.11646/zootaxa.3911.3.2>
- Pešić, V., Smit, H. & Mary, N. (2015) Third contribution to the knowledge of water mites from the Comoros, with the description of two new species (Acari: Hydrachnidia). *Zootaxa*, 3964 (4), 445–459.
<http://dx.doi.org/10.11646/zootaxa.3964.4.4>
- Pešić, V.G. & Smit, H. (2015) Fourth contribution to the knowledge of water mites from the Comoros, with the description of two new species (Acari: Hydrachnidia). *Zootaxa*, 4052 (5), 589–594.
<http://dx.doi.org/10.11646/zootaxa.4052.5.8>
- Petanović, R.U., Amrine, J., Chetverikov, P.E. & Cvrković, T.K. (2015) *Eriocaenus* (Acari: Trombidiformes: Eriophyoidea), a new genus from *Equisetum* spp. (Equisetaceae): morphological and molecular delimitation of two morphologically similar species. *Zootaxa*, 4013 (1), 51–66.
<https://doi.org/10.11646/zootaxa.4013.1.3>
- Rajput, S., Han, X., Xue, X.-F. & Hong, X.-Y. (2014) A new genus and three new species of Diptilomiopidae from Zhejiang Province, China. *Systematic & Applied Acarology*, 19, 223–235.
<http://dx.doi.org/10.11158/saa.19.2.13>
- Rajput, S., Li, H., Xue, X.-F. & Hong, X.-Y. (2015) Descriptions of five new eriophyoid mite species of the Diptilomiopidae (Acari: Trombidiformes: Eriophyoidea) from Zhejiang Province, China. *Zootaxa*, 4020 (2), 335–356.
<http://dx.doi.org/10.11646/zootaxa.4020.2.6>
- Rajput, S., Zuo, Y., Wang, Z. & Hong, X.-Y. (2015) Five new species of the genus *Tetra* Keifer (Acari: Eriophyidae: Phyllocoptinae) from China. *Systematic & Applied Acarology*, 20, 203–219.
<http://dx.doi.org/10.11158/saa.20.2.7>
- Ramirez-Sánchez, M.M. & Rivas, G. (2013) New species of subgenus *Megaluracarus* (Acari: Hydrachnidiae: Arrenuridae: Arrenurus) from Mexico. *Zootaxa*, 3718 (4), 317–330.
<http://dx.doi.org/10.11646/zootaxa.3718.4.2>
- Ripka, G. (2014) Eriophyoid mites (Acari: Prostigmata: Eriophyoidea) from Hungary: a new species on *Agrimonia eupatoria* (Rosaceae) and new record on *Convolvulus arvensis* (Convolvulaceae). *Zootaxa*, 3900 (2), 263–270.
<http://dx.doi.org/10.11646/zootaxa.3900.2.6>
- Ripka, G., Laniecka, I. & Kazmierski, A. (2013) On the arboreal acarofauna of Hungary: some new and rare species of prostigmatic mites (Acari: Prostigmata: Tydeidae, Iolinidae and Stigmeidae). *Zootaxa*, 3702 (1), 1–50.
<http://dx.doi.org/10.11646/zootaxa.3702.1.1>
- Rocha, M.D.S., Skvarla, M.J. & Ferla, N.J. (2013) A new species of *Scutopalus* (Acari: Cunaxidae: Cunaxoidinae) from Rio Grande do Sul State, Brazil with a key to world species. *Zootaxa*, 3734 (1), 38–44.
<http://dx.doi.org/10.11646/zootaxa.3734.1.4>
- Rocha, M.S., Rodrigues, E.N.L. & Ferla, N.J. (2015) New species and records of cunaxid mites (Acari: Cunaxidae) from soil in Southern Brazil. *Zootaxa*, 3981 (1), 56–70.
<http://dx.doi.org/10.11646/zootaxa.3981.1.2>
- Saboori, A., Pešić, V. & Šundic, M. (2015) First record of *Podothrombium* (Acari: Podothrombiidae) from Serbia with description of a new species based on larvae. *Systematic & Applied Acarology*, 20, 121–128.
<http://dx.doi.org/10.11158/saa.20.1.11>
- Saboori, A., Wohltmann, A., Hakimitabar, M. & Shirvani, A. (2013) A new species of the genus *Achaemenothrombium* (Acari: Achaemenothrombiidae) from Iran. *Zootaxa*, 3694 (2), 143–152.
<http://dx.doi.org/10.11646/zootaxa.3694.2.3>
- Samadpour, M., Khanjani, M. & Asalifayaz, B. (2014) *Neognathus sinaei* sp.nov. (Acari: Caligonellidae), a new species from western Iran. *Systematic & Applied Acarology*, 19, 144–153.
<http://dx.doi.org/10.11158/saa.19.2.4>
- Sevsay, S. & Karakurt, I. (2013) A new species of the genus *Empitrombium* Southcott, 1994 (Acari: Microtrombidiidae) from Turkey. *Zootaxa*, 3709 (3), 255–266.
<http://dx.doi.org/10.11646/zootaxa.3709.3.4>
- Sikora, B., Kaszewska, K. & Skoracki, M. (2014) Two new quill mites of the family Syringophilidae (Acari: Prostigmata) parasitising the tapaculos (Passeriformes: Rhinocryptidae) in South America. *Zootaxa*, 3895 (3), 419–426.
<http://dx.doi.org/10.11646/zootaxa.3895.3.6>
- Silva-de la Fuente, M.C., Paredes-Leon, R., Casanueva, M.E., Escobar-Huerta, G. & Salas, L.M. (2015) A new genus and species of pterygosomatid mite (Acari: Pterygosomatidae) parasitizing *Callopistes maculatus* (Squamata: Teiidae) from Chile. *Zootaxa*, 3972 (1), 65–74.
<http://dx.doi.org/10.11646/zootaxa.3972.1.4>
- Silva, G., Cunha, U., Rocha, M.S., Panou, E.N. & Ferla, N.J. (2014) Tydeid and triophtydeid mites (Acari: Tydeoidea) associated with grapevine (Vitaceae: *Vitis* spp.) in Brazil, with the descriptions of species of *Prelorryia* (André, 1980) and *Tydeus* Koch, 1835. *Zootaxa*, 3814 (4), 495–511.

- http://dx.doi.org/10.11646/zootaxa.3814.4.3
- Skoracki, M., Hromada, M. & Unoeld, M. (2013) Three new quill mite species of the genus *Neoaulonastus* Skoracki (Acari: Syringophilidae) parasitizing passerines in Tanzania. *Zootaxa*, 3616 (4), 367–377.
http://dx.doi.org/10.11646/zootaxa.3616.4.5
- Skoracki, M., Kaszewska, K. & Kavetska, K. (2015) Two new species of the syringophilid quill mites (Acari: Prostigmata: Syringophilidae) parasitizing apodiform birds (Aves: Apodiformes). *Zootaxa*, 4052 (5), 583–588.
http://dx.doi.org/10.11646/zootaxa.4052.5.7
- Skoracki, M., Klimovičová, M., Muchai, M. & Hromada, M. (2014) New taxa of the family Syringophilidae (Acari: Prostigmata) from African barbets and woodpeckers (Piciformes: Lybiidae, Picidae). *Zootaxa*, 3768 (2), 178–188.
http://dx.doi.org/10.11646/zootaxa.3768.2.5
- Skoracki, M. & Mironov, S.V. (2013) New species and records of quill mites of the family Syringophilidae (Acari: Prostigmata) from the passerines (Aves: Passeriformes) from the Russian Far East. *Zootaxa*, 3641 (5), 554–564.
http://dx.doi.org/10.11646/zootaxa.3641.5.4
- Skoracki, M., Mironov, S.V. & Unoeld, M. (2013) The first records of quill mites of the family Syringophilidae (Acariformes: Prostigmata: Cheyletoidea) from trogoniform birds (Aves: Trogoniformes). *Zootaxa*, 3701 (2), 291–297.
http://dx.doi.org/10.11646/zootaxa.3701.2.9
- Skoracki, M. & Sikora, B. (2014) Two new quill mite species of the family Syringophilidae (Acari: Prostigmata) parasitising the house sparrow *Passer domesticus* (L.) (Aves: Passeriformes). *Zootaxa*, 3765 (2), 194–200.
http://dx.doi.org/10.11646/zootaxa.3765.2.7
- Smit, H. (2013) New records of water mites mainly from Vogelkop, New Guinea (Acari: Hydrachnidia), with the description of one new genus and thirteen new species. *Zootaxa*, 3716 (2), 207–235.
http://dx.doi.org/10.11646/zootaxa.3716.2.5
- Smit, H. (2014) Australian water mites of the genus *Hydrachna* Müller, with the description of five new species (Acari: Hydrachnidia: Hydrachnidae). *Zootaxa*, 3827 (2), 170–186.
http://dx.doi.org/10.11646/zootaxa.3827.2.3
- Smit, H. (2015a) A new notoaturnine genus from New Zealand (Acari: Hydrachnidia: Aturidae: Notoaturinae). *Zootaxa*, 4033 (3), 442–444.
http://dx.doi.org/10.11646/zootaxa.4033.3.9
- Smit, H. (2015b) The water mite family Hygrobatidae Koch in Australia. The genera *Aspidiobatella* Cook, *Australorivacarus* Viets, *Gondwanabates* Imamura and *Rhynchaustrobates* Cook (Acari: Hydrachnidia). *Zootaxa*, 4033 (4), 567–583.
http://dx.doi.org/10.11646/zootaxa.4033.4.7
- Smit, H. (2015c) The water mite genus *Pro corticacarus* K.O. Viets in Australasia (Acari: Hydrachnidia, Hygrobatidae). *Zootaxa*, 3956 (1), 56–78.
http://dx.doi.org/10.11646/zootaxa.3956.1.3
- Smit, H. & Pešić, V. (2014) The first Asian record of the water mite genus *Thoracophoracarus* K. Viets (Hydrachnidia: Arrenuridae). *Systematic & Applied Acarology*, 19, 431–434.
http://dx.doi.org/10.11158/saa.19.4.6
- Smit, H. & Pešić, V. (2014) Water mites from Mount Kinabalu and the Crocker Range, Borneo, Malaysia (Acari: Hydrachnidia), with the description of 34 new species. *Zootaxa*, 3876 (1), 1–71.
http://dx.doi.org/10.11646/zootaxa.3876.1.1
- Sousa, J.M., Lofego, A.C. & Gondim, M. (2014) Two new species of tarsonemid mites (Acari: Tarsonemidae) from northeastern Brazil. *Zootaxa*, 3889 (3), 429–441.
http://dx.doi.org/10.11646/zootaxa.3889.3.5
- Stathakis, T., Kapaxidi, E. & Papadoulis, G. (2014) A new record and a new species of the genus *Agistemus* Summers (Acari: Stigmaeidae) from Greece. *Zootaxa*, 3780 (1), 153–170.
http://dx.doi.org/10.11646/zootaxa.3780.1.6
- Stekolnikov, A.A. (2013) *Leptotrombidium* (Acari: Trombiculidae) of the world. *Zootaxa*, 3728 (1), 1–173.
http://dx.doi.org/10.11646/zootaxa.3728.1.1
- Stekolnikov, A.A. & González-Acuña, D. (2015) A review of Chilean chiggers (Acari: Trombiculidae), with the description of a new genus and ten new species. *Zootaxa*, 3964 (1), 1–43.
http://dx.doi.org/10.11646/zootaxa.3964.1.1
- Tan, M.-C., Yang, J., Lai, X.-S. & Wang, G.-Q. (2015) Four new species of the family Diptilomiopidae (Acari, Trombidiformes, Eriophyoidea) from Hainan Island, South China. *Systematic & Applied Acarology*, 20, 769–781.
http://dx.doi.org/10.11158/saa.20.7.5
- Tuzovskij, P.V. (2013a) New water mite species (Acari, Hydrachnidia) from Vietnam. *Zootaxa*, 3700 (4), 547–560.
http://dx.doi.org/10.11646/zootaxa.3700.4.3
- Tuzovskij, P.V. (2013b) New water mite species of the genus *Piona* Koch, 1842 (Acari, Hydrachnidia: Pionidae) from Northeastern Russia. *Zootaxa*, 3664 (3), 321–334.
http://dx.doi.org/10.11646/zootaxa.3664.3.3
- Tuzovskij, P.V. & Semenchenko, K.A. (2015) Water mites of the genus *Unionicola* Haldeman, 1842 (Acari, Hydrachnidia,

- Unionicolidae) in Russia. *Zootaxa*, 3919 (3), 401–456.
<http://dx.doi.org/10.11646/zootaxa.3919.3.1>
- Vidović, B. (2014) A new species and record of *Aceria* (Acari: Prostigmata: Eriophyoidea) on *Carlina* spp. (Asteraceae) from Serbia. *Zootaxa*, 3838 (4), 486–494.
<http://dx.doi.org/10.11646/zootaxa.3838.4.7>
- Wang, Q., El-Halawany, A., Xue, X.-F. & Hong, X.-Y. (2014) New species and records of eriophyoid mites from Saudi Arabia (Acari: Eriophyoidea). *Systematic & Applied Acarology*, 19, 409–430.
<http://dx.doi.org/10.11158/saa.19.4.5>
- Xu, Y., Fan, Q.-H. & Huang, J. (2014) Description of a new species of *Terminalichus* (Acari: Trombidiformes: Tenuipalpidae) from China. *Zootaxa*, 3753 (3), 233–250.
<http://dx.doi.org/10.11646/zootaxa.3753.3.3>
- Xu, Y., Fan, Q.-H. & Huang, J. (2015) Description and ontogeny of *Krugeria fuzhouensis* sp.nov. (Acari: Trombidiformes: Tenuipalpidae). *Systematic & Applied Acarology*, 20, 87–109.
<http://dx.doi.org/10.11158/saa.20.1.9>
- Xu, Y., Fan, Q.-H. & Zhang, Z.-Q. (2013) A new species of *Ultratenuipalpus* (Acari: Tenuipalpidae) from Cook Islands, with a key to the known species. *Zootaxa*, 3731 (2), 223–233.
<http://dx.doi.org/10.11646/zootaxa.3731.2.3>
- Xu, Y. & Zhang, Z.-Q. (2013) New Zealand Tenuipalpidae (Acari: Trombidiformes): A new species of *Acaricis* from Cyperaceae and its ontogenetic patterns in chaetotaxy. *Systematic & Applied Acarology*, 18, 357–388.
<http://dx.doi.org/10.11158/saa.18.4.6>
- Xu, Y. & Zhang, Z.-Q. (2014) *Prolixus* (Acari: Trombidiformes: Tenuipalpidae) newly recorded from New Zealand: A new species from Cyperaceae and its ontogenetic patterns in chaetotaxy. *Zootaxa*, 3900 (1), 1–20.
<http://dx.doi.org/10.11646/zootaxa.3900.1.1>
- Xue, X.-F., Cheng, L.-S. & Hong, X.-Y. (2013) Eriophyoid mites from Hainan Province, China VII: Descriptions of four new species (Acari: Diptilomiopidae). *Systematic & Applied Acarology*, 18, 177–189.
<http://dx.doi.org/10.11158/saa.18.2.10>
- Xue, X.-F., Guo, J.-F. & Hong, X.-Y. (2013) Eriophyoid mites from Northeast China (Acari: Eriophyoidea). *Zootaxa*, 3689 (1), 1–123.
<http://dx.doi.org/10.11646/zootaxa.3689.1.1>
- Xue, X.-F., Han, X. & Zhang, Z.-Q. (2015) Correct identification and biosecurity decision-making: Two species instead of one in *Aceria genistae* complex (Acari: Eriophyidae) in New Zealand. *Systematic & Applied Acarology*, 20, 71–86.
<http://dx.doi.org/10.11158/saa.20.1.8>
- Xue, X.-F., Sadeghi, H., Hong, X.-Y. & Sinaie, S. (2013) New species and records of eriophyid mites from Iran (Acari: Eriophyidae). *Systematic & Applied Acarology*, 18, 41–52.
<http://dx.doi.org/10.11158/saa.18.1.5>
- Yi, T.-C. & Zhang, Z.-Q. (2013) A new species of the genus *Sonotetranychus* (Acari: Tetranychidae) from New Zealand. *Zootaxa*, 3721 (4), 334–350.
<http://dx.doi.org/10.11646/zootaxa.3721.4.2>
- Zhang, Z.Q. (2014) Continued growth of *Systematic and Applied Acarology*, and hot spots and shelf life of new species in 2013. *Systematic & Applied Acarology*, 19, 110–112.
<http://dx.doi.org/10.11158/saa.19.1.9>
- Zhang, Z.-Q. (2015) New development and scope for *Systematic & Applied Acarology*. *Systematic & Applied Acarology*, 20, 153–154.
<http://dx.doi.org/10.11158/saa.20.1.14>
- Zhang, Z.Q. (2016) The rise of Systematic & Applied Acarology during its second decade. *Systematic & Applied Acarology*, 21, 146–146.
<http://dx.doi.org/10.11158/saa.21.1.10>
- Zhang, Z.Q., Fan, Q.-H., Pesic, V., Smit, H., Bochkov, A.V., Khaustov, A.A., Baker, A., Wohltmann, A., Wen, T.-H., Amrine, J.W., Beron, P., Lin, J.-Z., Gabrys, G. & Husband, R. (2011) Order Trombidiformes Reuter, 1909. In: Zhang, Z.Q. (Ed.) *Animal biodiversity: an outline of higher-level classification and survey of taxonomic richness*. *Zootaxa*, 3148, 129–138.
- Zuo, Y., Rajput, S., Xue, X.-F. & Hong, X.-Y. (2015) Three new species of Nothopodinae (Acari: Eriophyidae) from Guangdong Province, China. *Systematic & Applied Acarology*, 20, 286–296.
<http://dx.doi.org/10.11158/saa.20.3.6>