



Zootaxa 5561 (1): 001–081

<https://www.mapress.com/zt/>

Copyright © 2024 Magnolia Press

Monograph

ISSN 1175-5326 (print edition)

ZOOTAXA

ISSN 1175-5334 (online edition)

<https://doi.org/10.11646/zootaxa.5561.1.1>

<http://zoobank.org/urn:lsid:zoobank.org:pub:27863A22-14CF-4AC3-9BD5-25CE994BADA3>

ZOOTAXA

5561

Lech Borowiec's collection of tortoise beetles (Coleoptera: Chrysomelidae: Cassidinae) in the Museum of Natural History, University of Wrocław

JOLANTA ŚWIĘTOJAŃSKA

Department of Biodiversity and Evolutionary Taxonomy, University of Wrocław, Przybyszewskiego 65, 51-148 Wrocław, Poland,

[✉ jolanta.swietojanska@uwr.edu.pl](mailto:jolanta.swietojanska@uwr.edu.pl); [ORCID](https://orcid.org/0000-0002-6701-8809) <https://orcid.org/0000-0002-6701-8809>



Magnolia Press
Auckland, New Zealand

Accepted by W. Flowers: 16 Nov. 2024; published: 27 Dec. 2024

Licensed under Creative Commons Attribution-N.C. 4.0 International <https://creativecommons.org/licenses/by-nc/4.0/>

JOLANTA ŚWIĘTOJAŃSKA

Lech Borowiec's collection of tortoise beetles (Coleoptera: Chrysomelidae: Cassidinae) in the Museum of Natural History, University of Wrocław
(*Zootaxa* 5561)

81 pp.; 30 cm.

27 Dec. 2024

ISBN 978-1-77973-241-5 (paperback)

ISBN 978-1-77973-242-2 (Online edition)

FIRST PUBLISHED IN 2024 BY

Magnolia Press

P.O. Box 41-383

Auckland 1041

New Zealand

e-mail: magnolia@mapress.com

<https://www.mapress.com/zt>

© 2024 Magnolia Press

All rights reserved.

No part of this publication may be reproduced, stored, transmitted or disseminated, in any form, or by any means, without prior written permission from the publisher, to whom all requests to reproduce copyright material should be directed in writing.

This authorization does not extend to any other kind of copying, by any means, in any form, and for any purpose other than private research use.

ISSN 1175-5326 (Print edition)

ISSN 1175-5334 (Online edition)

Table of Contents

Abstract	4
Introduction	4
List of collection	4
Tribe Aspidimorphini	4
Tribe Cassidini	13
Tribe Delocraniini	41
Tribe Dorynotini	41
Tribe Eugenysini	42
Tribe Goniocheniini	43
Tribe Hemisphaerotini	44
Tribe Ischyrosonychini	44
Tribe Mesomphaliini	46
Tribe Notosacanthini	55
Tribe Omocerini	60
Tribe Spilophorini	62
Appendix	63
References	72
A list of papers with descriptions of new taxa whose types are preserved in the Lech Borowiec collection.	79
Reference cited in Introduction.	81

Abstract

Lech Borowiec's collection of tortoise beetles, recently stored in the Museum of Natural History, University of Wrocław is reviewed in detail. The collection contains 21 292 specimens of 2 491 species, including 229 primary types and 979 secondary types (paratypes, syntypes, paralectotypes). Additionally, a complete list of 202 papers on Cassidinae published by Prof. Lech Borowiec, a list of publications written by other authors with descriptions of new species whose descriptive types are deposited in the collection of Lech Borowiec, and a checklist of all taxa described by Prof. Lech Borowiec are given.

Key words: Coleoptera, Chrysomelidae, Cassidinae, museum collection

Introduction

In 2023, Prof. Lech Borowiec reached the age of 71. In 1984–2023 he conducted research on the biodiversity and taxonomy of the tortoise beetles (Coleoptera: Chrysomelidae: Cassidinae, mostly sensu stricto, except most of the former Hispinae). During this time, he published 202 papers, including the monumental 6-part monograph of the African Cassidinae (Borowiec 1994 c, 1997 g, 2002; Borowiec & Świętojańska 2017 b, 2018 c, 2022), two volumes of the monograph of Madagascan Cassidinae (Borowiec & Świętojańska 2011 h, 2013 h), the catalog of world Cassidinae (Borowiec 1999 k), and dozens of faunistic and taxonomic works, in which he proposed 12 new genera or subgenera and 417 species-level taxa (see appendix). On retirement, Prof. Lech Borowiec decided to focus his scientific activity on studying taxonomy and zoogeography of Balkan and Mediterranean ants and donated his Cassidinae collection to the Museum of Natural History, University of Wrocław (MNHW, apart from representatives of the former Imatidiini tribe, which were donated to the collection of Dr. Lukas Sekerka, Prague, Czech Republic).

The collection is now available for study under the standard procedures used at MNHW. To properly present the collection's richness, a detailed checklist of species therein deposited is presented below. The checklist also includes annotations on the number of specimens of each recorded species and their type status. In addition, collected specimens are listed per country. The checklist is organized alphabetically—tribes, genera, and species. For taxa whose descriptive types are present in the collection, the annotation includes the author and year of the description and the page number of the original publication.

References consist of a complete list of papers on Cassidinae authored by Lech Borowiec. Papers with descriptions of new taxa whose types are deposited in MNHW are marked in bold. Additionally, a list of papers with descriptions of new taxa whose types are preserved in Lech Borowiec's collection is also included.

The nomenclature of genera and species is consistent with that used in the collection at the time of its transfer to the Museum of Natural History, University of Wrocław (MNHW), in 2022. As for the Palearctic and Oriental taxa, it does not take into account the changes proposed in the Catalogue of Palearctic Coleoptera (Sekerka & Świętojańska 2024), to enable the curator of the collection to quickly find a specific species in the collection. MNHW includes data from the collection in the central museum database, which will use both the valid names, according to the current revision work, and the names used on labels in the L. Borowiec collection.

List of collection

(abbreviations: HT—holotype, PT—paratype, ST—syntype, PLT—paralectotype)

Tribe Aspidimorphini

Genus *Aspidimorpha* Hope, 1840

Subgenus *Aspidimorpha* s. str.

adhaerens (Weber, 1801): Indonesia (new Guinea): 7, Papua new Guinea: 36, Solomon Islands: 15; Total **58**

adjecta Spaeth, 1940: Cameroon: 1, Dem. Rep. of Congo: 1; Total **2**

adumbrata Spaeth, 1917: Kenya: 11; Total **11**

amabilis Boheman, 1854: Indonesia: 20; Total **20**
andrei Borowiec, 1997 g: 206: Dem. Rep. of Congo: 3 (**1 PT**); Total **3**
angoramensis Borowiec, 1992 c: 131: Papua New Guinea: 6 (**1 PT**); Total **6**
assimilis Boheman, 1854: Indonesia: 45, Malaysia: 5, Thailand: 1; Total **51**
astraea Spaeth, 1917: Angola: 2, Botswana: 1; Total **3**
atrodorsata Borowiec, 1997 g: 212: Dem. Rep. of Congo: 1 (**1 HT**); Total **1**
aurata (Montrouzier, 1855): Indonesia (New Guinea): 7, Papua New Guinea: 40, Solomon Islands: 33; Total **80**
australasiae (Boisduval, 1835): Indonesia: 1, Papua New Guinea: 57; Total **58**
bataviana Świętojańska, 2001 c: 45: Indonesia: 49 (**20 PT**); Total **49**
bertiae Borowiec, 1997 g: 215: Madagascar: 5 (**5 PT**); Total **5**
biguttata (Fabricius, 1775): Ethiopia: 1, Kenya: 6, Malawi: 1, Mozambique: 4, Tanzania: 29, no country: 1; Total **42**
bilobata Boheman, 1856: Philippines: 12; Total **12**
bimaculata (Fabricius, 1772): Cameroon: 1, Central African Rep.: 41, Dem. Rep. of Congo: 27, Ghana: 6, Ivory Coast: 1, Sierra Leone: 1, Tanzania: 1, Togo: 3, no country: 1; Total **82**
biradiata Boheman, 1854: Philippines: 5; Total **5**
biremis Boheman, 1854: Indonesia: 9; Total **9**
birmanica Spaeth, 1914: Myanmar: 2, Nepal: 2; Total **4**
castaneipennis Spaeth, 1912: Indonesia: 7, Malaysia: 1, Philippines: 8; Total **16**
chandrika Maulik, 1918: India: 5, Laos: 1, Thailand: 7, Vietnam: 2; Total **15**
collarti Borowiec, 1997 g: 235: Dem. Rep. of Congo: 2 (**1 HT, 1 PT**), Gabon: 1 (**1 PT**); Total **3**
convolvuli (Boheman, 1862): New Caledonia: 2; Total **2**
corrugata Borowiec, 1997 g: 237: Madagascar: 2 (**1 HT, 1 PT**); Total **2**
curtidens Hincks, 1962: Madagascar: 3; Total **3**
delitescens Weise, 1903: Cameroon: 1, Malawi: 4, Zambia: 3; Total **8**
densepicta Hincks, 1962: Madagascar: 3; Total **3**
deusta (Fabricius, 1775): Australia: 5, Indonesia: 33; Total **38**
diaphana (Sahlberg, 1823): Cameroon: 19, Central African Rep.: 4, Dem. Rep. of Congo: 27, Equatorial Guinea: 2, Gabon: 1, Nigeria: 1, Sierra Leone: 1, Tanzania: 1, Togo: 5; Total **61**
difformis (Motschulsky, 1860): China: 25, Japan: 4, North Korea: 1, Russia: 2; Total **32**
dilecta Boheman, 1854: RSA: 3, Zambia: 5; Total **8**
dissentanea Boheman, 1862: Angola: 7, Cameroon: 35, Central African Rep.: 11, Dem. Rep. of Congo: 62 (**1 PLT** *Aspidomorpha burgeoni* Spaeth, 1932: 9), Equatorial Guinea: 3, Gabon: 1, Ghana: 2, Guinea: 3, Ivory Coast: 1, Liberia: 1, Malawi: 1, Rwanda: 1, Sierra Leone: 2, South Sudan: 1, Tanzania: 3, Togo: 6, Uganda: 11; Total **151**
dorsata (Fabricius, 1787): Ceylon: 1, India: 5, Indonesia: 28, Laos: 6, Malaysia: 2, Thailand: 5, Vietnam: 1; Total **48**
dulcicula Boheman, 1862: Indonesia: 5; Total **5**
elevata (Fabricius, 1801): Indonesia: 37, Malaysia: 11, Vietnam: 1; Total **49**
equatoriensis Borowiec, 1986 a: 799: Dem. Rep. of Congo: 3, Ethiopia: 8, Kenya: 6 (**5 PT**), Somalia: 1, South Sudan: 2 (**2 PT**), Uganda: 4 (**1 PT**); Total **24**
ertli Spaeth, 1906: Tanzania: 3, Zambia: 7; Total **10**
expansa Spaeth, 1917: Tanzania: 4, Uganda: 1; Total **5**
extumida Spaeth, 1915: Madagascar: 2, Vietnam (mislabelled!): 1; Total **3**
fabricii Sekerka, 2008: Burkina Faso: 2, Cameroon: 1, Central African Rep.: 6, Dem. Rep. of Congo: 1, Gambia: 1, Mali: 1, Niger: 1, Senegal: 4, Togo: 1; Total **18**
fampanamboensis Borowiec, 1997 g: 289: Madagascar: 2 (**1 HT**); Total **2**
fausta Spaeth, 1932: Cameroon: 3, Dem. Rep. of Congo: 1; Total **4**
fenestrata (Olivier, 1808): Ivory Coast: 3, Nigeria: 2; Total **5**
filiola Borowiec, 1997 g: 299: Zimbabwe: 2 (**1 PT**); Total **2**
firma Weise, 1912: Tanzania: 1; Total **1**
fraterna Baly, 1863: Cambodia: 2, China: 2, Thailand: 1, Vietnam: 4; Total **9**

furcata (Thunberg, 1789): Ceylon: 1, China: 1, India: 24, Indonesia: 1, Laos: 4, Malaysia: 1, Nepal: 1, Philippines: 8, Thailand: 8, Vietnam: 29; Total **78**

fusca Shaw, 1961: Dem. Rep. of Congo: 5, Senegal (mislabelled!): 1, Zambia: 1; Total **7**

fusconotata Boheman, 1854: Philippines: 10; Total **10**

fuscopunctata Boheman, 1854: Gabon (mislabelled!): 1, India: 4, Indonesia: 11, Laos: 1, Myanmar: 1, Philippines: 1, Thailand: 5, Vietnam: 6; Total **30**

gruevi Borowiec, 1985 d: 451: Ethiopia: 3, Saudi Arabia: 2, Yemen: 6 (**1 PT**); Total **11**

hexaspilota Baly, 1863: Thailand: 6; Total **6**

honesta Spaeth, 1902: Dem. Rep. of Congo: 45; Total **45**

hyalina Spaeth, 1934: Dem. Rep. of Congo: 1, Zambia: 1; Total **2**

icterica Boheman, 1854: Ethiopia: 1, Kenya: 2, Malawi: 1, RSA: 13, Tanzania: 29; Total **46**

incerta Borowiec, 1997 g: 332: Cameroon: 3 (**3 PT**), Dem. Rep. of Congo: 2 (**2 PT**), Equatorial Guinea: 4 (**4 PT**); Total **9**

indica Boheman, 1854: India: 17, Indonesia: 8, Laos: 5, Malaysia: 23, Nepal: 3, Thailand: 2; Total **58**

indistincta Boheman, 1854: Burkina Faso: 5, Cameroon: 1, Central African Rep.: 3, Dem. Rep. of Congo: 3, Gambia: 4, Ivory Coast: 1, Nigeria: 1, Senegal: 4; Total **22**

infusca Spaeth, 1909: Dem. Rep. of Congo: 4; Kenya: 1, Tanzania: 6, Uganda: 4; Total **15**

ingens Duvivier, 1891: Cameroon: 1, Dem. Rep. of Congo: 6, Tanzania: 1; Total **8**

inquinata Boheman, 1854: Indonesia: 8; Total **8**

intermedia Świętojańska, 2001 c: 111: Philippines: 3 (**1 HT, 2 PT**); Total **3**

interrupta (Fabricius, 1775): Australia: 8; Total **8**

intricata Weise, 1903: Dem. Rep. of Congo: 1, Malawi: 1, Zambia: 4; Total **6**

inuncta Boheman, 1854: India: 15 (**1 ST** *Aspidomorpha spaethi* Maulik, 1918: 324); Total **15**

isparetta Boheman, 1854: Benin: 2, Cameroon: 35, Dem. Rep. of Congo: 78, Equatorial Guinea: 13, Gabon: 7, Ghana: 42, Guinea: 3, Ivory Coast: 13, Nigeria: 7, Rwanda: 2, Senegal (mislabelled!): 2, Sierra Leone: 2, Togo: 5, Uganda: 19, Upper Volta: 5, Vietnam (mislabelled!): 3, no country: 1; Total **239**

karamojana Borowiec & Sekerka, 2010 a: 271: Ethiopia: 2, Uganda: 7 (**6 PT**); Total **9**

katangana Spaeth, 1932: Dem. Rep. of Congo: 4, Zambia: 7, Zimbabwe: 1; Total **12**

kilimana Weise, 1903: Tanzania: 1, Zimbabwe: 1; Total **1**

laevigata Spaeth, 1905: Cameroon: 7, Equatorial Guinea: 2; Total **9**

levissima Borowiec, 1997 g: 374: Cameroon: 2 (**1 HT, 1 PT**); Total **2**

limbata (Goldfuss, 1805): Indonesia: 2; Total **2**

limbipennis Boheman, 1854: Indonesia: 2; Total **2**

lobata Boheman, 1854: Bangladesh: 1, Ceylon: 1, India: 13; Total **15**

luzonica Świętojańska, 2001 c: 125: Indonesia: 2 (**1 HT, 1 PT**); Total **2**

maculatissima (Boheman, 1856): Australia: 11; Total **11**

madagascariensis Boheman, 1854: Botswana: 2, Central African Rep.: 21, Dem. Rep. of Congo: 11, Ethiopia: 1, Kenya: 6, Madagascar: 30, Malawi: 4, RSA: 3, Tanzania: 27, Uganda: 2, Zambia: 1, Zimbabwe: 2; Total **110**

maffinbayensis Borowiec, 1992 c: 147: Indonesia (New Guinea): 4 (**1 PT**); Total **4**

malaccana Spaeth, 1937: Malaysia: 5, Thailand: 5; Total **10**

miliaris (Fabricius, 1775): China: 4, India: 13, Indonesia: 24, Laos: 19, Malaysia: 15, Nepal: 10, Papua New Guinea: 13, Philippines: 33, Thailand: 13, Vietnam: 10; Total **154**

mirabilis Borowiec, 1997 g: 388: Burundi: 1 (**1 PT**), Rwanda: 1 (**1 PT**), Uganda: 1 (**1 PT**); Total **3**

momboensis Weise, 1899: Kenya: 3, Tanzania: 1; Total **4**

montanella Borowiec, 1997 g: 395: Dem. Rep. of Congo: 18 (**18 PT**); Total **18**

mrogorensis Weise, 1899: Mozambique: 2, Tanzania: 4; Total **6**

muehlei Borowiec, 1997 g: 404: Dem. Rep. of Congo: 2 (**1 HT, 1 PT**); Total **2**

musta Spaeth, 1914: India: 1, Thailand: 3; Total **4**

mutata Boheman, 1854: Cameroon: 14, Dem. Rep. of Congo: 25, Gabon: 4, Guinea: 2, Rep. of Congo: 3, Rwanda: 1, Uganda: 2, no country: 1; Total **52**

mutilata Boheman, 1854: Indonesia: 29, Malaysia: 1; Total **30**

natalensis Boheman, 1854: Malawi: 2, RSA: 8, Tanzania: 2; Total **12**

nigropunctata (Klug, 1835): Angola: 1, Benin: 1, Burundi: 1, Cameroon: 3, Dem. Rep. of Congo: 21, Ghana: 3, Ethiopia: 3, Ivory Coast: 1, Kenya: 2, Malawi: 2, Mozambique: 1, RSA: 1, Somalia: 2, Tanzania: 16, Togo: 1, Zambia: 1, Zimbabwe: 3, no country: 1; Total **64**
novaguineensis (Boisduval, 1835): India (mislabeled): 1, Indonesia: 1, Myanmar (mislabeled): 1, Papua New Guinea: 32; Total **35**
oblectans Spaeth, 1924: Tanzania: 2; Total **2**
obovata (Klug, 1835): Cameroon: 10, Dem. Rep. of Congo: 12, Ghana: 13, Kenya: 2, Togo: 3, Uganda: 1; Total **41**
obtusangula Spaeth, 1906: Tanzania: 7; Total **7**
obuduensis Borowiec, 1997 g: 441: Cameroon: 1, Dem. Rep. of Congo: 1, Nigeria: 1 (**1 HT**); Total **3**
officiosa Boheman, 1862: Cameroon: 16, Central African Rep.: 3, Dem. Rep. of Congo: 29; Total **48**
orbicularis Boheman, 1854: Philippines: 6; Total **6**
orbifera Spaeth, 1934: Dem. Rep. of Congo: 3; Total **3**
orientalis Boheman, 1856: India: 4; Total **4**
pacalis Spaeth, 1926: Indonesia: 12; Total **12**
palleago Boheman, 1854: RSA: 4, Tanzania: 1; Total **5**
pallescens Spaeth, 1917: Dem. Rep. of Congo: 5, Zambia: 4; Total **9**
ponderosa Boheman, 1862: Indonesia: 2; Total **2**
pontifex Boheman, 1854: India (mislabeled!): 1, Madagascar: 5; Total **6**
potens Weise, 1903: Dem. Rep. of Congo: 2, RSA: 1, Tanzania: 2, Zambia: 5; Total **10**
procax Spaeth, 1932: 15: Cameroon: 3, Dem. Rep. of Congo: 5 (**2 ST**), Gabon: 1, Nigeria: 1; Total **10**
proszynskii Borowiec, 1985 b: 227: Ghana: 2 (**1 HT**), Togo: 1; Total **3**
punctum (Fabricius, 1801): Australia: 2, Indonesia (New Guinea): 15, Papua New Guinea: 45; Total **62**
quadrilobata Boheman, 1856: Philippines: 7; Total **7**
quadriradiata Boheman, 1854: Australia: 4; Total **4**
quadriramosa Boheman, 1854: Cameroon: 31, Central African Rep.: 8, Dem. Rep. of Congo: 21, Gabon: 8, Togo: 2; Total **70**
quadriremis (Gyllenhal, 1808): RSA: 40; Total **40**
quinquefasciata (Fabricius, 1801): Cameroon: 5, Central African Rep.: 2, Dem. Rep. of Congo: 1, Gambia: 9, Ghana: 1, Guinea Bissau: 1, Liberia: 4, Madagascar: 1, Neu Caledonia (introduced): 2, Nigeria: 9, Reunion: 2, São Tomé and Príncipe: 4, Senegal: 5, Togo: 4; Total **50**
reflexa Spaeth, 1917: Botswana: 2, Namibia: 5, RSA: 5; Total **12**
rubroornata Borowiec, 1997 g: 490 Madagascar: 6 (**1 HT, 2 PT**); Total **6**
sanctaecrucis (Fabricius, 1792): Cambodja: 1, China: 8, India: 39, Indonesia: 4, Laos: 1, Malaysia: 1, Nepal: 1, Vietnam: 21, Thailand: 7; Total **83**
sankuruensis Borowiec, 1997 g: 492: Dem. Rep. of Congo: 2 (**1 HT, 1 PT**); Total **2**
sarsinorum Spaeth, 1932: Indonesia: 1; Total **1**
sarawacensis Spaeth, 1904: Malaysia (Borneo): 4; Total **4**
sassii Borowiec, 1997 g: 495: Kenya: 5 (**4 PT**); Total: **5**
semiramosa Wagnere, 1880: Cameroon: 1, Kenya: 1, Togo: 1; Total **3**
sessarum Spaeth, 1911: Central African Rep.: 1, Dem. Rep. of Congo: 6, Uganda: 4; Total **11**
setosa Borowiec, 1997 g: 506: Dem. Rep. of Congo: 4 (**1 PT**), Equatorial Guinea: 1 (**1 PT**), Republic of Congo: 2 (**1 PT**); Total **7**
silfverbergi Borowiec, 1985: Guinea: 2, no country: 1; Total **3**
sjoestedti Spaeth, 1903: Cameroon: 5; Total **5**
snizeki Świętojańska, 2001 c: 173: China: 2 (**1 HT**); Total **2**
splendidula Spaeth, 1961: Dem. Rep. of Congo: 2; Total **2**
sternalis Weise, 1896: Cameroon: 4, Dem. Rep. of Congo: 25, Kenya: 2, Uganda: 2; Total **33**
stevensi Baly, 1863: Vietnam: 1 (**1 PT** *Aspidomorpha vietnamica* Medvedev & Eroshkina, 1982: 113); Total **1**
strigosa (Gorham, 1892): Dem. Rep. of Congo: 1, Tanzania: 7, Zambia: 27; Total **35**
suavis Spaeth, 1912: Malaysia (Borneo): 3; Total **3**
subcruciata Boheman, 1854: Philippines: 1; Total **1**

submutata Weise, 1899: Burkina Faso: 2, Cameroon: 2, Central African Rep.: 3, Equatorial Guinea: 2, Ethiopia: 1, Ivory Coast: 1, Malawi: 1, RSA: 2, Tanzania: 6, Togo: 2, Uganda: 1, Zambia: 2, Zimbabwe: 1, no country: 1; Total **27**

sulawesiensis Świętojańska, 2001 c: 183: Indonesia: 1 (**1 HT**); Total **1**

sulfuripennis Spaeth, 1932:16: Cameroon: 1 (**1 ST**), Dem. Rep. of Congo: 2; Total **3**

sumbawaensis Świętojańska & Borowiec, 2013 e: 218: Indonesia: 2 (**2 PT**); Total **2**

tamdaoensis Świętojańska, 2001 c: 185: Vietnam: 2 (**1 HT, 1 PT**); Total **2**

tanganikana Spaeth, 1916 c: Dem. Rep. of Congo: 11, Kenya: 3, Rwanda: 6, Uganda: 10; Total: **30**

tibetana Świętojańska & Borowiec, 2006 g: 542: China: 1 (**1 PT**); Total **1**

timorensis Świętojańska, 2001 c: 188: Indonesia: 2 (**1 HT, 1 PT**); Total **2**

togata Thomson, 1858: Cameroon: 58, Central African Rep.: 21, Dem. Rep. of Congo: 30, Gabon: 3, Kenya: 1, Uganda: 9; Total **122**

togoensis Weise, 1904: Cameroon: 1, Dem. Rep. of Congo: 1, Guinea: 1, Ivory Coast: 1, Sierra Leone: 1, Togo: 2; Total **7**

tortuosa Boheman, 1862: Cameroon: 50; Total **50**

toxopei Spaeth, 1926: Indonesia: 3; Total **3**

transparipennis (Motschulsky, 1860): China: 7, Japan: 2, North Korea: 2, Russia: 2; Total **13**

tuberosa Borowiec, 1997 g: 567: Dem. Rep. of Congo: 2 (**1 HT, 1 PT**); Total **2**

uelensis Spaeth, 1932: Cameroon: 4, Dem. Rep. of Congo: 6; Total **10**

uluguruensis Borowiec, 1997 g: 573: Tanzania: 14 (**6 PT**); Total **14**

undulatipennis Spaeth, 1911: Madagascar: 5; Total **5**

vernicata Fairmaire, 1901: Madagascar: 8; Total **8**

wagneri Borowiec & Sekerka, 2014 a: 198: Uganda: 1 (**1 HT**); Total **1**

wahlbergi Boheman, 1854: RSA: 1, Tanzania: 2; Total: **3**

westwoodii Boheman, 1854: Australia: 5; Total **5**

Aspidimorpha s. str. sp. 1: Gabon: 1; Total **1**

Subgenus *Afroaspidimorpha* Borowiec, 1997

areata (Klug, 1835): Dem. Rep. of Congo: 7, Ethiopia: 1, Kenya: 4, Malawi: 1, Namibia: 4, RSA: 5, Somalia: 2, South Sudan: 1, Tanzania: 3, Uganda: 4, Urundi, 1, Zambia: 2, Zimbabwe: 1; Total **36**

fallaciosa (Fairmaire, 1904): Madagascar: 5; Total **5**

loennebergi Spaeth, 1915: Kenya: 6; Total **6**

longifrons Shaw, 1961: Dem. Rep. of Congo: 6; Total **6**

multiguttata Spaeth, 1935: 173: Angola: 3 (**3 ST**); Total **3**

nigritarsis Spaeth, 1932: Malawi: 3; Total **3**

nigromaculata (Herbst, 1799): Angola: 1, Burkina Faso: 10, Cameroon: 9, Chad: 26, Dem. Rep. of Congo: 16, Ethiopia: 5, Gambia: 8, Ghana: 9, Guinea: 1, Kenya: 6, Namibia: 2, Nigeria: 2, Senegal: 4, Sierra Leone: 1, Sudan: 2, Uganda: 2, Zimbabwe: 4; Total **108**

polyspila Spaeth, 1911: 264: Madagascar: 17 (**1 ST**); Total **17**

pseudoareata Borowiec, 1997 g: 49: Kenya: 1 (**1 PT**), Uganda: 2 (**2 PT**); Total **3**

rainoni Borowiec & Coache, 2018 b: 1: Benin: 7 (**7 PT**); Total **7**

severini Spaeth, 1902: Dem. Rep. of Congo: 1, Malawi: 1, Namibia: 4, RSA: 2, Tanzania: 2, Zambia: 2; Total **12**

Subgenus *Aspidocassis* Borowiec, 1997

apicalis (Klug, 1833): Madagascar: 15, Seychelles: 3; Total **18**

confinis (Klug, 1835): Angola: 1, Burkina Faso: 3, Cameroon: 4, Dem. Rep. of Congo: 20, RSA: 4, Ghana: 6, Guinea: 1, Kenya: 3, Rwanda: 1, Sierra Leone: 1, South Sudan: 2, Uganda: 9, Zambia: 1; Total **56**

hiekei Borowiec, 1997 g: 74: Ethiopia: 1, Senegal: 3 (**1 PT**), Togo: 2 (**2 PT**); Total **6**

tanolaensis Borowiec, 1997 g: 76: Madagascar: 3 (**1 HT, 2 PT**); Total **3**

Subgenus *Dianaspis* Chen & Zia, 1984

denticollis Spaeth, 1932: Vietnam: 1; Total **1**

Subgenus *Megaspidomorpha* Hincks, 1952

angolensis Weise, 1896: Dem. Rep. of Congo: 3, Ethiopia: 4, Guinea (introduced?): 1, Kenya: 1, Mozambique: 4, RSA: 4, Tanzania: 3, Zambia: 14, Zimbabwe: 2; Total **36**

chlorotica (Olivier, 1808): Cameroon: 16, Central African Rep.: 2, Dem. Rep. of Congo: 15, Sierra Leone: 2, Uganda: 1, no country: 3; Total **39**

puncticosta Boheman, 1854: Kenya: 3, Mozambique: 3, RSA: 22, Tanzania: 12, Zambia: 3, no country: 1; Total **44**

Subgenus *Neoaspidomorpha* Borowiec, 1997

septemcostata Wagener, 1881: Australia: 7; Total **7**

Subgenus *Semiaspidomorpha* Borowiec, 1997

chlorina Boheman, 1854: Cameroon: 22, Central African Rep.: 1, Dem. Rep. of Congo: 29, Equatorial Guinea: 3, Uganda: 3; Total **58**

fatua Spaeth, 1917: Cameroon: 3, Dem. Rep. of Congo: 4, Sierra Leone: 1, Uganda: 2; Total **10**

irrorata Weise, 1898: Kenya: 4, Malawi: 3, Mozambique: 1, Tanzania: 22; Total **30**

kasaiensis Spaeth, 1932: Angola: 1, Cameroon: 2; Total **3**

kolbei Weise, 1899: Tanzania: 4; Total **4**

pseudochlorina Borowiec, 1986 a: 793: Benin: 1, Burkina Faso: 2, Cameroon: 1, Ghana: 6 (**2 PT**), Togo: 4 (**3 PT**); Total **14**

salazarensis Borowiec, 1997 g: 121: Angola: 1 (**1 HT**); Total **1**

viridula Weise, 1903: Dem. Rep. of Congo: 2, Ethiopia: 1, Guinea: 1, South Sudan: 1; Total **5**

Subgenus *Spaethia* Berg, 1899

alluaudi Spaeth, 1924: Dem. Rep. of Congo: 3; Total **3**

cepaecolor (Fairmaire, 1898): Madagascar: 3; Total **3**

citrina Spaeth, 1934: Zambia: 8, Zimbabwe: 1; Total **9**

exacta Spaeth, 1932: Tanzania: 1; Total **1**

ganglbaueri (Spaeth, 1898): RSA: 1, Tanzania: 1, Zambia: 23; Total **25**

gausapina Spaeth, 1924: RSA: 5; Total **5**

heroni Borowiec, 1997 g: 147: Kenya: 2 (**1 HT, 1 PT**); Total **2**

pellucida Weise, 1899: Dem. Rep. of Congo: 6, Tanzania: 1, Zambia: 1; Total **8**

scalena Spaeth, 1917: Zambia: 4; Total **4**

siticulosa Spaeth, 1917: 428: Dem. Rep. of Congo: 4 (**1 PT**), Ethiopia: 4, Tanzania: 1, Uganda: 4; Total **13**

socrus Spaeth, 1917: Dem. Rep. of Congo: 2, Zambia: 2; Total **4**

zambiana Borowiec, 2006 c: 249: Zambia: 1 (**1 HT**); Total **1**

Subgenus *Spaethiomorpha* Borowiec, 1997

haefligeri Spaeth, 1906: Malawi: 8, Tanzania: 1, Zimbabwe: 2; Total **11**

Subgenus *Weiseocassis* Spaeth, 1932

sculpturata Borowiec, 1997 g: 166: Tanzania: 1 (**1 PT**); Total **1**

striata Weise, 1896: Cameroon: 4, Dem. Rep. of Congo: 4, Kenya: 2, Tanzania: 17; Total **27**

Genus *Conchyloctenia* Spaeth, 1902

adpersa (Fabricius, 1801): Dem. Rep. of Congo: 10, Ethiopia: 1, Republic of Congo: 1, Sierra Leone: 3; Total **15**

aruwimiensis (Gorham, 1892): Dem. Rep. of Congo: 10, Uganda: 3; Total **13**

aspidiformis Borowiec, 1994 c: 53: RSA: 1 (**1 HT**); Total **1**

bipuncticollis (Boheman, 1854): RSA: 1, Suazi: 1, Zimbabwe: 1; Total **3**

bonnyana (Gorham, 1892): Cameroon: 8, Central African Rep.: 1, Dem. Rep. of Congo: 20, Nigeria: 3, Republic of Congo: 3; Total **35**

capensis Świętojańska & Borowiec, 2002 d: 212: RSA: 1 (**1 HT**); Total **1**

fibrata Spaeth, 1912: Dem. Rep. of Congo: 18, Ethiopia: 3, South Sudan: 1, Uganda: 5; Total **27**

hepatica (Boheman, 1854): Benin: 1, Cameroon: 13, Ethiopia: 1, Gabon: 5, Guinea: 1, Ivory Coast: 1, Togo: 1; Total **23**

hybrida (Boheman, 1854): Dem. Rep. of Congo: 14, Ethiopia: 2, Kenya: 1, Mozambique: 5, RSA: 6, Rwanda: 4, Tanzania: 8, Zambia: 7, no country: 3; Total **50**

illota (Boheman, 1854): RSA: 5; Total **5**

mouffleti (Boheman, 1854): Cameroon: 3, Central African Rep.: 1, Dem. Rep. of Congo: 25, Ethiopia: 1, Gabon: 1, Liberia: 3; Total **34**

multimaculata Spaeth, 1917: Angola: 1, Dem. Rep. of Congo: 4; Total **5**
nigrovittata (Boheman, 1854): India: 5; Total **5**
praecox (Boheman, 1854): Dem. Rep. of Congo: 3, Kenya: 1, Malawi: 7, Mozambique: 1, RSA: 3, Tanzania: 2, Uganda: 1, Zambia: 4, Zimbabwe: 4; Total **26**
punctata (Fabricius, 1787): Angola: 1, Botswana: 1, Dem. Rep. of Congo: 5, Ethiopia: 3, Gabon: 1, Kenya: 3, Malawi: 1, Mozambique: 1, Namibia: 11, RSA: 13, Tanzania: 6, Uganda: 1, Zambia: 8, Zimbabwe: 16, no country: 3; Total **74**
signatipennis (Boheman, 1854): Cameroon: 5, Central African Rep.: 2, Dem. Rep. of Congo: 26, Ethiopia: 2, Ghana: 12, Guinea: 1, Nigeria: 3, Sierra Leone: 1, Uganda: 4; Total **56**
tripuncticollis (Boheman, 1862): Botswana: 5, Kenya: 2, Mozambique: 1, Namibia: 16, RSA: 17, Tanzania: 4, Zambia: 1, Zimbabwe: 3; Total **49**

Genus *Hybosinota* Spaeth, 1909

nodulosa (Boheman, 1854): Botswana: 1, Namibia: 3, RSA: 1; Total **5**
turrigera (Boheman, 1862): Rwanda: 1, Tanzania: 1, Zambia: 1, Zimbabwe: 1; Total **4**

Genus *Laccoptera* Boheman, 1855

Subgenus *Laccoptera* s. str.

aurosa Fairmaire, 1891: Kenya: 3, Tanzania: 26; Total **29**
brancsiki Spaeth, 1919: 186: Zimbabwe: 3 (**1 ST**); Total **3**
excavata Boheman, 1855: RSA: 9, no country: 1; Total **10**
rugosicollis (Spaeth, 1902): Namibia: 8, RSA: 8, Tanzania: 1, Zambia: 9, Zimbabwe: 3, no country: 1; Total **30**
weisei (Spaeth, 1902): RSA: 4; Total **4**
zambesiaca Spaeth, 1919: 187: Malawi: 5 (**1 ST**), Mozambique: 2, RSA: 3, Tanzania: 1; Total **11**

Subgenus *Asphalesia* Weise, 1899

confragosa Weise, 1899: Madagascar: 2; Total **2**
pallicolor (Fairmaire, 1901): Madagascar: 3; Total **3**
perrieri Fairmaire, 1898: Madagascar: 5; Total **5**
regularis Fairmaire, 1898: Madagascar: 2; Total **2**
spectrum Boheman, 1855: Madagascar: 1; Total **1**
undulata (Spaeth, 1919): 191: Madagascar: 1 (**1 ST**); Total **1**

Subgenus *Laccopteroidea* Spaeth in Hincks, 1952

cheni Świętojańska, 2001 c: 209: China: 1 (**1 HT**); Total **1**
discreta Boheman, 1855: Indonesia: 1; Total **1**
fallax Weise, 1910: Indonesia: 18; Total **18**
fruhstorferi Spaeth, 1905: Laos: 2, Myanmar: 2, Thailand: 1; Total **5**
impressa (Blanchard, 1863): Australia: 3, Indonesia (New Guinea): 2, Papua New Guinea: 20; Total **25**
nepalensis Boheman, 1855: China: 8, India: 26, Laos: 19, Nepal: 2, Malaysia: 50, Taiwan: 2, Thailand: 24, Vietnam: 26; Total **157**
permodica (Boheman, 1862): Indonesia: 7; Total **7**
prominens Chen & Zia, 1964: China: 6; Total **6**
quatuordecimnotata Boheman, 1855: India: 8; Total **8**
sculpturata Boheman, 1855: 78: Indonesia: 1 (**1 PLT**); Total **1**
sedecimnotata Boheman, 1862: Australia: 1, Indonesia: 13; Total **14**
sutteri Hincks, 1953: Indonesia: 2; Total **2**
tredecimguttata Wagener, 1877: Philippines: 34; Total **34**
tredecimpunctata (Fabricius, 1801): India: 1, Indonesia: 34, Philippines: 13; Total **48**
yunnanica Spaeth, 1914 c: 226: China: 4 (**1 ST**); Total **4**

Subgenus *Orphnoda* Weise, 1899

cancellata Boheman, 1855: RSA: 10; Total **10**
intertexta Boheman, 1862: RSA: 4; Total **4**

Subgenus *Orphnodella* Spaeth, 1902

atrata Spaeth, 1905: Dem. Rep. of Congo: 8, Kenya: 1, Rwanda: 4; Total **13**

basalis Weise, 1899: Cameroon: 9, Gabon: 1; Total **10**

bicolor Spaeth, 1937: Cameroon: 5, Dem. Rep. of Congo: 2; Total **7**

burorum (Spaeth, 1902): RSA: 3; Total **3**

caduca Borowiec, 1994 c: 202: Rep. of Congo: 5 (**5 PT**); Total **5**

cicatricosa Boheman, 1855: Botswana: 2, Cameroon: 1, Erytrea: 3, Ethiopia: 3, Kenya: 1, Malawi: 1, Mali: 1, Mozambique: 1, Namibia: 5, RSA: 2, Senegal: 3, Somalia: 1, Zambia: 2, Zimbabwe: 3; Total **29**

corrugata (Sahlberg, 1823): Cameroon: 27, Central African Rep.: 3, Dem. Rep. of Congo: 49, Gabon: 6, Ghana: 2, Guinea: 2, Ivory Coast: 1, São Tomé and Príncipe: 3, Togo: 2; Total **95**

deremensis Weise, 1899: Kenya: 1, Malawi: 1, Tanzania: 5, Zimbabwe: 2; Total **9**

foveolata (Boheman, 1856): India: 3; Total **3**

kapiriana Spaeth, 1932: Dem. Rep. of Congo: 3; Total **3**

montivaga Spaeth, 1909: Cameroon: 2, Central African Rep.: 6, Dem. Rep. of Congo: 92, Ethiopia: 9, Ghana: 1, Kenya: 3, Rwanda: 3, Tanzania: 1, Uganda: 7; Total **124**

nigricornis Wagener, 1877: Gabon: 2; Total **2**

nunbergi Borowiec, 1994 c: 232: Cameroon: 7 (**6 PT**), Gabon: 2 (**2 PT**); Total **9**

obscurumaculata Borowiec, 2010 c: 287: Dem. Rep. of Congo: 5 (**4 PT**); Total **5**

rotundicollis Borowiec, 1997 c: 664: Burkina Faso: 1 (**1 HT**), Senegal: 6 (**1 PT**), Togo: 3 (**2 PT**); Total **10**

rubricollis Spaeth, 1932: Dem. Rep. of Congo: 4; Total **4**

rugicollis Thomson, 1858: Cameroon: 5, Dem. Rep. of Congo: 2, Ghana: 9; Total **16**

ruginosa Boheman, 1855: Mozambique: 1, RSA: 6, Tanzania: 2, Zimbabwe: 2; Total **11**

sassana Spaeth, 1912: Cameroon: 2, Central African Rep.: 3, Dem. Rep. of Congo: 6; Total **11**

triangula Spaeth, 1912: Cameroon: 3, Dem. Rep. of Congo: 6, Equatorial Guinea: 3, Gabon: 2; Total **14**

Subgenus *Orphnodina* Spaeth, 1932

distans (Spaeth, 1902): Zambia: 4, Zimbabwe: 5; Total **9**

rustica (Weise, 1899): Mozambique: 2, RSA: 6; Total **8**

Subgenus *Patrisma* Fairmaire, 1891

murrayi Boheman, 1862: Cameroon: 12, Central African Rep.: 6, Dem. Rep. of Congo: 17; Total **35**

Subgenus *Sindia* Weise, 1897

depressa Świętojańska, 2001 c: 267: India: 4 (**1 HT, 3 PT**); Total **4**

sulcata (Olivier, 1808): India: 17; Total **17**

Subgenus *Sindiola* Spaeth, 1903

burmensis (Spaeth, 1938): 231: Myanmar: 7 (**1 ST**), Thailand: 4; Total **11**

hospita Boheman, 1855: China: 2, Laos: 5, Thailand: 6, Vietnam: 9; Total **22**

meghalayaensis Świętojańska, 2001 c: 280: India: 5 (**1 HT, 4 PT**); Total **5**

multinotata Boheman, 1855: Indonesia: 6; Total **6**

novemdecimnotata Boheman, 1855: Indonesia: 9; Total **9**

parallelipennis (Spaeth, 1903): Myanmar: 2; Total **2**

vigintisexnotata Boheman, 1855: India: 1, Laos: 1, Malaysia: 1, Myanmar: 1, Thailand: 8, Vietnam: 16, no country: 1; Total **29**

Subgenus *Sindiolina* Świętojańska, 2001

sedecimaculata (Boheman, 1856): India: 9, Laos: 1, Nepal: 2; Total **12**

Genus *Mahatsinia* Spaeth, 1919

nodulosa (Weise, 1910): Madagascar: 5; Total **5**

Genus *Nilgiraspis* Spaeth, 1932

andrewesi (Spaeth, 1932): India: 4; Total **4**

Tribe: Basiprionotini Hincks, 1952

Genus *Androya* Spaeth, 1911

longula (Fairmaire, 1901): 84: Madagascar: 4 (1 ST); Total 4
obscuricollis (Fairmaire, 1903): 10: Madagascar: 7 (1 ST); Total 7
rubrocostata (Fairmaire, 1898): 258: Madagascar: 13 (1 ST); Total 13
tenuecostata (Fairmaire, 1899): 556: Madagascar: 4 (1 ST); Total 4

Genus *Basiprionota* Chevrolat, 1836

amboinica (Spaeth, 1925): Indonesia: 2; Total 2
amitina (Spaeth, 1932): Thailand: 1; Total 1
angusta (Spaeth, 1914): Taiwan: 4; Total 4
atricornis (Spaeth, 1912): Malaysia (Borneo): 1; Total 1
bakeri (Spaeth, 1925): Philippines: 2; Total 2
bimaculata (Thunberg, 1789): China: 1, Laos: 2, Myanmar: 2, Thailand: 1; Total 6
bisignata (Boheman, 1862): China: 15, Vietnam: 3; Total 18
cerata (Spaeth, 1925): Vietnam: 5; Total 5
chinensis (Fabricius, 1798): China: 12, Vietnam: 7; Total 19
decemmaculata (Boheman, 1850): India: 6 (1 ST *Prioptera 10-signata* Boheman, 1850: 62), Indonesia: 2, Laos: 1, Malaysia: 1, Myanmar: 2, Nepal: 3, Thailand: 5; Total 20
decempustulata (Boheman, 1850): Indonesia: 8, Malaysia: 2; Total 10
flavicornis Borowiec, 1993: 84: Vietnam: 1 (1 PT); Total 1
gressitti Medvedev, 1957: China: 2; Total 2
immaculata (Wagner, 1881): Philippines: 5; Total 5
impacata (Spaeth, 1925): Indonesia: 2; Total 2
lata Chen & Zia, 1964: China: 1; Total 1
latissima (Wagner, 1881): Philippines: 6; Total 6
maerkeli (Boheman, 1850): Indonesia: 1; Total 1
morigera (Spaeth, 1925): Philippines: 1; Total 1
multiplagiata (Wagner, 1881): India (Andanam Is.): 3; Total 3
nigricollis (Weise, 1897): Indonesia: 3; Total 3
octomaculata (Boheman, 1850): Indonesia: 1; Total 1
octonotata (Boheman, 1850): Indonesia: 1; Total 1
octopunctata (Fabricius, 1787): Indonesia: 7; Total 7
octopustulata (Boheman, 1856): Philippines: 4; Total 4
opima (Spaeth, 1925): China: 1; Total 1
privigna (Boheman, 1862): Indonesia: 1; Total 1
prognata (Spaeth, 1925): China: 1, Vietnam: 1; Total 2
pudica (Spaeth, 1925): China: 5; Total 5
quadriimpressa (Boheman, 1850): Indonesia: 8; Total 8
ramigera (Boheman, 1862): 25: Indonesia: 1 (1 ST); Total 1
rugosipennis (Spaeth, 1901 c): 3: Indonesia: 1 (1 ST); Total 1
sarawacensis (Spaeth, 1912): Malaysia (Borneo): 6; Total 6
schultzei (Weise, 1908): Philippines: 2; Total 2
secretata (Spaeth, 1925): China: 5, Vietnam: 3; Total 8
sexmaculata (Boheman, 1850): India: 2, Nepal: 1, Thailand: 10; Total 13
sinuata (Olivier, 1790): Philippines: 10; Total 10
sospes (Spaeth, 1925): Philippines: 5; Total 5
subopaca (Spaeth, 1925): Philippines: 3; Total 3
sumba Borowiec, 2006 b: 245: Indonesia: 2 (1 HT, 1 PT); Total 2
tibetana (Spaeth, 1914): China: 3; Total 3
westermanni (Mannerheim, 1844): China: 1, India: 1, Myanmar: 1, Thailand: 13, Vietnam: 1; Total 17
whitei (Boheman, 1856): China: 10; Total 10

Genus *Cassidopsis* Fairmaire, 1899

basipennis Fairmaire, 1899: 557: Madagascar: 3 (1 ST); Total 3

perrieri Fairmaire, 1900: 89: Madagascar: 3 (1 ST); Total 3

Genus *Craspedonta* Chevrolat, 1837

andrewesi (Weise, 1897): India: 10; Total 10

laotica Świątojańska & Borowiec, 2000 a: Laos: 4 (1 HT); Total 4

leayana (Latreille, 1807): Ceylon: 1, India: 10, Laos: 4, Thailand: 4, Vietnam: 15, no country: 1; Total: 35

levis Borowiec, 1993: 79: China: 6 (6 PT); Total 6

mouhoti (Baly, 1863): Cambodia: 5, Vietnam: 1; Total 6

obscura (Weise, 1897): Myanmar: 5; Total 5

Genus *Epistictina* Hincks, 1950

fulvonigra (Maulik, 1913): Laos: 2; Total 2

perplexa (Baly, 1863): Myanmar: 1, Thailand: 15; Total 16

reicheana (Guérin, 1844): India: 11; Total 11

viridimaculata (Boheman, 1850): China: 4, India: 5, Laos: 3, Malaysia: 2, Myanmar: 3, Thailand: 14, Vietnam: 3;

Total 34

Genus *Megapyga* Boheman, 1850

angulicollis Spaeth, 1898: Indonesia: 2; Total 2

brevis Spaeth, 1904: Indonesia: 1; Total 1

chinensis Spaeth, 1936: Thailand: 1; Total 1

coeruleomaculata Boheman, 1850: Philippines: 6; Total 6

eximia Boheman, 1850: 43: Philippines: 6 (1 ST); Total 6

minima Borowiec, 1998 a: 103: Indonesia: 1 (1 HT); Total 1

obscuricollis Borowiec, 1993: 80: Philippines: 1 (1 HT); Total 1

terminalis Boheman, 1862: 16: Philippines: 1 (1 ST); Total 1

Genus *Metriopepla* Fairmaire, 1882

inornata (Waterhouse, 1877): Dem. Rep. of Congo: 2, Mozambique: 2, Tanzania: 2; Total 6

Genus *Pseudandroya* Spaeth, 1952

livingstonei Spaeth, 1952: Tanzania: 1; Total 1

Tribe Cassidini

Old World genera

Genus *Acrocassis* Spaeth, 1922

Subgenus s. str.:

flavescens Weise, 1904: Kenya: 15; Total 15

gibbipennis (Boheman, 1854): Botswana: 2, Malawi: 1, Namibia: 1, RSA: 6, Tanzania: 4, Zambia: 1, Zimbabwe: 1; Total 16

intermedia Borowiec, 2002 i: 28: Angola: 3 (3 PT); Total 3

roseomarginata (Boheman, 1854): Burkina Faso: 11, Cameroon: 1, Danmark (introduced!): 1, Dem. Rep. of Congo: 2, Gabon: 2, Ghana: 5, Ivory Coast: 2, Sierra Leone: 1, Tanzania: 3; Total 28

rufula (Thomson, 1858): Burkina Faso: 1, Cameroon: 5, Dem. Rep. of Congo: 6, Nigeria: 3, Tanzania: 2, Uganda: 1; Total 18

sudanensis Spaeth, 1929: Senegal: 3, Yemen: 2; Total 5

undulatipennis Borowiec, 2002 i: 45: Namibia: 1 (1 HT); Total 1

zavattarii Spaeth, 1938: Ethiopia: 1; Total **1**

Subgenus *Bassamia* Spaeth, 1924

paeminosa (Boheman, 1856): Gabon: 2, Ivory coast: 1, no country: 1; Total **4**

Genus *Aethiopocassis* Spaeth, 1922

angolensis Spaeth, 1926: Angola: 1; Total **1**

angulicollis Borowiec & Świetojańska, 2018 c: 10: Tanzania: 1 (**1 PT**); Total **1**

cinctidorsum (Weise, 1912): Dem. Rep. of Congo: 39, Rwanda: 2, Tanzania: 1, Uganda: 2; Total **44**

dewittei Borowiec & Świetojańska, 2018 c: 15: Dem. Rep. of Congo: 16 (**16 PT**); Total **16**

flaccida (Spaeth, 1924): Kenya: 1; Total **1**

fugax (Spaeth, 1906): Botswana: 1, Dem. Rep. of Congo: 1, Malawi: 4, Zambia: 1, Zimbabwe: 2; Total **9**

gallarum (Spaeth, 1906): Dem. Rep. of Congo: 1, Erytrea: 1, Ethiopia: 8, Kenya: 2, Malawi: 4, South Sudan: 4,

Tanzania: 21; Total **41**

garambana Borowiec & Świetojańska, 2018 c: 24: Dem. Rep. of Congo: 20 (**20 PT**); Total **20**

guineensis Borowiec & Świetojańska, 2018 c: 25: Guinea: 2 (**2 PT**); Total **2**

imbuta (Spaeth, 1924): 331: Dem. Rep. of Congo: 13 (**1 ST**), Mozambique: 1, Zambia: 4; Total **18**

kasaiensis (Spaeth, 1933): Equatorial Guinea: 1; Total **1**

kraatzii (Weise, 1898): Cameroon: 28; Total **28**

longidoana Borowiec & Świetojańska, 2018 c: 32: Tanzania: 1 (**1 PT**); Total **1**

pauli (Weise, 1898): Tanzania: 10; Total **10**

plagipennis (Spaeth, 1916): Rwanda: 1, Tanzania: 8; Total **9**

planipennis (Spaeth, 1933): Tanzania: 1; Total **1**

quadrioculata (Spaeth, 1916): Burundi: 1, Dem. Rep. of Congo: 6, Rwanda: 1, Uganda: 6; Total **14**

revulsa (Spaeth, 1906): Dem. Rep. of Congo: 40, Kenya: 1, Malawi: 3, Mozambique: 4, Tanzania: 4, Zambia: 2, no country: 1; Total **55**

rhodesiana (Spaeth, 1924): Dem. Rep. of Congo: 1, Malawi: 6, Zambia: 11; Total **18**

scripta (Fabricius, 1798): RSA: 2; Total **2**

sensualis (Spaeth, 1924): Dem. Rep. of Congo: 1; Total **1**

silphoides (Spaeth, 1906): Dem. Rep. of Congo: 8, Tanzania: 8, Zambia: 1, no country: 1; Total **18**

sparsuta (Weise, 1904): Kenya: 2, Tanzania: 3; Total **5**

steindachneri (Spaeth, 1916): Dem. Rep. of Congo: 2; Total **2**

stictica (Harold, 1879): Angola: 6; Total **6**

suaheorum (Weise, 1899): Malawi: 4, Tanzania: 22, Zimbabwe: 4; Total **30**

suspiciosa (Weise, 1903): Dem. Rep. of Congo: 85 (**1 PT** *Aethiopocassis maynei* Spaeth, 1933: 351), Kenya: 1, Malawi: 17, Mozambique: 2, Tanzania: 9, Zambia: 47, Zimbabwe: 8; Total **169**

transvaalensis Borowiec & Świetojańska, 2018 c: 61: RSA: 2 (**2 PT**); Total **2**

vigintimaculata (Thunberg, 1789): Mozambique: 1, RSA: 22; Total **23**

Aethiopocassis sp.: Dem. Rep. of Congo: 2, Tanzania: 8; Total **10**

Genus *Andevocassis* Spaeth, 1924

picta (Spaeth, 1905): Madagascar: 6; Total **6**

Genus *Austropsecadia* Hincks, 1950

chlorina (Weise, 1899): Australia: 7; Total **7**

Genus *Basipta* Chevrolat, 1849

glauca Chevrolat, 1849: RSA: 20; Total **20**

luteocincta Boheman, 1854: RSA: 2; Total **2**

pilosella Boheman, 1854: RSA: 4; Total **4**

sebastiani Borowiec, 2002 i: 69: RSA: 4 (**4 PT**); Total **4**

stolida Boheman, 1854: RSA: 13; Total **13**

Genus *Capelocassis* Spaeth, 1952

dorsata (Duvivier, 1891): India: 6; Total **6**

Genus *Cassida* Linnaeus, 1758

achardi (Spaeth, 1926): Vietnam: 3; Total **3**

aethiopica Boheman, 1854: Cameroon: 1, Chad: 1, Dem. Rep. of Congo: 1, Ghana: 1, Mali: 6, Togo: 1; Total **11**

agilis Spaeth, 1915: Madagascar: 2; Total **2**

albertisi Spaeth, 1903: Papua New: 4; Total **4**

algerica Lucas, 1849: Algeria: 13, Italy: 9, Spain: 3; Total **25**

alidagiense Özdikmen, Sahin, & Bal, 2020: Turkey: 2; Total **2**

alpina Bremsi-Wolf, 1855: France: 5, Italy: 7; Total **12**

altiuscula Spaeth, 1916: Cameroon: 2, Dem. Rep. of Congo: 2, Equatorial Guinea: 1; Total **5**

amaranthica Medvedev & Eroshkina, 1988: 140: Vietnam: 5 (**2 PT**), Thailand: 1; Total **6**

ambrica Borowiec, 1999 j: 450: Madagascar: 1 (**1 HT**); Total **1**

amorifica Boheman, 1862: Dem. Rep. of Congo: 1, Namibia: 8, Rwanda: 2, RSA: 2, Zambia: 1; Total **14**

amurensis (Kraatz, 1879): China: 1, North Korea: 1, Russia: 8; Total **10**

ancorifera Boheman, 1856: Central African Rep.: 1, Dem. Rep. of Congo: 4, Tanzania: 1, Zambia: 1, Zimbabwe: 5; Total **12**

andapaensis Borowiec, 1988 b: 560: Madagascar: 1 (**1 HT**); Total **1**

andohahelana Borowiec & Świętojańska, 2013 h: 33: Madagascar: 3 (**1 HT, 2 PT**); Total **3**

andreinii Spaeth, 1933: Nigeria: 1, RSA: 4; Total **5**

andrewesi Weise, 1897: 111: India: 4 (**1 ST**); Total **4**

angulicollis Borowiec & Świętojańska, 2013 h: 35: Madagascar: 1 (**1 HT**); Total **1**

angusta Marseul, 1876: Algeria: 1, Morocco: 4; Total **5**

angustifrons Weise, 1891: Klein-Asien (mislabeled!): 1; Total **1**

anosyennesensis Borowiec & Świętojańska, 2013 h: 37: Madagascar: 1 (**1 HT**); Total **1**

antoni Borowiec & Świętojańska, 1997 d: 278: India: 8 (**1 HT, 1 PT**); Total **8**

appluda Spaeth, 1926 c: 86: China: 1, Laos: 1, Thailand: 1, Vietnam: 1 (**1 ST**); Total **4**

astrolabiana (Spaeth, 1903): Papua New Guinea: 11; Total **11**

atrata Fabricius, 1787: Austria: 4, Bulgaria: 6, Greece: 1, Poland: 6, Ukraine: 1; Total **18**

atroannulus Borowiec & Świętojańska, 2013 h: 39: Madagascar: 1 (**1 PT**); Total **1**

atrofemorata Borowiec & Sassi, 2002 c: 145: China: 1 (**1 HT**); Total **1**

atromarginata Borowiec & Świętojańska, 2013 h: 40: Madagascar: 14 (**5 PT**); Total **14**

atropunctata Borowiec & Świętojańska, 2013 h: 42: Madagascar: 2 (**1 HT, 1 PT**); Total **2**

atrorubra Borowiec, 1999 j: 470: Madagascar: 3 (**1 HT, 1 PT**); Total **3**

aureola (Spaeth, 1915): Australia: 2; Total **2**

auropustulata (Fairmaire, 1899): Madagascar: 5; Total **5**

aurora Weise, 1907: Austria: 8, Poland: 1, Ukraine: 1; Total **10**

australica (Boheman, 1855): China: 1, India: 9, Nepal: 2, Thailand: 3; Total **15**

avia (Weise, 1897): India: 3; Total **3**

azurea Fabricius, 1801: Albania: 1, France: 20, Germany: 1, Hungary: 3, Slovakia: 8, Turkey: 1; Total **34**

bakeri Spaeth, 1925: Philippines: 2; Total **2**

bamendana Borowiec & Świętojańska, 2021 (in litt.): Cameroon: 1 (**1 PT**); Total **1**

bella Faldermann, 1837: Georgia: 2, Russia: 5, Turkey: 2; Total **9**

belli Weise, 1897: India: 11; Total **11**

belliformis Maulik, 1919: India: 3; Total **3**

benaadirensis Borowiec & Świętojańska, 2021 (in litt.): Somalia: 1 (**1 HT**); Total **1**

beniowski Borowiec, 1988 b: 552: Madagascar: 1 (**1 HT**); Total **1**

beondrokana Borowiec & Świętojańska, 2013 h: 49: Madagascar: 2 (**2 PT**); Total **2**

bergeali Bordy, 1995 b: 244: Bulgaria: 1, Czech Rep.: 1, France: 3 (**1 PT**), Poland: 5, Slovakia: 2; Total **12**

berolinensis Suffrian, 1844: Czech Rep.: 2, Mongolia: 4, Poland: 1, Russia: 1, no country: 1; Total **9**

bezdeki Borowiec, 2002 h: 567: Laos: 2 (**1 HT**); Total **2**

bicallosa Spaeth, 1915: Madagascar: 11; Total **11**
biguttata (Spaeth, 1903): Papua New Guinea: 14; Total **14**
brevis Weise, 1884: Afghanistan: 6, Lebanon: 1, Turkey: 32; Total **39**
brooksi Borowiec, 1992 a: 283: Madagascar: 7 (**1 PT**); Total **7**
bulirschi Borowiec & Świętojańska, 2013 h: 55: Madagascar: 2 (**1 HT, 1 PT**); Total **2**
butterwecki Borowiec, 2007 d: 48: Madagascar: 1 (**1 PT**); Total **1**
callosicollis Spaeth, 1926: Tanzania: 4; Total **4**
calvaria (Weise, 1900): Dem. Rep. of Congo: 1, Zambia: 1; Total **2**
camerunensis (Spaeth, 1903): Cameroon: 3; Total **3**
canaliculata Laicharting, 1781: Austria: 4, Germany: 1, Romania: 1, Russia: 18, Slovakia: 4, Ukraine: 5, no country: 1; Total **34**
capensis Borowiec, 2005 f: 116: RSA: 2 (**2 PT**); Total **2**
catenata (Boheman, 1855): India: 7, Indonesia: 44, Malaysia (continental and Borneo): 17, Vietnam: 2; Total **70**
ceylonica (Boheman, 1855): Ceylon: 2; Total **2**
cherrapunjiensis Maulik, 1919: India: 2, Laos: 7, Vietnam: 1; Total **10**
chiangmaiensis Borowiec, 2001 k: 551: Thailand: 2 (**1 HT, 1 PT**); Total **2**
chrysanthemoides Borowiec & Świętojańska, 2001 a: 155: RSA: 4 (**4 PT**); Total **4**
circumdata Herbst, 1799: China: 1, India: 10 (**1 HT** *Cassia nilgiriensis* Borowiec & Takizawa, 1991 d: 642), Indonesia: 7, Laos: 1, Malaysia (continental and Borneo): 6, Philippines: 29, Thailand: 3, Vietnam: 22, USA (Hawaii, introduced): 3; Total **82**
circumsepta Spaeth, 1915: Madagascar: 1; Total **1**
coagulata Boheman, 1854: Kenya: 1, Malawi: 1, RSA: 5, Tanzania: 3, Zambia: 1, Zimbabwe: 1; Total **12**
coelebs Borowiec, 1999 j: 455: Madagascar: 2 (**2 PT**); Total **2**
cognobilis Spaeth, 1926: Thailand: 7; Total **7**
collucens Spaeth, 1915: Madagascar: 14; Total **14**
compuncta (Boheman, 1855): Australia: 4, Fiji (introduced): 19; Total **23**
concallescens Spaeth, 1915: Madagascar: 1; Total **1**
concha Solsky, 1872: North Korea: 1, South Korea: 3, Taiwan: 1; Total **5**
conchyliata (Spaeth, 1914): China: 4, India: 2, Malaysia: 1, Thailand: 1, Vietnam: 3; Total **11**
consobrina Spaeth, 1915: Madagascar: 3; Total **3**
conspurcata Boheman, 1854: India: 9; Total **9**
contracta (Spaeth, 1915): Madagascar: 2; Total **2**
corallina Boheman, 1862: Algeria: 1, Spain: 3; Total **4**
corbetti (Weise, 1897): China: 1, Laos: 1; Total **2**
cordula Boheman, 1854: RSA: 1; Total **1**
corollata Spaeth, 1940: Indonesia: 4, Malaysia (Borneo): 4; Total **8**
crucifera (Kraatz, 1879): China (Hongkong): 1, Japan: 4, Laos: 3, Taiwan: 1, Vietnam: 1; Total **10**
culminis (Chen & Zia, 1964): India: 1, Laos: 8, Myanmar: 1, Thailand: 1; Total **11**
currax Spaeth, 1915: Madagascar: 8; Total **8**
deflorata Suffrian, 1844: Algeria: 1, France: 5, Italy: 4, Morocco: 3, Spain: 1, no country: 1; Total **15**
delenifica Boheman, 1862: Benin: 1, Cameroon: 4; Total **5**
delesserti Boheman, 1854: India: 17; Total **17**
dembickyi Borowiec, 2001 k: 545: Thailand: 2 (**1 HT, 1 PT**); Total **2**
densestriata Borowiec & Świętojańska, 2013 h: 73: Madagascar: 2 (**1 HT, 1 PT**); Total **2**
denticollis Suffrian, 1844: Poland: 58, Slovakia: 1, Ukraine: 2; Total **61**
denticulata Boheman, 1856: Australia: 7; Total **7**
depicta Boheman, 1862: Cameroon: 4, Dem. Rep. of Congo: 9, Ethiopia: 3, Kenya: 3, Tanzania: 2, Togo: 1, Uganda: 2; Total **24**
derasa Spaeth, 1940: China: 1, Vietnam: 4; Total **5**
desultrix (Spaeth, 1914): China: 1, India: 8, Laos: 8, Nepal: 1, Thailand: 3; Total **21**
devalaensis Borowiec & Takizawa, 1991 d: 642: India: 4 (**1 PT**); Total **4**
devylderi Spaeth, 1928: Botswana: 1, Namibia: 5, Zimbabwe: 1; Total **7**

dionna Boisduval, 1835: Australia: 3, Indonesia (New Guinea): 1, Papua New Guinea: 49, Samoa (introduced): 3;
 Total **56**
distinguenda Spaeth, 1928: RSA: 4; Total **4**
diversepunctata Borowiec & Świętojańska, 2001 a: 157: RSA: 2 (**1 PT**); Total **2**
dolens Borowiec, 1999 j: 465: Madagascar: 6 (**6 PT**); Total **6**
dorsonotata Boheman, 1854: India: 9; Total **9**
dorsovittata Boheman, 1854: Dem. Rep. of Congo: 3, Madagascar: 6, Namibia: 1, RSA: 17, Rwanda: 1, Uganda:
 1, Zambia: 1; Total **30**
dulcis (Boheman, 1862): Madagascar: 1; Total **1**
ellipticollis Spaeth, 1914: India: 7; Total **7**
eluta Boheman, 1862: Ethiopia: 2, South Sudan: 1; Total **3**
emontinensis Borowiec & Świętojańska, 2021 (in litt.): RSA: 3 (**1 HT, 2 PT**); Total **3**
eo (Spaeth, 1928): China: 1 (**1 ST**), Laos: 2; Total **3**
exilis (Boheman, 1854): India: 15; Total **15**
eximia Borowiec & Ghate, 2004 c: 250: India: 3 (**1 HT, 2 PT**); Total **3**
expressa (Spaeth, 1914): China: 6; Total **6**
fausti Spaeth & Reitter, 1926: Bulgaria: 1, Turkey: 2; Total **3**
fae Spaeth, 1904: Laos: 2, Myanmar: 1, Thailand: 3; Total **6**
ferranti Spaeth, 1915: Madagascar: 5; Total **5**
ferruginea Goeze, 1777: Bulgaria: 1, Czech Rep.: 2, France: 6, Italy: 2, Poland: 1, Russia: 2, Slovenia: 1, South
 Korea: 1, Spain: 1, no country: 1; Total **18**
flaveola Thunberg, 1794: Bulgaria: 2, Finland: 1, Poland: 32, Russia: 1 (**1 PT** *Cassia kusnetzovi* Matis, 1974: 1101);
 Total **36**
flavoguttata Spaeth, 1914: India: 3; Total **3**
flavoscutata Spaeth, 1914: India: 1; Total **1**
flavosignata Spaeth, 1932: Tanzania: 2; Total **2**
foveolatipennis Borowiec & Świętojańska, 2001 a: 157: RSA: 1 (**1 PT**); Total **1**
franklinmuelleri Spaeth, 1925: RSA: 5, Zimbabwe: 1; Total **6**
frontalis Boheman, 1856: Madagascar: 13; Total **13**
fukhanica Medvedev & Eroshkina, 1988: 136: Vietnam: 2 (**1 PT**); Total **2**
fumida Spaeth, 1914: Myanmar: 1; Total **1**
fuscocaula Borowiec, 1988: Madagascar: 2; Total **2**
fuscorufa Motschulsky, 1866: China: 8, Japan: 2, North Korea: 11, Russia: 3, South Korea: 6; Total **30**
fuscocaula Boheman, 1854: Gambia: 1, Senegal: 1; Total **2**
gansuica Chen & Zia, 1964: China: 2; Total **2**
gentilis Spaeth, 1926 c: 87: Vietnam: 12 (**1 ST**); Total **12**
ghesquieri Spaeth, 1943: Dem. Rep. of Congo: 1; Total **1**
glebicolor Spaeth, 1926 d: 14: Myanmar: 1 (**1 ST**); Total **1**
goudoti (Boheman, 1855): Madagascar: 20; Total **20**
granula Boheman, 1856: Dem. Rep. of Congo: 9, Ghana: 1; Total **10**
granulicollis Spaeth, 1905: RSA: 13; Total **13**
guttipennis Boheman, 1862: RSA: 4; Total **4**
hablitziae Motschulsky, 1838: Georgia: 14, Turkey: 4; Total **18**
hemisphaerica Herbst, 1799: Bulgaria: 3, France: 1, Poland: 10, Spain: 2; Total **16**
heroni Borowiec & Świętojańska, 2021 (in litt.): Mozambique: 3 (**3 PT**), RSA: 3 (**1 HT, 2 PT**); Total **6**
hexastigma Suffrian, 1844: France: 11, Italy: 1, Morocco: 1, Portugal: 6, no country: 2; Total **21**
hova (Weise, 1910): Madagascar: 3; Total **3**
hovacassiformis Borowiec, 1999 j: 447: Madagascar: 1 (**1 HT**); Total **1**
humeralis Kraatz, 1874: France: 2, Portugal: 2, Spain: 1; Total **5**
humerosa Spaeth, 1902: Cameroon: 3, Dem. Rep. of Congo: 12, Rwanda: 2, South Sudan: 1, Uganda: 63; Total **81**
hyalina Weise, 1891: Spain: 9; Total **9**
icterica Boheman, 1854: India: 7 (**1 ST** *Cassia petulans* Spaeth, 1914 b: 550), Nepal: 4; Total **11**

ictericiformis Borowiec, 2009 f: 476: India: 3 (**1 HT, 2 PT**); Total **3**
imbecilla (Boheman, 1862): Ceylon: 3; Total **3**
imitatrix Spaeth, 1916: Dem. Rep. of Congo: 1, Ethiopia: 2, Rwanda: 1, Uganda: 9; Total **13**
impressipennis Borowiec & Świętojańska, 2013 h: 95: Madagascar: 2 (**1 HT, 1 PT**); Total **2**
inaequalis Thomson, 1858: Cameroon: 12; Dem. Rep. of Congo: 5, Rep. of Central Africa: 2, Uganda: 4; Total **23**
incallida Spaeth, 1938: Laos: 2, Thailand: 1, Vietnam: 1; Total **4**
inciens Spaeth, 1926 c: 85: Laos: 1, Myanmar: 1, Vietnam: 4 (**1 ST**); Total **6**
inconstans (Fairmaire, 1899): 506: Madagascar: 2 (**1 ST**); Total **2**
indochinensis (Spaeth, 1919): China: 2, Laos: 3, Myanmar: 1, Vietnam: 8; Total **14**
inflaccens Spaeth, 1940: Laos: 1, Vietnam: 1; Total **2**
inflata Gressitt, 1952: China: 2; Total **2**
informis Boheman, 1862: India: 26; Total **26**
innotata Boheman 1854: Kenya: 2, Malawi: 1, Namibia: 2, Nigeria: 1, RSA: 5, Tanzania: 1, Zambia: 2; Total **14**
inopinata Sassi & Borowiec, 2006 h: 547: Greece: 1 (**1 PT**), Italy: 2 (**2 PT**); Total **3**
inquinata Brullé, 1832: Algeria: 4, Bulgaria: 8, France: 3, Italy: 3, Morocco: 13, Portugal: 2, Spain: 4, Tunisia: 1;
 Total **38**
insulana Gressitt, 1952: Taiwan: 4; Total **4**
irregularis Boheman, 1854: RSA: 17; Total **17**
irrorata Weise, 1898: Kenya: 1, Tanzania: 4; Total **5**
isarogensis Borowiec, 2009 f: 479: Philippines: 1 (**1 HT**); Total **1**
jacobsoni Spaeth, 1914: China: 14; Total **14**
japana Baly, 1874: China: 6, Japan: 9, South Korea: 2, Vietnam: 9; Total **26**
juglans Gressitt, 1942: Russia: 1; Total **1**
justa Spaeth, 1914: India: 3; Total **3**
kantnerorum Borowiec & Świętojańska, 2021 (in litt.): Malawi: 40 (**1 HT, 39 PT**); Total **40**
kinabaluensis Borowiec, 1999 a: 2: Malaysia (Borneo): 1 (**1 HT**); Total **1**
koreana Borowiec & Cho, 2011 b: 447: South Korea: 4 (**1 HT, 1 PT**); Total **4**
kunminica (Chen & Zia, 1964): Thailand: 2; Total **2**
labiatophaga Medvedev & Eroshkina, 1988: 132: Vietnam: 1 (**1 PT**); Total **1**
laccopteroides Borowiec & Świętojańska, 2013 h: 100: Madagascar: 3 (**1 HT, 2 PT**); Total **3**
lacrymosa Boheman, 1854: Dem. Rep. of Congo: 7, Kenya: 3, Namibia: 1, RSA: 6, Rwanda: 1, Uganda: 1,
 Zimbabwe: 2; Total **21**
laetabilis Spaeth, 1915: Madagascar: 4; Total **4**
langeri Borowiec, 2009 f: 474: Myanmar: 1 (**1 HT**); Total **1**
laotica Borowiec, 2002 h: 569: Laos: 3 (**1 HT**); Total **3**
latecincta Fairmaire, 1904: Madagascar: 17; Total **17**
lateritia Fairmaire, 1904: Madagascar: 18; Total **18**
lateritioides Borowiec & Świętojańska, 2013 h: 108: Madagascar: 8 (**8 PT**); Total **8**
leleupi Borowiec & Świętojańska, 2021 (in litt.): Rwanda: 2 (**2 PT**); Total **2**
leucanthemi Bordy, 1995 a: 377: Czech Rep.: 1, France: 5 (**1 PT**), Italy: 1, Poland: 5, Switzerland: 2; Total **14**
liliputana Borowiec & Świętojańska, 2013 h: 110: Madagascar: 1 (**1 HT**); Total **1**
limpopoana Borowiec & Świętojańska, 2001 a: 158: Malawi: 1, Mozambique: 1, Zimbabwe: 3 (**5 PT**); Total **5**
lineola Creutzer, 1799: Bulgaria: 10, China: 1, Kazakhstan: 12, Mongolia: 3, Poland: 3, Romania: 2, Russia: 5,
 South Korea: 2, Ukraine: 2; Total **40**
linnavuorii Borowiec, 1986: Lebanon: 3; Total **3**
litigiosa Boheman, 1854: Botswana: 2, Namibia: 9, RSA: 20, Zambia: 2, Zimbabwe: 1; Total **34**
lombocensis (Spaeth, 1919): Indonesia: 10; Total **10**
luxuriosa Spaeth, 1940: Cameroon: 1, Kenya: 1; Total **2**
lycii Borowiec & Świętojańska, 2001 a: 160: RSA: 2 (**1 PT**); Total **2**
lyrica Fairmaire, 1904: Madagascar: 1; Total **1**
madagascariensis Borowiec, 1999 j: 452: Madagascar: 1 (**1 HT**); Total **1**
major Kraatz, 1874: Algeria: 9, Spain: 1; Total **10**

malaysiana Borowiec, 2010 b: 280: Malaysia: 10 (**1 HT, 9 PT**); Total **10**
mandli Spaeth, 1921: Russia: 6, South Korea: 8; Total **14**
manipuria Maulik, 1923: Myanmar: 2; Total **2**
margaritacea Schaller, 1783: Bulgaria: 6, Croatia: 1, Poland: 21, Turkey: 2; Total **30**
mariaeadelheidae Spaeth, 1915: Madagascar: 4; Total **4**
mashonensis Spaeth, 1928: Kenya: 1, Namibia: 1; Total **2**
medvedevi Lopatin, 1965: Kyrgyzstan: 1, Uzbekistan: 2; Total **3**
melanophthalma Boheman, 1854: Namibia: 4; RSA: 23; Total **27**
mera Germar, 1848: Australia: 13; Total **13**
mindanaoensis Spaeth, 1933: Philippines: 12; Total **12**
minutissima Borowiec & Świętojańska, 2021 (in litt.): Somalia: 3 (**1 HT, 2 PT**); Total **3**
mishmiensis Sekerka & Borowiec, 2008 d: 612: India: 1 (**1 PT**); Total **1**
mongolica Boheman, 1854: China: 4, Mongolia: 4, South Korea: 1; Total **9**
moori Boheman, 1856: Kyrgyzstan: 4; Total **4**
morondaviana Borowiec, 2007 d: 51: Madagascar: 7 (**1 HT, 6 PT**); Total **7**
mroczkowskii Borowiec & Świętojańska, 1997 d: 275: Vietnam: 1 (**1 HT**); Total **1**
multistrigata Borowiec & Świętojańska, 2013 h: 124: Madagascar: 1 (**1 PT**); Total **1**
murraea Linnaeus, 1767: Bulgaria: 10, Mongolia: 1, Poland: 16, Russia: 10, Turkey: 1; Total **38**
murraea ssp. *halophila* Spaeth, 1926: Russia: 3, Ukraine: 2; Total **5**
mysorensis Borowiec & Świętojańska, 1996 e: 467: India: 1 (**1 HT**); Total **1**
namibiensis Borowiec, 2005d f: 119: Namibia: 1 (**1 PT**); Total **1**
natalensis Spaeth, 1932: RSA: 7; Total **7**
navicella Boheman, 1862: Australia: 3; Total **3**
nebulosa Linnaeus, 1758: Bulgaria: 3, China: 1, Czech Rep.: 1, France: 1, Poland: 15, Russia: 48, Slovakia: 3;
 Total **72**
nepalica Medvedev, 1997: India: 3, Nepal: 2; Total **5**
nigeriana Borowiec & Świętojańska, 2021 (in litt.): Nigeria: 2 (**1 HT, 1 PT**); Total **2**
nigriventris Boheman, 1854: India: 20, Nepal: 4, Vietnam: 2; Total **26**
nigrodentata Medvedev & Eroshkina, 1988: Thailand: 2; Total **2**
nigroflavens Borowiec, 1988: Madagascar: 2; Total **2**
nigrogibbosa Spaeth, 1901: Indonesia: 3; Total **3**
nigrohumeralis Borowiec & Ghate, 2004 c: 247: India: 1 (**1 HT**); Total **1**
nigropunctata Borowiec & Świętojańska, 2013 h: 129: Madagascar: 1 (**1 HT**); Total **1**
nigroscutata Fairmaire, 1904: Madagascar: 1; Total **1**
nobilis Linnaeus, 1758: Bulgaria: 1, Czech Rep.: 1, France: 14, Mongolia: 1, Poland: 15, Russia: 3, Slovakia: 1,
 Spain: 2; Total **38**
nosybeensis Borowiec & Świętojańska, 2013 h: 13: Madagascar: 1 (**1 HT**); Total **1**
nucula Spaeth, 1914: China: 2, Myanmar: 1, Vietnam: 1; Total **4**
nuwara Maulik, 1919: Ceylon: 2; Total **2**
nysea Spaeth, 1926: India: 3; Total **3**
obenbergeri (Spaeth, 1928): Thailand: 1 (**1 HT** *Cassia crucipennis* Borowiec, 2003 c: 121); Total **1**
oberlaenderi Spaeth, 1916: Cameroon: 1, Uganda: 14; Total **15**
obtusata Boheman, 1854: Ceylon: 4, India: 4, Indonesia: 3, Japan: 2, Laos: 11, Philippines: 3, Vietnam: 14;
 Total **41**
occursans Spaeth, 1914: India: 13; Total **13**
olympica Sekerka, 2005: 285: Greece: 9 (**9 PT**); Total **9**
oromiaensis Borowiec & Świętojańska, 2021 (in litt.): Ethiopia: 1 (**1 HT**); Total **1**
ovalis Spaeth, 1914: Tunisia: 11; Total **11**
overlaeti Borowiec & Świętojańska, 2021 (in litt.): Dem. Rep. of Congo: 1 (**1 ST**); Total **1**
pacholatkoii Sekerka & Borowiec, 2008 d: 616: India: 1 (**1 PT**); Total **1**
paiensis Borowiec, 2001 k: 548: Thailand: 5 (**1 HT, 4 PT**), Vietnam: 1; Total **6**
palaestina Reiche, 1858: Cyprus: 1, Iran: 13, Iraq: 1, Israel: 1, Kazakhstan: 1, Lebanon: 4, Turkey: 8, Turkmenistan:
 6, Ukraine: 2, Uzbekistan: 24; Total **61**

palijsi Matis, 1970: Kazakhstan: 1; Total **1**
pallidicollis Boheman, 1856: China: 3, Russia: 7, South Korea: 4; Total **14**
pannonica Suffrian, 1844: Austria: 2, Bulgaria: 17, China: 1, Croatia: 1, Germany: 2, Hungary: 1, Poland: 41, Russia: 9, Slovakia: 1, Turkey: 6, Ukraine: 1, Uzbekistan: 1; Total **83**
panzeri Weise, 1907: Austria: 1, Bulgaria: 1, France: 2, Poland: 15, Ukraine: 2; Total **21**
papuana (Spaeth, 1903): Indonesia (New Guinea): 7, Papua New Guinea: 10, Solomon Islands: 7; Total **24**
parvula Boheman, 1854: Bulgaria: 1, China: 12, Kazakhstan: 1; Total **14**
patruelis Spaeth, 1935: Dem. Rep. of Congo: 1, Zambia: 1; Total **2**
pauliani Borowiec, 1999 j: 442: Madagascar: 1 (**1 PT**); Total **1**
pauxilla Boheman, 1854: India: 1; Total **1**
pellegrini Marseul, 1868: Cyprus: 7, Israel: 2, Tunisia: 1; Total **10**
pernix Spaeth, 1917: 433: Dem. Rep. of Congo: 2 (**1 ST**); Total **2**
perplexa (Chen & Zia, 1961): India: 2, Laos: 1, Thailand: 3; Total **6**
persica Spaeth, 1926: Iran: 2, Iraq: 1; Total **3**
pfefferi Sekerka, 2006: 253: Cyprus: 49 (**2 PT**); Total **49**
physodes (Boheman, 1855): Indonesia: 14, Malaysia (continental and Borneo): 2, Philippines: 1, Singapore: 1, Thailand: 2, Vietnam: 5 (**1 ST** *Cassia expromta* Spaeth, 1914 b: 567); Total **25**
pipinata Hope, 1842: China: 9, Japan: 8, South Korea: 5; Total **22**
plausibilis (Boheman, 1862): China: 1, Malaysia: 2, Myanmar: 1, Thailand: 3; Total **7**
polymeriae Borowiec & Burwell, 2011 a: 286: Australia: 2 (**2 PT**); Total **2**
postarcuata (Chen & Zia, 1964): China: 1; Total **1**
praensis Borowiec, 2001 k: 556: Thailand: 2 (**1 HT, 1 PT**); Total **2**
praetimida Spaeth, 1912: Benin: 1, Dem. Rep. of Congo: 2, Ethiopia: 2, Gambia: 2, Ghana: 1, Kenya: 1, Nigeria: 1, Rep. of Central Africa: 3, Senegal: 2, Somalia: 1, Sudan: 1, Tanzania: 2, Yemen: 1; Total **20**
prasina Illiger, 1798: Bulgaria: 7, Czech Rep.: 1, Germany: 1, Hungary: 1, Kazakhstan: 1, Poland: 33, Russia: 9, Serbia: 1, Slovakia: 1, Turkey: 1, Ukraine: 5; Total **61**
probata Spaeth, 1914: China: 3; Total **3**
prospera Spaeth, 1915: Madagascar: 3; Total **3**
pseudolateritia Borowiec & Świętojańska, 2013 h: 141: Madagascar: 10 (**10 PT**); Total **10**
pseudosyratica Medvedev & Eroshkina, 1988: 138: Vietnam: 1 (**1 PT**), Thailand: 1; Total **2**
pubipennis Borowiec, 1999 j: 475: Madagascar: 1 (**1 HT**); Total **1**
pudens Boheman, 1854: RSA: 7; Total **7**
pulpa Spaeth, 1915: Madagascar: 4; Total **4**
pulvinata Boheman, 1854: India: 3; Total **3**
purpuricollis (Spaeth, 1914): China: 2; Total **2**
pusilla Walzl, 1839: Algeria: 2, Croatia: 1, France: 15, Italy: 3, Spain: 1; Total **22**
pusio Spaeth, 1915: Madagascar: 3; Total **3**
pyrenaica Weise, 1893: France: 2, Portugal: 3, Spain: 2, no country: 2; Total **9**
quadriramosa Gressitt, 1952: China: 3, Taiwan: 1; Total **4**
quatuordecimsignata Spaeth, 1899: RSA: 10; Total **10**
quinaria (Chen & Zia, 1964): Laos: 1, Vietnam: 1 (**1 PT** *Cassia quinqueasteriza* Medvedev & Eroshkina, 1988: 133); Total **2**
quinquemaculata Boheman, 1854: Philippines: 8; Total **8**
rabaiensis Borowiec & Świętojańska, 2021 (in litt.): Kenya: 1 (**1 HT**); Total **1**
rati Maulik, 1923: China: 5, Indonesia: 1, Myanmar: 1, Vietnam: 6; Total **13**
recondita (Boheman, 1862): Indonesia: 2, Philippines: 17; Total **19**
reitteri Weise, 1892: Armenia: 3, Turkey: 2; Total **5**
relicta Spaeth, 1927: USA: 3; Total **3**
reticulicosta (Chen & Zia, 1964): Laos: 1; Total **1**
reticulipennis Borowiec & Świętojańska, 2001 a: 162: RSA: 6 (**5 PT**); Total **6**
ribbei (Spaeth, 1919): Indonesia: 2; Total **2**
rimosa (Boheman, 1854): Madagascar: 12; Total **12**

rogezensis Borowiec & Świętojańska, 2013 h: 158: Madagascar: 1 (**1 HT**); Total **1**
rothschildi Spaeth, 1922: Kenya: 1; Sudan: 1, Yemen: 7; Total **9**
rubiginosa Müller, 1776: Bulgaria: 5, Canada: 2, China: 1, France: 5, Greece: 8, Italy: 7, Japan: 5, Kazakhstan: 5, Poland: 49, Russia: 4, Slovakia: 2, South Korea: 3, Spain: 1, Turkey: 6, Ukraine: 1, USA: 2; Total **106**
rubripennis Borowiec, 2002 h: 571: Laos: 1 (**1 HT**); Total **1**
rubromaculata Spaeth, 1918: Madagascar: 3; Total **3**
rubroornata (Boheman, 1855): Madagascar: 1; Total **1**
rudicollis (Spaeth, 1915): Madagascar: 9; Total **9**
rufomicans Fairmaire, 1904: Madagascar: 19; Total **19**
rufovirens Suffrian, 1844: Bosnia and Hercegovina: 1, Czech Rep.: 4, France: 7, Poland: 5, Russia: 1, Turkey: 1; Total **19**
rugipennis (Boheman, 1855): Madagascar: 6; Total **6**
ruralis (Boheman, 1862): Indonesia: 8, Malaysia: 1, Vietnam: 1; Total **10**
sabahensis Świętojańska & Borowiec, 2002 e: 360: Malaysia (Borneo): 1 (**1 HT**); Total **1**
saginata Spaeth, 1914: India: 6; Total **6**
salomonina (Spaeth, 1936): Papua New Guinea (Bougainville): 5, Solomon Islands: 6; Total **11**
sanguineoguttata Spaeth, 1915: Madagascar: 4; Total **4**
sanguinicollis (Spaeth, 1926): Madagascar: 1; Total **1**
sanguinolenta Müller, 1776: Armenia: 1, Austria: 1, Bulgaria: 5, Czech Rep.: 1, Poland: 24, Russia: 1, Slovakia: 1, Ukraine: 5; Total **39**
sanguinosa Suffrian, 1844: England: 1, France: 9, Poland: 24, Slovakia: 1; Total **35**
sappho (Boheman, 1862): Australia: 2; Total **2**
sareptana Kraatz, 1874: Kazakhstan: 2, Russia: 5; Total **7**
satanas Borowiec & Świętojańska, 2021 (in litt.): Uganda: 1 (**1 HT**); Total **1**
saucia Weise, 1889: Azerbaijan: 12, Turkey: 1, Turkmenistan: 5; Total **18**
sauteri (Spaeth, 1913): Taiwan: 11; Total **11**
schawalleri Medvedev, 1990: India: 1, Nepal: 1; Total **2**
schenklingi (Spaeth, 1915): Madagascar: 2; Total **2**
schoutedeni Spaeth, 1932: Dem. Rep. of Congo: 2, Kenya: 3, Uganda: 2; Total **7**
sculpturipennis Borowiec & Świętojańska, 2013 h: 158: Madagascar: 1 (**1 PT**); Total **1**
seladonia Gyllenhal, 1827: Algeria: 3, Bulgaria: 1, Czech Rep.: 1, France: 12, Germany: 2, Morocco: 1, Poland: 2, Spain: 4, no country: 1; Total **27**
senegalensis Borowiec & Świętojańska, 2021 (in litt.): Senegal: 3 (**3 PT**); Total **3**
senicula (Spaeth, 1915): Madagascar: 8; Total **8**
seniculoides Borowiec, 1999 j: 472: Madagascar: 1 (**1 PT**); Total **1**
seraphina Ménétries, 1836: Turkey: 10, no country: 1; Total **11**
setosa Chapuis, 1880: Tanzania: 1; Total **1**
sexguttata Boisduval, 1835: Indonesia (New Guinea): 5, Papua New Guinea: 35; Total **40**
shimba Borowiec & Świętojańska, 2021 (in litt.): Kenya: 3 (**1 HT, 2 PT**); Total **3**
sigillata (Gorham, 1885): South Korea: 5, Taiwan: 2; Total **7**
silvicola Borowiec, 1988 b: 562: Madagascar: 3 (**1 PT**); Total **3**
simanica (Chen & Zia, 1961): Thailand: 1; Total **1**
smaragdocruciata Medvedev & Eroshkina, 1982: 114: Vietnam: 1 (**1 PT**); Total **1**
socialis Spaeth, 1926 c: 86: India: 8 (**4 ST**); Total **8**
spaethi Weise, 1900: China: 1, North Korea: 3, Russia: 5; Total **9**
spaethiana (Gressitt, 1952): China: 8; Total **8**
sparteiformis Borowiec & Świętojańska, 2021 (in litt.): Angola: 1 (**1 HT**); Total **1**
spatiosa Spaeth, 1928: Namibia: 3; Total **3**
sphaerula Boheman, 1854: RSA: 5; Total **5**
spissa Weise, 1897: Ceylon: 1; Total **1**
stevensi Sekerka, 2011: India: 1 (**1 PT**); Total **1**
stigmatica Suffrian, 1844: Bulgaria: 1, Germany: 2, Kyrgyzstan: 1, Mongolia: 1, Poland: 69, Russia: 30, Slovakia: 1; Total **105**

stipidosa Borowiec & Świętojańska, 2021 (in litt.): Ethiopia: 1 (**1 PT**); Total **1**
strigaticollis Borowiec, 1988 b: 553: Madagascar: 1 (**1 PT**); Total **1**
strumosa (Spaeth, 1915): Madagascar: 2; Total **2**
suaveola (Spaeth, 1915): Madagascar: 1; Total **1**
subacuticollis Borowiec, 1999 j: 457: Madagascar: 9 (**1 PT**); Total **9**
sublesta (Weise, 1904): Dem. Rep. of Congo: 5, Kenya: 2, Uganda: 2, Zambia: 3; Total **12**
subprobata (Chen & Zia, 1964): Thailand: 6; Total **6**
subreticulata Suffrian, 1844: Austria: 1, Bulgaria: 3, Croatia: 2, France: 6, Poland: 3, Portugal: 2, Turkey: 1, Ukraine: 12; Total **30**
subtilis Weise, 1897: India: 10; Total **10**
successiva Spaeth, 1924: Dem. Rep. of Congo: 2, Kenya: 2, Rwanda: 7, Tanzania: 4, Uganda: 1; Total **16**
sulphurago Boheman, 1854: Namibia: 5; Total **5**
taediosa Boheman, 1856: Madagascar: 2; Total **2**
tarda Weise, 1899: Tanzania: 1, Uganda: 2; Total **3**
tenasserimensis (Spaeth, 1926): Laos: 2, Thailand: 1; Total **3**
tenax Spaeth, 1915: Madagascar: 3; Total **3**
thailandica Borowiec, 2001 k: 558: Laos: 2, Thailand: 4 (**1 HT, 2 PT**); Total **6**
thomsoni Boheman, 1862: Former Rhodesia: 1, Namibia: 4, RSA: 2, Zambia: 2 (**1 ST** of *Cassida pentheri* Spaeth, 1905: 109); Total **9**
tianshanica Borowiec & Świętojańska, 2001 e: 70: Kyrgyzstan: 7 (**1 HT, 6 PT**); Total **7**
timorensis Borowiec, 1995 c: 95: Indonesia: 7 (**5 PT**); Total **7**
tosta Klug, 1835: Cameroon: 12, Congo Brazzaville: 1, Dem. Rep. of Congo: 16, Gabon: 1, Ivory Coast: 2, Uganda: 5; Total **37**
transcaucasica Borowiec & Świętojańska, 2001 e: 72: Georgia: 1 (**1 HT**), Iran: 1, no country (Caucasus): 2 (**2 PT**); Total **4**
trianguliformis Borowiec & Świętojańska, 2013 h: 196: Madagascar: 1 (**1 HT**); Total **1**
triangulum (Weise, 1897): India: 1, Indonesia: 10, Thailand: 1; Total **12**
troglydites Boheman, 1854: Africa: 1, Dem. Rep. of Congo: 11, Ethiopia: 1, Ghana: 1, Nigeria: 1, Tanzania: 3, Togo: 2, Uganda: 2; Total **22**
trossula (Spaeth, 1916): Madagascar: 12; Total **12**
truncatipennis (Spaeth, 1914): China: 3, Laos: 1, Myanmar: 1, Thailand: 3, Vietnam: 1; Total **9**
tryznai Borowiec & Świętojańska, 2013 h: 200: Madagascar: 1 (**1 PT**); Total **1**
tsaratanana Borowiec, 1998: Madagascar: 1; Total **1**
tsinlinica Chen & Zia, 1964: China: 2; Total **2**
tuberculata Medvedev & Eroshkina, 1988: 133: Vietnam: 5 (**1 PT**); Total **5**
tumidicollis (Chen & Zia, 1961): China: 2, Laos: 1; Total **3**
turcmenica (Weise, 1892): Kazakhstan: 11, Kyrgyzstan: 4, Turkmenistan: 1, no country: 1; Total **17**
ultima Borowiec & Świętojańska, 2013 h: 2004: Madagascar: 1 (**1 HT**); Total **1**
umbonata Borowiec, 1999 j: 445: Madagascar: 1 (**1 HT**); Total **1**
undecimnotata Gebler, 1841: China: 6, Kazakhstan: 3, Kyrgyzstan: 1, Tadjikistan: 2, Turkmenistan: 38, Uzbekistan: 4; Total **54**
unica Świętojańska & Borowiec, 2002 e: 357: Malaysia (Borneo): 3 (**1 HT, 1 PT**); Total **3**
unicatenata Weise, 1910: Madagascar: 2; Total **2**
unimaculata Boheman, 1854: RSA: 12; Total **12**
uniorbis (Chen & Zia, 1961): China: 1, Laos: 8, Thailand: 1; Total **10**
variabilis (Chen & Zia, 1961): China: 1, Laos: 2, Thailand: 2; Total **5**
varians Herbst, 1799: Ceylon: 1, India: 128, Nepal: 2; Total **131**
varicornis (Spaeth, 1912): Indonesia: 2, Malaysia (Borneo): 1; Total **3**
velaris Weise, 1896: Japan: 2, Russia: 5, South Korea: 1; Total **8**
verrucata (Boheman, 1855): Madagascar: 1; Total **1**
versicolor (Boheman, 1855): China: 9, Japan: 5, South Korea: 1, Vietnam: 4; Total **19**
vespertillo Boheman, 1862: RSA: 2; Total **2**

vespertina Boheman, 1862: China: 12, Japan: 3, South Korea: 2; Total **17**
vibex Linnaeus, 1767: Bulgaria: 6, Kyrgyzstan: 1, Poland: 28, Russia: 2, Slovakia: 3; Total **40**
vietnamica Medvedev & Eroshkina, 1988: 138: Laos: 1, Vietnam: 5 (**1 PT**); Total **6**
villiersi Borowiec & Świętojańska, 2021 (in litt.): Burkina Faso: 1 (**1 PT**), Niger: 9 (**1 HT, 8 PT**); Total **10**
virguncula Weise, 1889: China: 8; Total **8**
viridiguttata Chen & Zia, 1964: Thailand: 1; Total **1**
viridipennis Boheman, 1854: RSA: 21, Zimbabwe: 2; Total **23**
viridis Linnaeus, 1758: Bulgaria: 1, Poland: 23, Russia: 4, Serbia: 2, South Korea: 2; Total **32**
vitalisi (Spaeth, 1928): Laos: 2, Thailand: 2, Vietnam: 3; Total **7**
vittata Villers, 1789: Bulgaria: 3, Israel: 1, Italy: 2, Poland: 21, Spain: 8; Total **35**
vittata ssp. *atlantica* Escalera, 1914: Morocco: 11, Tunisia: 2; Total **13**
wanati Borowiec & Świętojańska, 2021 (in litt.): RSA: 1 (**1 HT**); Total **1**
weinmanni Chapuis, 1880: Ethiopia: 23; Total **23**
weisei (Jacobson, 1894): Kazakhstan: 1; Total **1**
wittmeri Borowiec & Świętojańska, 2021 (in litt.): RSA: 1 (**1 PT**); Total **1**
Cassia sp. 1: China: 1; Total **1**
Cassia sp. 2: China: 1; Total **1**
Cassia sp. 3: Kazakhstan: 2, Uzbekistan: 1; Total **3**
Cassia sp. (Oriental Region): probably 6-10 undescribed species; Total **53**

Genus *Chelysida* Fairmaire, 1882

obtectata Fairmaire, 1883: Ethiopia: 7, Uganda: 2; Total **9**

Genus *Chiridopsis* Spaeth, 1922

acutangula (Weise, 1905): India: 3; Total **3**
aequinoctialis (Olivier, 1808): Cameroon: 4, Dem. Rep. of Congo: 28, Gabon: 4, Gambia: 1, Ivory Coast: 1, Uganda: 2, South Sudan: 4; Total **44**
andamanica (Dohrn, 1880: 370): India: 3 (**2 ST**); Total **3**
atricollis Borowiec, 2005 g: 385: Madagascar: 1 (**1 HT**); Total **1**
aubei (Boheman, 1855): Angola: 4, Burundi: 2, Cameroon: 12, Central African Rep.: 3, Dem. Rep. of Congo: 70, Ethiopia: 6, Gabon: 1, Ivory Coast: 1 (**1 ST Chiridopsis signaticollis** Spaeth, 1924: 346), Kenya: 5, Liberia: 1, Rwanda: 5, São Tomé and Príncipe: 1, South Sudan: 2, Tanzania: 2, Togo: 1, Uganda: 9, Zambia: 5; Total **130**
bipunctata (Linnaeus, 1767): Bangladesh: 1, Ceylon: 2, India: 43; Total **46**
bistrimaculata (Boheman, 1855): India: 20; Total **20**
boutareli (Spaeth, 1917: 442): Chad: 1, Dem. Rep. of Congo: 4, Republic of Congo: 1 (**1 ST**); Total **6**
bowringii (Boheman, 1855): China: 4, Laos: 1, Vietnam: 9; Total **14**
circe (Weise, 1896): Angola: 1; Total **1**
congoana (Spaeth, 1912): Dem. Rep. of Congo: 3; Total **3**
convexula (Boheman, 1854): RSA: 2, Swaziland: 6; Total **8**
coorta Spaeth, 1926: Laos: 1, Vietnam: 1; Total **2**
defecta Medvedev & Eroshkina, 1988: Vietnam: 5; Total **5**
elongata Spaeth, 1924: Dem. Rep. of Congo: 3; Total **3**
flavipennis (Spaeth, 1902): Dem. Rep. of Congo: 1, Kenya: 1 (**1 ST Chiridopsis rothschildi** Spaeth, 1922: 1004), Rwanda: 7, Uganda: 5; Total **14**
ghatei Borowiec & Świętojańska, 2000 c: 53: India: 5 (**1 HT, 4 PT**); Total **5**
gregaria (Weise, 1905): India: 1; Total **1**
irrorata (Spaeth, 1917): Malawi: 1, Zambia: 11, Zimbabwe: 1; Total **13**
klapperichi Borowiec & Świętojańska, 2017 b: 34: RSA: 1 (**1 HT**); Total **1**
laetifica (Boheman, 1862): Cameroon: 26; Total **26**
leopardina (Boheman, 1855): Madagascar: 22; Total **22**
levis Borowiec, 2005 g: 387: Madagascar: 6 (**1 HT, 5 PT**); Total **6**
maculata Borowiec, 2005 g: 388: Madagascar: 3 (**1 HT, 2 PT**); Total **3**

marginata (Weise, 1901): Ceylon: 6, India: 3; Total **9**
marginepunctata Borowiec, 2005 g: 390: Madagascar: 3 (**1 HT, 2 PT**); Total **3**
melanocephala Spaeth, 1924: RSA: 5; Total **5**
mimica (Weise, 1905): India: 3, Myanmar: 6; Total **9**
nickerli (Spaeth, 1911: 276): Madagascar: 4 (**1 ST**); Total **4**
nigropunctata Borowiec & Ghate, 1999 f: 103: India: 10 (**1 HT, 9 PT**); Total **10**
nigroreticulata Borowiec, 2005 g: 391: Madagascar: 1 (**1 HT**); Total **1**
nigrosepta (Fairmaire, 1891): Ethiopia: 3, Kenya: 3, Senegal: 1, Tanzania: 2, Zambia: 16; Total **25**
novemkalankita (Maulik, 1919): India: 9; Total **9**
observabilis (Spaeth, 1916): Dem. Rep. of Congo: 1, Kenya: 2, Rwanda: 1; Total **4**
octoguttata (Boheman, 1854): Nigeria: 1, Senegal: 4; Total **5**
opposita (Boheman, 1854): Burkina Faso: 2, Ghana: 3, Ivory Coast: 3, Nigeria: 1, Senegal: 9; Total **18**
parellina Spaeth, 1937: Tanzania: 10; Total **10**
picturata Spaeth, 1939: Tanzania: 1; Total **1**
punctata (Weber, 1801): China: 1, Indonesia: 11, Malaysia: 4, Vietnam: 34; Total **50**
quadriguttata (Boheman, 1854): Ghana: 6, Liberia: 1; Total **7**
regionalis Spaeth, 1924: RSA: 2; Total **2**
ruandana Spaeth, 1934: Rwanda: 1; Total **1**
rubromaculata Borowiec, Ranade, Rane & Ghate, 2001 i: India: 2 (**1 HT, 1 PT**); Total **2**
scalaris (Weber, 1801): Inonesia: 14, Malaysia: 11, Thailand: 1; Total **26**
selecta (Weise, 1905): India: 19; Total **19**
septemnotata (Boheman, 1855): India: 10, Myanmar: 1, Vietnam: 5, no country: 1; Total **17**
sexplagiata (Spaeth, 1919): India: 12; Total **12**
spadix (Spaeth, 1917): Dem. Rep. of Congo: 3; Total **3**
suffriani (Boheman, 1854): RSA: 1, Zimbabwe: 3; Total **4**
tanganikana (Spaeth, 1917): Dem. Rep. of Congo: 1, Tanzania: 1; Total **2**
tanzaniensis Borowiec & Świętojańska, 2017 b: 60: Tanzania: 2 (**1 HT, 1 PT**); Total **2**
tessellata (Spaeth, 1917): Dem. Rep. of Congo: 3, Zambia: 2; Total **5**
tredecimnotata (Boheman, 1855): Ceylon: 1; Total **1**
tredecimsignata (Boheman, 1855): Ceylon: 3; Total **3**
trizonata (Fairmaire, 1904): Madagascar: 4; Total **4**
undecimnotata (Boheman, 1855): India: 5; Total **5**
ventralis (Boheman, 1855): India: 24; Total **24**
westringi (Boheman, 1862): Philippines: 6; Total **6**
zambiana Borowiec & Świętojańska, 2017 b: 64: Zambia: 5 (**1 HT, 4 PT**); Total **5**
Chiridopsis sp. nov. sp. 1: Madagascar: 1; Total **1**

Genus *Chiridula* Weise, 1889

semenovi Weise, 1899: Afghanistan: 1, Kazakhstan: 1; Total **2**

Genus *Emdenia* Spaeth, 1915

maxima (Blackburn, 1896): Australia: 2; Total **2**

Genus *Erbolaspis* Spaeth, 1924

passaria (Weise, 1900 b: 215): Ethiopia: 1, Kenya: 2, Tanzania: 13 (**1 ST**); Total **16**

Genus *Fornicocassis* Spaeth, 1917

obdurans (Spaeth, 1917): Dem. Rep. of Congo: 2; Total **2**

rufocincta Spaeth, 1917: RSA: 3; Total **3**

unicolor Spaeth, 1934: Ethiopia: 1, Tanzania: 1; Total **2**

Genus *Glyphocassis* Spaeth, 1914

Subgenus *s. str.*

szechuana Gressitt, 1952: China: 2; Total **2**

trilineata (Hope, 1831): India: 11, Vietnam: 6; Total **17**

Subgenus *Hebdomecosta*

lepida (Spaeth, 1914): China: 7; Total **7**

spilota (Gorham, 1885): China: 7; Total **7**

Genus *Hovacassis* Spaeth, 1952

brunneofasciata Borowiec, 2002 i: 98: Madagascar: 2 (**1 HT, 1 PT**); Total **2**

discolor (Boheman, 1855): Madagascar: 16; Total **16**

flavonigra Borowiec, 2002 i: 104: Madagascar: 2 (**2 PT**); Total **2**

murzini Borowiec, 2002 i: 109: Madagascar: 1 (**1 HT**); Total **1**

pulchra (Spaeth, 1915): Madagascar: 2; Total **2**

rubromaculata Borowiec, 2002 i: 114: Madagascar: 1 (**1 HT**); Total **1**

rubrovittata Borowiec, 2002 i: 117: Madagascar: 2 (**1 HT, 1 PT**); Total **2**

Genus *Hypocassida* Weise, 1893

convexipennis Borowiec, 2000 i: 601: Iran: 1 (**1 HT**); Total **1**

cornea (Marseul, 1868): Armenia: 1, Bulgaria: 7, Italia: 1, UAE: 1; Total **10**

grossepunctata Bordy, 2009: 52: Bulgaria: 3, France: 2 (**2 PT**), Spain: 1; Total **6**

meridionalis (Suffrian, 1844): France: 14, Italy: 1, Spain: 2; Total **17**

subferruginea (Schrank, 1776): Bulgaria: 2, Czech Rep.: 5, Egypt: 1, Hungary: 1, Italy: 1, Kazakhstan: 7, Kyrgyzstan: 2, Poland: 9, Russia: 10, Slovakia: 18, Spain: 2, Ukraine: 11; Total **69**

Genus *Ischiocassis* Spaeth, 1917

convexa (Boheman, 1854): Dem. Rep. of Congo: 1, Namibia: 3, RSA: 18, Zambia: 5; Total **27**

stabilis (Weise, 1899): Namibia: 10, South Sudan: 1, Zambia: 2; Total **13**

tragardhi (Spaeth, 1928): RSA: 1; Total **1**

umbrata (Boheman, 1854): RSA: 3; Total **3**

Genus *Ischyronota* Weise, 1891

brisouti (Reitter, 1889): Algeria: 3; Total **3**

conicicollis (Weise, 1890): China: 1, Mongolia: 1, Turkmenistan: 1; Total **3**

desertorum (Gebler, 1834): Armenia: 4, Azerbaijan: 5, Georgia: 2, Kazakhstan: 5, Kyrgyzstan: 7, Pakistan: 1, Russia: 3, Turkey: 1, Turkmenistan: 1, no country: 3; Total **32**

elevata (Weise, 1890): Azerbaijan: 4, Iran: 2, Kazakhstan: 2, Russia: 4, Turkmenistan: 1; Total **13**

jordanensis Borowiec, 1986 b: 575: Jordan: 1 (**1 PT**); Total **1**

schusteri Spaeth, 1914: Mongolia: 1; Total **1**

spaethi Reitter, 1901: Kazakhstan: 2, Russia: 3; Total **5**

Genus *Limnocassis* Spaeth, 1952

compacta Borowiec & Sekerka, 2015 c: 276: RSA: 1 (**1 PT**); Total **1**

pumilio (Boheman, 1854): RSA: 5; Total **5**

regularis Borowiec & Sekerka, 2015 c: 273: RSA: 1 (**1 PT**); Total **1**

Genus *Lorentzocassis* Spaeth, 1913

aruensis Spaeth, 1926: Indonesia: 2; Total **2**

papuana Spaeth, 1913: Indonesia: 1, Papua New Guinea: 1; Total **2**

riedeli Borowiec, 2003 g: 235: Papua New Guinea: 1 (**1 PT**); Total **1**

Genus *Macromonycha* Spaeth, 1911

apicalis (Gebler, 1845): Iran: 1, Russia: 7, Uzbekistan: 1; Total **9**

Genus *Malayocassis* Spaeth, 1952

hilaris (Boheman, 1855): Indonesia: 6, Malaysia (Borneo): 2; Total **8**

manilensis (Weise, 1910): Philippines: 8; Total **8**

Genus *Meroscalsis* Spaeth, 1903

blackburni (Spaeth, 1913): Australia: 3; Total **3**

radiata (Boheman, 1855): Indonesia: 2, Papua New Guinea: 2; Total **4**

selecta Spaeth, 1903: Papua New Guinea: 3; Total **3**

weisei (Spaeth, 1903): Australia: 4, Papua New Guinea: 1; Total **5**

Meroscalsis sp. nov. sp. 1: Papua New Guinea: 1; Total **1**

Meroscalsis sp. nov. sp. 2: Papua New Guinea: 1; Total **1**

Meroscalsis sp. nov. sp. 3: Papua New Guinea: 5; Total **5**

Meroscalsis sp. nov. sp. 4: Papua New Guinea: 1; Total **1**

Genus *Nabathaea* Spaeth, 1911 (1)

pygmaea Spaeth, 1911: Egypt: 5, Yemen: 2; Total **7**

Genus *Oocassida* Weise, 1897

ceylonica Weise, 1901: Ceylon: 1; Total **1**

cruenta (Fabricius, 1792): Bangladesh: 2, India: 15; Total **17**

pudibunda (Boheman, 1856): India: 6, Nepal: 1; Total **7**

schultzei Spaeth, 1917: Burkina Faso: 1, Niger: 2, Senegal: 19; Total **22**

tunisienis (Boheman, 1854): Algeria: 5, no country: 1; Total **6**

Genus *Orobiocassis*, Spaeth, 1934

gabonica (Spaeth, 1933): RSA: 2; Total **2**

testudinea (Boheman, 1854): Namibia: 1, RSA: 5; Total **6**

Genus *Oxylepus* Desbrochers, 1884

bituberculatus Borowiec, 2002 i: 162: RSA: 4 (**2 PT**); Total **4**

boroveci Borowiec, 2001 g: 350: Tunisia: 4 (**1 HT, 3 PT**); Total **4**

capensis Spaeth, 1933: RSA: 7; Total **7**

cicatricosa Spaeth, 1936 b: 10: Namibia: 1 (**1 ST**); Total **1**

convexicollis Borowiec, 2002 i: 168: RSA: 7 (**7 PT**); Total **7**

cuneipennis Spaeth, 1936: RSA: 2; Total **2**

deflexicollis (Boheman, 1862): Algeria: 5, Croatia: 1, France: 3, Greece: 4, Morocco: 1, Spain: 4; Total **18**

grobbellaarae Borowiec, 2002 i: 173: Namibia: 2 (**2 PT**), RSA: 6 (**6 PT**); Total **8**

impressipennis Borowiec, 2002 i: 176: Namibia: 8 (**8 PT**); Total **8**

intermedius Borowiec, 2002 i: 178: Namibia: 2 (**1 HT, 1 PT**); Total **2**

kossmati Spaeth, 1901: Djibuti: 1, Erytrea: 1, Oman: 1, Tanzania: 1, Yemen: 4, no country: 1; Total **9**

planicollis Borowiec, 2002 i: 184: RSA: 3 (**3 PT**); Total **3**

sextuberculatus Borowiec, 2002 i: 186: RSA: 20 (**20 PT**); Total **20**

Genus *Pilemostoma* Desbrochers, 1891

fastuosa (Schaller, 1783): Bulgaria: 7, Poland: 6, Turkey: 2, Ukraine: 4; Total **19**

Genus *Psalidoma* Spaeth, 1899

holubi Spaeth, 1899: Botswana: 8; Total **8**

knirshi Spaeth, 1919: Malawi: 4, Mozambique: 2, Zimbabwe: 1; Total **7**

Genus *Rhacocassis* Spaeth, 1904

balyi (Boheman, 1855): Ceylon: 2; Total **2**

flavoplagiata (Baly, 1863): Indonesia: 1, Malaysia: 8; Total **9**

Genus *Rhytidocassis* Spaeth, 1941

angulipennis Borowiec, 2002 i: 201: Namibia: 1 (**1 HT**); Total **1**

indicola (Duvivier, 1892): Bangladesh: 1, India: 1; Total **2**

iranella (Lopatin, 1984): Iran: 1; Total **1**

limbiventris (Boheman, 1854): Dem. Rep. of Congo: 2, Ethiopia: 4, Namibia: 2, Tanzania: 28; Total **36**

lopatini Borowiec & Świętojańska, 2001 b: 325: Iran: 7 (**1 HT**); Total **7**

minuta (Boheman, 1854): RSA: 1; Total **1**

scutellaris (Klug, 1835): Angola: 1, Ghana: 1, Niger: 1, Senegal: 8, Sudan: 5; Total **16**

Genus *Seminabathea* Borowiec, 1994

arabica (Spaeth, 1911): Oman: 1; Total **1**

Genus *Silana* Spaeth, 1914

farinosa (Boheman, 1856): Ceylon: 1, Malaysia: 1; Total **2**

Genus *Smeringaspis* Spaeth, 1924

hirsuta Spaeth, 1952: Zambia: 1; Total **1**

setifera (Boheman, 1854): Namibia: 5, RSA: 1, Senegal: 8; Total **14**

Genus *Sphenocassis* Spaeth, 1911

anosibensis Borowiec, 2002 i: 230: Madagascar: 2 (**1 HT, 1 PT**); Total **2**

humerosa (Fairmaire, 1898): Madagascar: 6; Total **6**

imerina (Spaeth, 1926): Madagascar: 2; Total **2**

impressipennis Borowiec, 2002 i: 238: Madagascar: 2 (**1 HT, 1 PT**); Total **2**

incisicollis (Spaeth, 1911): Madagascar: 2; Total **2**

punctatissima (Weise, 1910): Madagascar: 4; Total **4**

Genus *Tegocassis* Spaeth, 1924 (1)

corpulenta (Weise, 1904): Cameroon: 3, Madagascar: 2; Total **5**

Genus *Thlaspida* Weise, 1899

biramosa (Boheman, 1855): China: 10, Japan: 5, Malaysia: 3, Taiwan: 3, Vietnam: 53; Total **74**

cribrosa (Boheman, 1854): India: 1, Laos: 1, Myanmar: 4, Vietnam: 1; Total **7**

lewisii (Baly, 1874): China: 1, North Korea: 5, Russia: 2; Total **8**

tsoui Borowiec & Lee, 2009 e: 350: Taiwan: 2 (**2 PT**); Total **2**

Genus *Thlaspidosoma* Spaeth, 1901

brevis Chen & Zia, 1964: Laos: 2, Vietnam: 1; Total **3**

celebensis Spaeth, 1933: Malaysia: 1; Total **1**

dohrni Spaeth, 1901: Indonesia: 3, Malaysia (Borneo): 2; Total **5**

fallaciosa Spaeth, 1901: Indonesia: 6; Total **6**

gracilicornis Spaeth, 1915: Malaysia (Borneo): 3; Total **3**

horsfieldi (Boheman, 1855): Indonesia: 3; Total **3**

philippina (Spaeth, 1916): Philippines: 4; Total **4**

rizalensis Spaeth, 1933: Philippines: 3; Total **3**

Genus *Thlaspidula* Spaeth, 1901

fimbriata Spaeth, 1901: Malaysia (Borneo): 1; Total **1**

moseri (Spaeth, 1915): Indonesia: 1; Total **1**
muelleri (Spaeth, 1903): Papua New Guinea: 5; Total **5**
riedeli Borowiec & Świętojańska, 2001 c: 335: Indonesia: 2 (**1 HT**); Total **2**

Genus *Trichaspis* Spaeth, 1911

brevicornis Borowiec, 2002 i: 257: Namibia: 1 (**1 HT**); Total **1**
erinacea Borowiec, 2002 i: 261: Democratic Rep. of Congo: 4 (**1 HT, 4 PT**); Total **4**
louwi Borowiec, 2002 i: 265: RSA: 7 (**7 PT**); Total **7**
minutissima Borowiec, 2002 i: 269: Zambia: 1 (**1 HT**); Total **1**
pilosa (Spaeth, 1911): RSA: 6; Total **6**
pilosula (Boheman, 1862): Namibia: 7; Total **7**
tomentosa Borowiec, 2002 i: 276: RSA: 5 (**5 PT**); Total **5**

Genus *Trigonocassis* Hincks, 1950

conducta (Boheman, 1862): Cameroon: 3, Democratic Rep. of Congo: 3, Togo: 2, Uganda: 2; Total **10**

Genus *Vietocassis* Medvedev & Eroshkina, 1988

viridis Medvedev & Eroshkina, 1988: 129: Vietnam: 4 (**1 PT**); Total **4**

New World genera

Genus *Agroiconota* Spaeth, 1913

atromaculata Borowiec, 2005 a: 62: Peru: 9 (**1 HT, 5 PT**); Total **9**
atropunctata Borowiec, 2005 a: 62: Peru: 2 (**2 PT**); Total **2**
aulatipennis Spaeth, 1936: Argentina: 2; Total **2**
bivittata (Say, 1827): USA: 12; Total **12**
carlobrivioi Borowiec, 2005 a: 64: Brazil: 2 (**2 PT**), Peru: 7 (**3 PT**); Total **9**
conflagrata (Boheman, 1855): Brazil: 1, French Guyana: 13, Peru: 1, Trinidad and Tobago: 2; Total **17**
cubana Blake, 1970: Cuba: 1; Total **1**
gibberosa (Boheman, 1855): Brazil: 2; Total **2**
gibbipennis Borowiec, 2005 a: 66: Brazil: 8 (**1 HT, 7 PT**); Total **8**
inedita (Boheman, 1855): Argentina: 1, Brazil: 15, French Guyana (?mislabelled): 1, Paraguay: 1, no country: 1;
Total **19**
judaica (Fabricius, 1781): Bolivia: 1, Brazil: 5, Colombia: 5, Costa Rica: 3, Ecuador: 6, French Guyana: 8, Panama:
14, Peru: 26, Guyana: 3, Surinam: 3, Trinidad and Tobago: 2; Total **76**
lateripunctata Spaeth, 1936: Bolivia: 1, Venezuela: 4; Total **5**
paraguayana Borowiec, 2005 a: 65: Paraguay: 2 (**1 HT, 1 PT**); Total **2**
parellina Spaeth, 1937: Argentina: 1, Uruguay: 1; Total **2**
propinqua (Boheman, 1855): Colombia: 8, Costa Rica: 5, Dominican Rep.: 4, El Salvador: 1, Jamaica: 1, Panama:
16; Total **35**
pullula (Boheman, 1855): Bolivia: 1, Brazil: 9, Peru: 1; Total **11**
punctipennis (Boheman, 1855): Brazil: 2; Total **2**
reimoseri Spaeth, 1936: Paraguay: 2; Total **2**
sanareensis Borowiec, 2005 a: 69: Venezuela: 6 (**6 PT**); Total **6**
sodalis Spaeth, 1936: Colombia: 1, Venezuela: 4, no country: 1; Total **6**
stupidula (Boheman, 1855): Bolivia: 6, Brazil: 6, Paraguay: 2, Venezuela: 2; Total **16**
subtriangularis Spaeth, 1936: Bolivia: 1, Peru: 7; Total **8**
subvittata (Boheman, 1855): Brazil: 1; Total **1**
tristriata (Fabricius, 1792): Argentina: 3, Brazil: 4, French Guyana: 3, Paraguay: 1, Peru: 1; Total **12**
vilis (Boheman, 1855): Costa Rica: 1, Guatemala: 2, Mexico: 5; Total **8**

Genus *Aidoia* Spaeth, 1952

nubilosa (Boheman, 1855): Colombia: 3, Ecuador: 1; Total **4**

Genus *Aporocassida* Spaeth, 1952

aemula (Boheman, 1854): Bolivia: 1, Brazil: 1; Total **2**

graphica (Germar, 1824): Argentina: 19, Brazil: 2, no country: 4; Total **25**

Genus *Bradycassis* Spaeth, 1952

agroiconotoides Borowiec, 2005 c: 44: Argentina: 10 (**8 PT**), Brazil: 11 (**11 PT**), Paraguay: 7 (**7 PT**); Total **28**

drewseni (Boheman, 1855): Argentina: 4, Bolivia: 3, Uruguay: 2; Total **9**

immersa (Spaeth, 1909): Argentina: 1, Paraguay: 1; Total **2**

matogrossoensis Świętojańska & Borowiec, 1996 f: 471: Brazil: 4 (**4 PT**); Total **4**

romani (Spaeth, 1931): Brazil: 6; Total **6**

sandaricina (Boheman, 1862): Brazil: 3; Total **3**

sordescens (Spaeth, 1926): Brazil: 5; Total **5**

Genus *Chacocassis* Spaeth, 1952

fuscovittata (Boheman, 1855): Argentina: 4; Total **4**

Genus *Charidotella* Weise, 1896**Subgenus *Charidotella* s. str.**

actiosa (Spaeth, 1926): Colombia: 1, Ecuador: 2; Total **3**

balteata (Champion, 1894): Costa Rica: 1; Total **1**

bifasciata (Linnaeus, 1758): Bolivia: 3, Brazil: 2, French Guyana: 7, Peru: 4, Surinam: 1; Total **17**

bifossulata (Boheman, 1855): Mexico: 11; Total **11**

bifoveata (Spaeth, 1926): Bolivia: 2, Brazil: 7, Paraguay: 3; Total **12**

bordoni Borowiec, 2002 a: 126: Venezuela: 3 (**2 PT**); Total **3**

canchamayana (Spaeth, 1926): Ecuador: 1; Total **1**

carnulenta (Erichson, 1847): Peru: 7, no country: 1; Total **8**

circumnotata (Boheman, 1862): Brazil: 2, Colombia: 3, Ecuador: 1, French Guyana: 2; Total **8**

conclusa (Boheman, 1855): Brazil: 2; Total **2**

discoidalis (Boheman, 1855): Brazil: 1; Total **1**

dominicanensis Borowiec, 2011 d: 496: Dominican Rep.: 2 (**1 HT**); Total **2**

duplex (Champion, 1894): Guatemala: 1; Total **1**

ecuadorica Borowiec, 1989 b: 206: Ecuador: 1 (**1 HT**); Total **1**

egregia (Boheman, 1855): Costa Rica: 1, Mexico: 1, Panama: 1; Total **3**

ferranti (Spaeth, 1926): Argentina: 1, Bolivia: 4; Total **5**

fumosa (Boheman, 1855): Bolivia: 1, French Guyana: 4, Paru: 3; Total **8**

glaucovittata (Erichson, 1847): Bolivia: 4, Ecuador: 11, Peru: 7; Total **22**

quadeloupensis (Boheman, 1855): Guadeloupe: 2; Total **2**

hoegbergi (Boheman, 1855): Argentina: 3, Brazil: 4, Colombia: 2, El Salvador: 2, French Guyana: 2, Honduras: 2, Mexico: 2, Panama: 1, Paraguay: 2; Total **20**

immaculata (Olivier, 1790): Bolivia: 3, Brazil: 25, Colombia: 2, French Guyana: 4, Paraguay: 1, Paru: 1, Venezuela: 1; Total **37**

incerta (Boheman, 1855): Brazil: 9; Total **9**

inconstans (Boheman, 1855): Brazil: 2, French Guyana: 13; Total **15**

incorrupta (Boheman, 1855): Brazil: 2, Colombia: 5, Panama: 1, Peru: 27, Venezuela: 1; Total **36**

jamaicensis (Blake, 1966): Jamaica: 2; Total **2**

kesseli Borowiec, 1989 b: 212: Brazil: 4 (**1 HT, 3 PT**); Total **4**

latevittata (Boheman, 1855): Dominican Rep.: 1; Total **1**

limpida (Boheman, 1855): Argentina: 1, Bolivia: 2, Brazil: 8, Paraguay: 14; Total **25**

liquida (Erichson, 1847): Bolivia: 1, Brazil: 4, Guyana: 1, Trinidad and Tobago: 1; Total **7**
moraguesi Borowiec, 2007 e: 60: French Guyana: 6 (**1 HT, 5 PT**); Total **6**
morio (Fabricius, 1801): Brazil: 17; Total **17**
myops (Boheman, 1855): Ecuador: 2, Peru: 1, no country: 2; Total **5**
nigripennis Borowiec, 2009 d: 344: Venezuela: 1 (**1 HT**); Total **1**
oblita (Suffrian, 1868): Cuba: 2; Total **2**
pacata Borowiec, 2007 e: 62: Bolivia: 3 (**1 HT, 2 PT**), Brazil: 2 (**2 PT**); Total **5**
pallescens (Boheman, 1855): Argentina: 2, Bolivia: 1, Brazil: 2, Paraguay: 1; Total **6**
pectoralis (Kirsch, 1883): Bolivia: 8, Brazil: 3, Paraguay: 2; Total **13**
pellucida (Boheman, 1855): Brazil: 1; Total **1**
posticata (Boheman, 1855): Brazil: 1; Total **1**
praeusta (Boheman, 1855): Dominican Rep.: 1, Haiti: 1; Total **2**
proxima (Boheman, 1855): Mexico: 5; Total **5**
pudica (Boheman, 1855): Brazil: 1, Paraguay: 2, no country: 1; Total **4**
purpurea (Linnaeus, 1758): Bolivia: 3, Brazil: 3, French Guyana: 1, Peru: 7; Total **14**
quadrisignata (Boheman, 1855): Cuba: 5; Total **5**
rasilis (Spaeth, 1926 c: 23): Peru: 1; Total **1**
recidiva (Spaeth, 1926: 23): Argentina: 2, Brazil: 11 (**1 ST**); Total **13**
semiatrata (Boheman, 1862): Guatemala: 1, Mexico: 1; Total **2**
sexpunctata (Fabricius, 1781): Argentina: 2, Brazil: 12, Canada: 2, Colombia: 16, Costa Rica: 1, El Salvador: 2,
 French Guyana: 12, Guadeloupe: 2, Guatemala: 2, Honduras: 4, Mexico: 4, Panama: 1, Paraguay: 7, Peru: 3,
 Saint Lucia: 1, USA: 13, Venezuela: 5, no country: 1; Total **90**
sinuata (Champion, 1894): Costa Rica: 5, Panama: 4; Total **9**
steinhauseni Borowiec, 1989 b: 209: Jamaica: 4 (**4 PT**); Total **4**
striatopunctata (Boheman, 1855): Brazil: 1; Total **1**
stulta (Boheman, 1855): Brazil: 1; Total **1**
subannulata (Boheman, 1862): Costa Rica: 1; Total **1**
submaculata (Boheman, 1855): Surinam: 1; Total **1**
succinea (Boheman, 1855): El Salvador: 1, Guatemala: 1, Honduras: 1, Mexico: 18, USA: 4; Total **25**
tuberculata (Fabricius, 1775): Costa Rica: 2, El Salvador: 1, Guatemala: 6, Mexico: 4, Panam: 1; Total **14**
tumida (Champion, 1894): Costa Rica: 1; Total **1**
ventricosa (Boheman, 1855): Guatemala: 1, Mexico: 5; Total **6**
vinula (Boheman, 1855): Bolivia: 1, Brazil: 2, Colombia: 2, French Guyana: 23, Honduras: 1, Mexico: 1, Venezuela:
 3; Total **33**
virgo (Boheman, 1855): Argentina: 1, Brazil: 5, Ecuador: 1; Total **7**
virgulata (Boheman, 1855): El Salvador: 3, Guatemala: 2, Mexico: 4; Total **9**
zona (Fabricius, 1801): Bolivia: 2, Costa Rica: 3, French Guyana: 34: Panama: 1, Peru: 1; Total **41**
Charidotella s. str.: undetermined: 16; Total **16**
Subgenus *Chaerocassis* (6):
annexa (Boheman, 1855): Honduras: 1, Mexico: 4; Total **5**
atromarginata Borowiec, 2009 d: 342: Mexico: 3 (**1 HT, 1 PT**); Total **3**
emarginata (Boheman, 1855): Costa Rica: 1, Guatemala: 6, Mexico: 2, USA: 1, no country: 2; Total **12**
marculenta (Boheman, 1855): El Salvador: 3, Guatemala: 1, Nicaragua: 1; Total **5**
ormondensis (Blatchley, 1920): USA: 2; Total **2**
purpurata (Boheman, 1855): Canada: 1, USA: 12; Total **13**
Subgenus *Chapadacassis* (1):
paradoxa Sekerka & Borowiec, 2015 b: 71: Brazil: 1 (**1 HT**); Total **1**
Subgenus *Metrionaspis* (2):
rubicunda (Guérin, 1844): Brazil: 12; Total **12**
santaremi Borowiec, 1995 g: 441: Brazil: 5 (**4 PT**); Total **5**
Subgenus *Philaspis* (6):
bistillata (Spaeth, 1936): Bolivia: 2; Total **2**

bivulnerata (Boheman, 1855): Mexico: 6; Total **6**
goyazensis (Spaeth, 1936): Brazil: 2; Total **2**
marginepunctata Borowiec, 2004 d: 253: Brazil: 2 (**1 HT, 1 PT**); Total **2**
nigriceps (Spaeth, 1936): Paraguay: 1; Total **1**
seriatopunctata Spaeth, 1900): Paraguay: 6; Total **6**
subnotata (Boheman, 1855): Argentina: 8, Paraguay: 2; Total **10**
Subgenus *Xenocassis* (8):
ambita (Champion, 1894): Costa Rica: 4, Panama: 2; Total **6**
amicula (Spaeth, 1936): Colombia: 2; Total **2**
amoena (Boheman, 1855): Costa Rica: 5, El Salvador: 1, Mexico: 3 (**1 PLT *Coptocyclus ephippiata* Boheman, 1855: 214**); Total **9**
atalanta (Boheman, 1862): Guatemala: 1; Total **1**
opulenta (Boheman, 1855): Guatemala: 4, Mexico: 3; Total **7**
puella (Boheman, 1855): Colombia: 16, Costa Rica: 7, Mexico: 2, Venezuela: 1, no country: 2; Total **28**
rubidoannula Sekerka in litt.: Panama: 1 (**1 PT**); Total **1**
Charidotella (Xenocassis) sp.: 3; Total **3**

Genus *Charidotis* Boheman, 1854

abrupta Boheman, 1855: Brazil: 2; Total **2**
admirabilis Boheman, 1855: Brazil: 1; Total **1**
aerosa Spaeth, 1936: 76: Brazil: 2 (**1 ST**); Total **2**
ambigua Boheman, 1855: Brazil: 5, French Guyana: 3; Total **8**
annularis (Boheman, 1855): Brazil: 6; Total **6**
atramentosa Boheman, 1855: Brazil: 3; Total **3**
atrocruentata Sekerka & Windsor in litt.: Panama: 1 (**1 PT**); Total **1**
aurofasciata (Erichson, 1847): Bolivia: 2, Costa Rica: 1, French Guyana: 2, Panama: 1, Peru: 6, Venezuela: 1; Total **13**
auroguttata Boheman, 1855: Argentina: 1, Brazil: 3, Costa Rica: 1, Mexico: 3, Paraguay: 2, Venezuela: 4; Total **14**
aurulenta (Fabricius, 1801): Brazil: 6, French Guyana: 1; Total **7**
beatula Spaeth, 1936: Brazil: 3; Total **3**
bellula Boheman, 1855: Brazil: 2; Total **2**
biarcuata (Boheman, 1855): Brazil: 4; Total **4**
bicallosula Spaeth, 1936: Brazil: 1; Total **1**
bicincta Boheman, 1855: Brazil: 1; Total **1**
bicingulata (Boheman, 1855): Bolivia: 1, Brazil: 3; Total **4**
bimarginata Boheman, 1855: Bolivia: 1; Total **1**
bipartita (Boheman, 1855): Bolivia: 1, Honduras: 1, no country: 1; Total **3**
brevicollis Spaeth, 1936: Brazil: 1; Total **1**
candens (Boheman, 1855): Brazil: 4; Total **4**
carbunculus Spaeth, 1926: Argentina: 1; Total **1**
centromaculata Boheman, 1855: Brazil: 1; Total **1**
chacoana Spaeth, 1936: Brazil: 2; Total **2**
chermesina Spaeth, 1936: Brazil: 1; Total **1**
cinctella Boheman, 1855: Brazil: 2; Total **2**
cincticulus (Boheman, 1855): Bolivia: 1, Brazil: 3, French Guyana: 1, Peru: 1; Total **6**
circinata (Boheman, 1855): Brazil: 1; Total **1**
circulifera Boheman, 1862: Brazil: 1; Total **1**
circulus (Boheman, 1855): Brazil: 1; Total **1**
circumducta (Boheman, 1855): Brazil: 10; Total **10**
circumscripata Boheman, 1855: Brazil: 12; Total **12**
circumtexta Boheman, 1862: Brazil: 1; Total **1**
clitelligera (Boheman, 1855): Brazil: 1; Total **1**

clypeolata Boheman, 1855: Brazil: 1, Paraguay: 2, no country: 1; Total **4**
coccinea Boheman, 1855: Ecuador: 1; Total **1**
cognata Boheman, 1855: Brazil: 2, French Guyana: 2; Total **4**
concentrica (Boheman, 1855): Brazil: 1, no country: 1; Total **2**
connexa (Boheman, 1855): Brazil: 4; Total **4**
consentanea (Boheman, 1855): Argentina: 2, Brazil: 9, no country: 1; Total **12**
consimilis (Boheman, 1855): Brazil: 1, Surinam: 1; Total **2**
contexta (Boheman, 1855): Brazil: 1; Total **1**
costaricea Spaeth, 1936: Costa Rica: 4; Total **4**
crenata (Boheman, 1855): Brazil: 2; Total **2**
cyclops (Fabricius, 1801): Surinam: 1, Venezuela: 1; Total **2**
diabolica Świetojańska & Borowiec, 2000 j: 610: Brazil: 1 (**1 HT**); Total **1**
divisa Weise, 1904: Bolivia: 2; Total **2**
drakei Weise, 1904: Argentina: 1, Brazil: 3, Paraguay: 7; Total **11**
erythraea Boheman, 1862: Brazil: 1, Peru: 1; Total **2**
erythro stigma Champion, 1894: Mexico: 2; Total **2**
exigua Boheman, 1855: Colombia: 1, Venezuela: 1; Total **2**
exquisita (Boheman, 1862): Brazil: 2, Mexico: 1 (mislabeled), Paraguay: 5; Total **8**
flavicans Boheman, 1855: Brazil: 3; Total **3**
formalis Spaeth, 1936: Colombia: 1; Total **1**
formosa (Boheman, 1855): Brazil: 2, Paraguay: 2; Total **4**
fugitiva Spaeth, 1936: Paraguay: 1; Total **1**
furunculus (Boheman, 1855): Brazil: 5, Paraguay: 2; Total **7**
furva Boheman, 1855: Brazil: 6; Total **6**
gemellata Boheman, 1855: Brazil: 8, Paraguay: 1; Total **9**
gibbipennis Spaeth, 1910: Brazil: 1, Paraguay: 1; Total **2**
glomerosa Boheman, 1855: Brazil: 2; Total **2**
incincta (Boheman, 1862): Brazil: 2, Colombia: 2, Panama: 10, Peru: 2; Total **16**
iris (Perty, 1830): Brazil: 1; Total **1**
jansonorum Sekerka & Wilson in litt.: Panama: 1 (**1 PT**); Total **1**
leprieuri (Boheman, 1855): Belize: 1, Brazil: 2, Costa Rica: 1, Ecuador: 3, Surinam: 1; Total **8**
luteola Boheman, 1855: Brazil: 6, Peru: 3, no country: 1; Total **10**
mansueta (Boheman, 1855): Brazil: 3, Paraguay: 6; Total **9**
marginalis Boheman, 1855: Brazil: 2; Total **2**
margarella (Fabricius, 1775): Brazil: 7; Total **7**
miniata Boheman, 1855: Brazil: 1; Total **1**
miniatula Boheman, 1862: Brazil: 4, Paraguay: 1; Total **5**
mrazi Spaeth, 1936: Brazil: 1; Total **1**
neglecta (Boheman, 1855): Brazil: 4; Total **4**
nevermanni Spaeth, 1936: Costa Rica: 2, Panama: 1; Total **3**
nigrocincta Boheman, 1855: Brazil: 3; Total **3**
nucleata Boheman, 1855: Brazil: 5; Total **5**
oblectans Spaeth, 1936: Brazil: 1; Total **1**
ocularis Boheman, 1855: Brazil: 3; Total **3**
odegardi Sekerka & Windsor in litt.: Panama: 1 (**1 PT**); Total **1**
pastica Spaeth, 1936: French Guyana: 2; Total **2**
petulans Spaeth, 1936: Brazil: 8; Total **8**
plicatula Boheman, 1855: Brazil: 9; Total **9**
porosula Spaeth, 1902: Bolivia: 2, Brazil: 5, Ecuador: 2, Peru: 1; Total **10**
procincta (Boheman, 1855): Brazil: 7; Total **7**
pulchra Spaeth, 1902: Bolivia: 1; Total **1**
punctostriata Boheman, 1856: Brazil: 9, Paraguay: 9, Peru: 3; Total **21**

pupillata (Boheman, 1855): Brazil: 7; Total **7**
pustulata Champion, 1894: Belize: 1; Total **1**
pygmaea (Klug, 1829): Brazil: 4, Paraguay: 1; Total **5**
quadrinaculata Kirsch, 1876: Brazil: 2; Total **2**
quadrioculata (Boheman, 1855): Paraguay: 5; Total **5**
reinecki Spaeth, 1936: Bolivia: 2; Total **2**
repanda (Boheman, 1855): Bolivia: 4, Brazil: 1; Total **5**
rotundata Boheman, 1855: Brazil: 1; Total **1**
rubrodiscoidalis (Boheman, 1855): Argentina: 1, Brazil: 3, Paraguay: 2; Total **6**
rubropicta (Boheman, 1855): Brazil: 1; Total **1**
rubrovittata (Boheman, 1855): Brazil: 1; Total **1**
scarlatina Spaeth, 1936: Ecuador: 1, Peru: 1; Total **2**
soror Boheman, 1855: Brazil: 1; Total **1**
speculum (Boheman, 1855): Brazil: 1; Total **1**
spretata Boheman, 1855: Brazil: 1; Total **1**
stockwelli Sekerka & Windsor in litt.: Panama: 1 (**1 PT**); Total **1**
strandi Spaeth, 1936: Brazil: 1; Total **1**
subrugosa (Boheman, 1855): Brazil: 2; Total **2**
surda (Boheman, 1855): Brazil: 1; Total **1**
tantilla Boheman, 1855: Brazil: 4; Total **4**
tessellata Boheman, 1855: Brazil: 3; Total **3**
tricolor Boheman, 1855: Guyana: 1; Total **1**
venusta Spaeth, 1936: Peru: 1; Total **1**
vitreata (Perty, 1830): Brazil: 3, Panama: 1; Total **4**
Charidotis sp. 1: Brazil: 1; Total **1**
Charidotis sp. 2: Brazil: 1; Total **1**
Charidotis sp. 3: Brazil: 1; Total **1**
Charidotis sp. 4: Ecuador: 1; Total **1**
Charidotis sp. 5: Brazil: 1; Total **1**
Charidotis sp. 6: Paraguay: 1; Total **1**
Charidotis sp. 7: Brazil: 1; Total **1**
Charidotis sp. 8: Argentina: 1; Total **1**

Genus *Chersinellina* Hincks, 1950

heteropunctata (Boheman, 1854): Colombia: 1, Panama: 23; Total **24**

Genus *Coptocyclus* Chevrolat, 1836

Subgenus s. str.

atropunctata Sekerka & Windsor in litt.: Panama: 1 (**1 PT**); Total **1**
auricoma Boheman, 1855: Brazil: 4; Total **4**
aurifera Boheman, 1855: Brazil: 6; Total **6**
bicolor (Germar, 1824): Brazil: 2; Total **2**
concolor Boheman, 1855: Brazil: 3, Colombia: 2, Venezuela: 2; Total **7**
conspicillata Boheman, 1855: Brazil: 3; Total **3**
dolosa Boheman, 1855: Bolivia: 6, Brazil: 7, Peru: 1; Total **14**
infans Spaeth, 1937: Bolivia: 1; Total **1**
laeta Boheman, 1855: Brazil: 5; Total **5**
mundula Boheman: 1862: Brazil: 2; Total **2**
orbiculata Champion, 1894: Panama: 1; Total **1**
placida Boheman, 1855: Bolivia: 1, Peru: 3; Total **4**
roseocincta Boheman, 1855: Brazil: 6; Total **6**
sordida Boheman, 1855: Costa Rica: 1, Mexico: 3, Sweden: 1 (imported with bananas); Total **5**

stigma (Germar, 1824): Brazil: 9; Total **9**
undecimpunctata (Fabricius, 1781): French Guayana: 19; Total **19**
virguncula Boheman, 1862: Brazil: 4; Total **4**
Subgenus *Coptocyclella*
adamantina (Germar, 1824): Argentina: 1, Bolivia: 4, Brazil: 4, Paraguay: 3, Peru: 1, Venezuela: 3; Total **16**
fastidiosa Boheman, 1855: Brazil: 7; Total **7**
Subgenus *Dyscineta*
robusta Spaeth, 1936 a: 261: Ecuador: 2 (**1 PT**), Paru: 2; Total **4**
Subgenus *Podostraba*
arcuata (Swederus, 1787): Brazil: 6; Total **6**
laqueata Spaeth, 1936: Brazil: 3; Total **3**
lunifera Boheman, 1855: Brazil: 2; Total **2**
paranensis Spaeth, 1936: Brazil: 4; Total **4**
ruficornis Spaeth, 1936: Brazil: 2; Total **2**
Subgenus *Psalidonota*
aerata (Boheman, 1855): Brazil: 1; Total **1**
atriceps (Boheman, 1855): Brazil: 2; Total **2**
bisbipustulata Boheman, 1855: Brazil: 1; Total **1**
constellata Boheman, 1855: Bolivia: 1, Brazil: 3, Peru: 1; Total **5**
contemta (Boheman, 1855): Argentina: 7, Bolivia: 2, Brazil: 15, Paraguay: 3, Peru: 2; Total **29**
dorsoplagiata Champion, 1894: Mexico: 2; Total **2**
dorsopunctata (Klug, 1829): Cuba: 11; Total **11**
elegans Boheman, 1855: Brazil: 4; Total **4**
excelsa Boheman, 1855: Brazil: 2; Total **2**
inopinata Sekerka & Windsor in litt.: Brazil: 1 (**1 PT**), Colombia 1 (**1 PT**), Costa Rica: 2 (**2 PT**), Panama: 2 (**2 PT**),
Venezuela: 1 (**1 PT**); Total **7**
jamaicana Spaeth, 1936: Jamaica: 2; Total **2**
leprosa (Boheman, 1855): Belize: 1, Brazil: 1, Costa Rica: 4, El Salvador: 5, Guatemala: 2, no country: 1; Total **14**
marmorata Champion, 1894: Mexico: 4; Total **4**
sagana Boheman, 1862: Ecuador: 1; Total **1**
strandii Spaeth, 1936: Ecuador: 6; Total **6**
texana (Schaeffer, 1933): Mexico: 3, USA: 8; Total **11**
vana Boheman, 1855: Brazil: 3; Total **3**

Genus *Crambelea* Spaeth, 1913

illudens (Boheman, 1854): Brazil: 2; Total **2**
minuta Borowiec, 2009 b: 121: Paraguay: 2 (**1 HT, 1 PT**); Total **2**

Genus *Cteisella* Weise, 1896

amicta (Boheman, 1855): Bolivia: 2, Brazil: 2, Peru: 1; Total **5**
centropunctata (Boheman, 1855): Panama: 1, Venezuela: 1; Total **2**
confusa (Boheman, 1855): Argentina: 16, Brazil: 5, Paraguay: 4, Peru: 1; Total **26**
egens Spaeth, 1910: Paraguay: 5; Total **5**
flava Borowiec, 2004 e: 387: Brazil: 1 (**1 HT**); Total **1**
flavocincta (Boheman, 1855): Brazil: 2; Total **2**
gentilis Spaeth, 1926: Argentina: 2, Brazil: 1; Total **3**
imitatrix Spaeth, 1901: Peru: 7; Total **7**
impura (Boheman, 1855): Brazil: 4; Total **4**
indecorata (Boheman, 1855): Brazil: 1; Total **1**
intricata Spaeth, 1926: Argentina: 1, Brazil: 2, no country: 1; Total **4**
magica (Boheman, 1855): Brazil: 3; Total **3**
ramosa Spaeth, 1926: Argentina: 7, Brazil: 8, Paru: 2; Total **17**

rotalis (Boheman, 1855): Brazil: 1; Total **1**
signatifera (Boheman, 1855): Brazil: 3; Total **3**
virescens (Boheman, 1855): Brazil: 31; Total **31**
zonata (Boheman, 1855): Brazil: 1; Total **1**

Genus *Ctenocassida* Spaeth, 1926

blandidica (Boheman, 1855): Argentina: 1, Brazil: 1; Total **2**
glareosa (Boheman, 1855): Brazil: 5; Total **5**
reimoseri Spaeth, 1926: Paraguay: 3; Total **3**
rustica (Boheman, 1855: 435): Brazil: 1 (**1 ST**); Total **1**
vitellina (Boheman, 1855): Brazil: 1; Total **1**

Genus *Ctenocharidotis* Spaeth, 1926

chlorion (Spaeth, 1926): Brazil: 1; Total **1**
crispata (Boheman, 1855): Brazil: 1; Total **1**
ornatipennis (Spaeth, 1926): Brazil: 1; Total **1**
roseopicta (Boheman, 1855): Brazil: 4; Total **4**
subplagiata (Boheman, 1855): Brazil: 1; Total **1**

Genus *Ctenophilaspis* Spaeth, 1926

ingenua (Boheman, 1855): Brazil: 1; Total **1**

Genus *Cyclocassis* Spaeth, 1913

circulata (Boheman, 1855): French Guyana: 2, Mexico: 1, Panama: 1; Total **4**
secunda Borowiec, 1998 d: 234: Colombia: 1, Ecuador: 1 (**1 HT**); Total **2**

Genus *Deloyala* Chevrolat, 1836

cruciata (Linnaeus, 1758): Brazil: 8, French Guyana: 1; Total **9**
fuliginosa (Olivier, 1790): Belize: 1, Costa Rica: 4, Cuba: 4, Dominican Rep.: 1, El Salvador: 1, Guatemala: 2, Honduras: 3, Jamaica: 1, Mexico: 3, Panama: 3, USA: 4; Total **27**
guttata (Olivier, 1790): USA: 28; Total **28**
insubida (Boheman, 1855): Colombia: 13, Panama: 3; Total **16**
lecontei (Crotch, 1873): USA: 11; Total **11**
trilineata (Spaeth, 1936): Costa Rica: 1; Total **1**
zetterstedti (Boheman, 1855): El Salvador: 1, Mexico: 5; Total **6**

Genus *Drepanocassis* Spaeth, 1936

profana (Boheman, 1855): Brazil: 10; Total **10**

Genus *Eremionycha* Spaeth, 1911

bahiana (Boheman, 1855): Brazil: 1; Total **1**

Genus *Erepsocassis* Spaeth, 1936

rubella (Boheman, 1862): USA: 1; Total **1**

Genus *Exestastica* Spaeth, 1909

ignobilis (Boheman, 1854): Bolivia: 2, Brazil: 2; Total **4**

Genus *Floridocassis* Spaeth, 1952

repudiata (Suffrian, 1868): USA: 4; Total **4**

Genus *Gratiana* Spaeth, 1913

boliviana Spaeth, 1926: Argentina: 2, Paraguay: 3, Paru: 2; Total **7**
conformis (Boheman, 1854): Argentina: 3, Brazil: 3, Paraguay: 4; Total **10**
graminea (Klug, 1829): Argentina: 7; Total **7**
lutescens (Boheman, 1854): Argentina: 3; 1; Total **3**
pallidula (Boheman, 1854): Mexico: 2, USA: 8; Total **10**
spadicea (Klug, 1829): Brazil: 5, Paraguay: 11; Total **16**

Genus *Helocassis* Spaeth, 1952

clavata (Fabricius, 1798): Canada: 2, Mexico: 1, USA: 8; Total **11**
crucipennis (Boheman, 1855): Costa Rica: 1, El Salvador: 1, Guatemala: 1, Mexico: 8; Total **11**
distorta (Boheman, 1855): Mexico: 2; Total **2**
flavorugosa (Boheman, 1855): Brazil: 3; Total **3**
incurva (Boheman, 1855): Brazil: 2; Total **2**
maculicollis (Champion, 1894): Mexico: 2; Total **2**
montana Sekerka & Windsor in litt.: Panama: 1 (**1 PT**); Total **1**
testudinaria (Boheman, 1855): Colombia: 2, Costa Rica: 1, El Salvador: 1, Guatemala: 6; Total **10**
Helocassis sp.: Honduras: 2; Total **2**

Genus *Hybosa* Duponchel, 1842

acutangula Spaeth, 1913 a: 163: Brazil: 4 (**1 ST**), Paraguay: 9; Total **13**
cynthia Boheman, 1855: Brazil: 4, Paraguay: 2; Total **6**
fornicata Boheman, 1855: Bolivia: 2, Brazil: 2, Paraguay: 1; Total **5**
galbanata Boheman, 1855: Venezuela: 2; Total **2**
gibbera Boheman, 1855: Brazil: 5; Total **5**
insculpta Spaeth, 1937: Brazil: 4; Total **4**
marginepunctata Borowiec, 2011: Brazil: 2; Total **2**
mellicula Boheman, 1862: Brazil: 8, Colombia: 1, Panama: 1; Total **10**
santaritae Borowiec & Takizawa, 2011 e: 469: French Guyana: 1 (**1 HT**); Total **1**

Genus *Ischnocodia* Spaeth, 1942

annulus (Fabricius, 1781): Argentina: 1, Costa Rica: 1, Ecuador: 1, Panama: 9, Peru: 6, Trinidad and Tobago: 1, Venezuela: 2, no country: 1; Total **22**
succincta (Boheman, 1855): Brazil: 3 (**2 PT** *Ischnocodia flavofasciata* Borowiec, 1995 b: 91); Total **3**

Genus *Jonthonota* Spaeth, 1913

mexicana (Champion, 1894): Mexico: 2, USA: 2; Total **4**
nigripes (Olivier, 1790): USA: 17; Total **17**

Genus *Leptocodia* Spaeth, 1952

luctifera (Boheman, 1855): Brazil: 1; Total **1**

Genus *Metriona* Weise, 1896

elator (Klug, 1829): Brazil: 24; Total **24**

Genus *Metrionella* Spaeth, 1932

angularis (Champion, 1894): Costa Rica: 9; Total **9**
bilimeki Spaeth, 1932: Costa Rica: 2, El Salvador: 4, Guatemala: 1, Mexico: 2, Venezuela: 1; Total **10**
calva (Boheman, 1855): Brazil: 3, Paraguay: 1; Total **4**
chiriquiana Sekerka & Windsor in litt.: Panama: 1 (**1 PT**); Total **1**
connata Spaeth, 1932: Brazil: 1; Total **1**
erratica (Boheman, 1855): El Salvador: 1, Guatemala: 7, Honduras: 1, Mexico: 5, no country: 2; Total **16**
glabrescens Spaeth, 1932: Bolivia: 1; Total **1**

irrorata Spaeth, 1932: Peru: 2; Total **2**
metrionelloides Sekerka & Windsor in litt.: Panama: 3 (**2 PT**); Total **3**
placans Spaeth, 1932: Argentina: 1, Brazil: 7, Ecuador: 1, French Guyana: 7, Guyana: 1, Paraguay: 4, no country: 1, Peru: 3; Total **25**
strandii Spaeth, 1932: Peru: 1; Total **1**
tucumana Borowiec, 2006 i: 568: Argentina: 1 (**1 HT**); Total **1**
tumacoensis Borowiec, 2002 b: 139: Colombia: 1 (**1 HT**), Ecuador: 1; Total **2**
Metrionella sp.: Bolivia: 2; Total **2**

Genus *Mexicaspis* Spaeth, 1936

azteca (Champion, 1894): Mexico: 2; Total **2**

Genus *Microctenochira* Spaeth, 1926

aberrata (Weise, 1904): Costa Rica: 1; Total **1**
achardi (Spaeth, 1926): Argentina: 2, Paraguay: 1; Total **3**
aciculata (Boheman, 1855): Brazil: 6; Total **6**
annulata (Spaeth, 1926): Brazil: 2; Total **2**
anxia (Boheman, 1855): Brazil: 1; Total **1**
aspersa (Champion, 1894): Costa Rica: 2, Panama: 2; Total **4**
belizensis Borowiec, 2007 g: 488: Belize: 1 (**1 HT**); Total **1**
bifenestrata (Boheman, 1855): Costa Rica: 4, Ecuador: 1, Peru: 1; Total **6**
biolleyi (Spaeth, 1926): Costa Rica: 9; Total **9**
bonvouloiri (Boheman, 1862): Mexico: 6; Total **6**
brasiliensis Świętojańska & Borowiec, 1999 b: 109: Brazil: 1 (**1 HT**); Total **1**
championi (Spaeth, 1926): Panama: 2; Total **2**
chapada Świętojańska & Borowiec, 1995 h: 451: Brazil: 2 (**2 PT**); Total **2**
circinaria (Erichson, 1847): Bolivia: 1, Brazil: 4, Peru: 5; Total **10**
circumcincta (Boheman, 1855): Brazil: 2; Total **2**
conscripta (Boheman, 1855): French Guyana: 6; Total **6**
coronata (Boheman, 1855): El Salvador: 4, Guatemala: 1; Total **5**
cruxflava (Champion, 1894): Costa Rica: 1; Total **1**
cumulata (Boheman, 1855): Costa Rica: 7, Ecuador: 4, Guatemala: 1; Total **12**
danielssoni Borowiec, 1995 in Świętojańska & Borowiec, 1995 h: 447: Ecuador: 1, Peru: 2 (**1 PT**); Total **3**
decora (Spaeth, 1926): Ecuador: 2; Total **2**
diabolica (Spaeth, 1926): Ecuador: 2, Peru: 1; Total **3**
difficilis (Boheman, 1855): Brazil: 6; Total **6**
diffinis (Boheman, 1855): Brazil: 2, Colombia: 1, Ecuador: 1, French Guyana: 1, Surinam: 1, no country: 1; Total **7**
discrepans (Spaeth, 1926): Brazil: 1, French Guyana: 2, Guyana: 3, Peru: 1; Total **7**
dissimilis (Boheman, 1855): Guatemala: 1, Mexico: 1; Total **2**
dissoluta (Spaeth, 1901): Bolivia: 5, Peru: 2, no country: 2; Total **9**
excelsa (Spaeth, 1926): Costa Rica: 5, Panama: 6; Total **11**
excurrens (Spaeth, 1926): Ecuador: 2; Total **2**
fairmairei (Boheman, 1855): Bolivia: 3, Peru: 2; Total **5**
ferranti (Spaeth, 1926): Costa Rica: 1, El Salvador: 3, Mexico: 4; Total **8**
flavonotata (Boheman, 1855): Colombia: 1, Costa Rica: 7, Honduras: 1, Panama: 5, Venezuela: 1; Total **15**
fraterna (Boheman, 1855): Colombia: 4, Costa Rica: 6, Venezuela: 3; Total **13**
freyi (Boheman, 1862): Brazil: 3; Total **3**
gemina (Boheman, 1855): Brazil: 5; Total **5**
gnata (Spaeth, 1926): Brazil: 1; Total **1**
hectica (Boheman, 1855): Costa Rica: 3, El Salvador: 4, Guatemala: 2, Mexico: 1; Total **10**
hieroglyphica (Boheman, 1855): Mexico: 7; Total **7**

infantula (Boheman, 1862): Guatemala: 1; Total **1**
jousselini (Boheman, 1855): Trinidad and Tobago: 1; Total **1**
libidinosa (Spaeth, 1926 a: 67): Colombia: 3 (**1 ST**); Total **3**
lindigi (Kirsch, 1865): Colombia: 3, Ecuador: 2, Peru: 1; Total **6**
liquidata (Spaeth, 1926): Brazil: 2; Total **2**
lugubris (Boheman, 1862): Costa Rica: 2, Panama: 2; Total **4**
mapiriensis Borowiec, 2002 f: 366: Bolivia: 2 (**1 HT, 1 PT**); Total **2**
marginata (Spaeth, 1909): Ecuador: 1, Peru: 1; Total **2**
media (Boheman, 1855): Brazil: 3; Total **3**
melanota (Boheman, 1855): Colombia: 2, Costa Rica: 1, Guatemala: 1; Total **4**
minax (Spaeth, 1926): Ecuador: 3; Total **3**
mystica (Boheman, 1855): French Guyana: 1, Peru: 2; Total **3**
nigrocincta (Wagener, 1877): Costa Rica: 7, Panama: 5; Total **12**
obscurata Świętojańska & Borowiec, 1999 g: 112: Brazil: 1 (**1 HT**); Total **1**
optata (Boheman, 1855): Brazil: 20; Total **20**
panamensis Świętojańska & Borowiec, 1999 g: 109: Panama: 2 (**1 HT**); Total **2**
papulosa (Boheman, 1855): French Guyana: 4, Surinam: 1; Total **5**
patruelis (Boheman, 1855): Brasil: 1; Total **1**
peltata (Boheman, 1855): Colombia: 2, Ecuador: 1, Peru: 4; Total **7**
plebeja (Boheman, 1855): Mexico: 5; Total **5**
plicata (Boheman, 1855): Costa Rica: 3, Mexico: 1, Panama: 3; Total **7**
porosa (Boheman, 1855): Brazil: 1, Colombia: 1; Total **2**
punicea (Boheman, 1855): Belize: 1, El Salvador: 1, Guatemala: 3, Mexico: 5; Total **10**
quadrata (Degeer, 1775): Brazil: 6, Colombia: 2, French Guyana: 7, Paraguay: 2, Surinam: 1; Total **18**
reticularis (Degeer, 1775): Bolivia: 2, Brazil: 7, Colombia: 1, Ecuador: 2, French Guyana: 2, Panama: 1, Peru: 20, Surinam: 1, Trinidad and Tobago: 8; Total **44**
rubrocincta (Boheman, 1855): Costa Rica: 2, Mexico: 8, Panama: 7; Total **17**
sagulata (Boheman, 1862): Costa Rica: 4; Total **4**
salebrata (Boheman, 1862): Costa Rica: 14; Total **14**
sanguinidorsis (Spaeth, 1926): Costa Rica: 10; Total **10**
semilobata (Wagener, 1877): Colombia: 1; Total **1**
semilunaris (Boheman, 1862): Bolivia: 1, Brazil: 3, Colombia: 1, Ecuador: 1, Peru: 2; Total **8**
sertata (Erichson, 1847): Bolivia: 2, Brazil: 3, Colombia: 5, Ecuador: 4, French Guyana: 2, Panama: 1, Peru: 14, Surinam: 1; Total **32**
severa (Boheman, 1855): Brazil: 1; Total **1**
similata (Boheman, 1855): Brazil: 7, Paraguay: 1; Total **8**
soleifera (Spaeth, 1926): Brazil: 3; Total **3**
stigmatica (Boheman, 1855): Brazil: 6, no country: 1; Total **7**
tabida (Boheman, 1855): Mexico: 6, Panama: 6; Total **12**
trepida (Boheman, 1855): Brazil: 5; Total **5**
villica (Boheman, 1855): Brazil: 1; Total **1**
vivida (Boheman, 1855): Costa Rica: 1, Mexico: 2, Panama: 1; Total **4**
Microctenochira sp.: Honduras: 1; Total **1**

Genus *Nuzonia* Spaeth, 1912

atromaculata Borowiec, 2000 d: 62: Brazil: 1 (**1 HT**); Total **1**
brevicornis Borowiec, 1998 d: 237: Ecuador: 1 (**1 HT**); Total **1**
cayennensis (Boheman, 1855): Brazil: 1, French Guyana: 1; Total **2**
gestatrix (Boheman, 1855): French Guyana: 4; Total **4**
laqueifera (Kirsch, 1876): Bolivia: 1; Total **1**
marginepunctata Borowiec, 2000 d: 64: Colombia: 2 (**1 HT, 1 PT**); Total **2**
uniformis (Boheman, 1855): Australia (misabeled): 1, French Guyana: 3; Total **4**

Nuzonia sp. 1: Brazil: 1; Total **1**

Nuzonia sp. 2: Brazil: 1; Total **1**

Genus *Opacinota* Riley, 1986

bisignata (Boheman, 1855): USA: 5; Total **5**

Genus *Orexita* Spaeth, 1911

blanchardi (Boheman, 1855): Bolivia: 2, Peru: 2; Total **4**

blattoides Spaeth, 1911: Ecuador: 3; Total **3**

complanata (Boheman, 1855): Brazil: 3; Total **3**

justini (Boheman, 1855): Colombia: 1, Ecuador: 1; Total **2**

maura (Boheman, 1855: 98): Brazil: 4 (**1 PLT**); Total **4**

minima Borowiec & Świętojańska, 1997 c: 679: Brazil: 4 (**4 PT**); Total **4**

picta (Boheman, 1855): Brazil: 11, French Guyana (mislabeled): 1, USA (California, introduced): 1; Total **13**

subgibbosa Spaeth, 1911: Colombia: 1; Total **1**

subopaca Spaeth, 1919: Peru: 3; Total **3**

suggesta Spaeth, 1928: Ecuador: 2; Total **2**

tripartita (Champion, 1894): Costa Rica: 1; Total **1**

varians (Guérin, 1844): Bolivia: 8, Brazil: 6; Total **14**

wagneri (Boheman, 1862): Colombia: 1, Costa Rica: 5, Panama: 1; Total **7**

Orexita sp. 1: Brazil: 1; Total **1**

Orexita sp. 2: Peru: 1; Total **1**

Genus *Parachirida* Hincks, 1952

flavolineata (Latreille, 1811): Colombia: 6; Total **6**

pustulata (Boheman, 1855): Brazil: 4, French Guyana: 2, Peru: 2; Total **8**

semiannulata (Boheman, 1855): Bolivia: 1, Brazil: 2, Colombia: 4, Peru: 3; Total **10**

subirrorata (Boheman, 1855): Colombia: 3, Panama: 7, Trinidad and Tobago: 1, Venezuela: 5; Total **16**

Genus *Parorectis* Spaeth, 1901

callosa (Boheman, 1854): USA: 3; Total **3**

rugosa (Boheman, 1854): Costa Rica: 1, El Salvador: 2, Honduras: 1, Mexico: 6, no country: 1; Total **11**

sublaevis (Barber, 1946): USA: 3; Total **3**

Genus *Plagiometriona* Spaeth, 1899

ambigena (Boheman, 1855): Brazil: 6; Total **6**

amplexa (Erichson, 1847): Bolivia: 2, Peru: 3; Total **5**

boliviana Spaeth, 1912: Bolivia: 1, Peru: 2; Total **3**

boschmai Spaeth, 1937: Colombia: 8, Ecuador: 2; Total **10**

buqueti (Boheman, 1855): Colombia: 2, Venezuela: 1; Total **3**

clandestina (Boheman, 1855): Brazil: 10; Total **10**

clarki (Boheman, 1862): Bolivia: 3, Ecuador: 4, Peru: 2; Total **9**

coccinelloides (Boheman, 1855): Brazil: 5; Total **5**

commixta Spaeth, 1937: 156: Costa Rica: 1 (**1 ST**), Panama: 1; Total **2**

constricta (Boheman, 1855): Brazil: 5; Total **5**

costaricensis Borowiec, 2001 h: 356: Costa Rica: 2 (**1 HT, 1 PT**); Total **2**

deyrollei (Boheman, 1855): Brazil: 3; Total **3**

dodonea (Boheman, 1855): Brazil: 3; Total **3**

dorsosignata (Boheman, 1855): Brazil: 9; Total **9**

eggi Spaeth, 1899: Peru: 1; Total **1**

flavescens (Boheman, 1855): Brazil: 11; Total **11**

foliata (Boheman, 1855): Bolivia: 3, Colombia: 1, Peru: 4; Total **8**

forcipata (Boheman, 1855): Brazil: 16; Total **16**
gibbifera (Champion, 1894): Costa Rica: 1; Total **1**
glyphica (Boheman, 1855): Colombia: 1; Total **1**
gyrata (Boheman, 1855): Brazil: 5; Total **5**
herbea (Boheman, 1855): Brazil: 6; Total **6**
inscripta (Boheman, 1855): Brazil: 2; Total **2**
jucunda (Kirsch, 1876): Ecuador: 1, peru: 2; Total **3**
latemarginata Borowiec, 2001 h: Panama: 1 (1 **HT**); Total **1**
latifoliata Spaeth, 1937: Ecuador: 1; Total **1**
liturata (Boheman, 1855): Brazil: 2; Total **2**
losi Borowiec, 1998 d: 239: Ecuador: 1 (1 **HT**); Total **1**
ludicra (Boheman, 1855): Brazil: 4; Total **4**
microcera (Boheman, 1855): Brazil: 1; Total **1**
nympha (Boheman, 1855): Brazil: 1; Total **1**
paleacea (Boheman, 1855): Brazil: 5; Total **5**
pehlkei Spaeth, 1912: Colombia: 8; Total **8**
pernix Spaeth, 1912: 10: Colombia: 1 (1 **ST**); Total **1**
perroudi (Boheman, 1862): Colombia: 1; Total **1**
peruana Spaeth, 1912: Bolivia: 4, Colombia: 1, Peru: 5; Total **10**
phoebe (Boheman, 1855): Brazil: 4, Colombia: 4, French Guyana: 7, no country: 1; Total **16**
praecincta (Boheman, 1855): Brazil: 3; Total **3**
punctatissima (Boheman, 1855): Argentina: 1, Brazil: 6, Paraguay: 1; Total **8**
sahlbergi (Boheman, 1855): Brazil: 7; Total **7**
seriata Spaeth, 1937: 147: Brazil: 1 (1 **ST**); Total **1**
steinheili (Wagener, 1877): Ecuador: 4 (1 **ST** *Plagiometriona steinheili* ab. *fulvitaris* Spath, 1937: 151); Total **4**
stillata (Boheman, 1855): Brazil: 3; Total **3**
subprasina (Spaeth, 1902): Brazil: 6; Total **6**
tenella (Klug, 1829): Argentina: 9, Brazil: 19, Costa Rica (introduced): 1, Paraguay: 10; Total **39**
tredecimguttata (Boheman, 1862): Brazil: 4; Total **4**
vesiculifera (Boheman, 1862): Ecuador: 1; Total **1**
vespertilio (Spaeth, 1902): Peru: 5; Total **5**
vigens (Boheman, 1855): Brazil: 6; Total **6**
zelleri (Boheman, 1855): Colombia: 3; Total **3**
Plagiometriona sp. 1: Brazil: 2; Total **2**
Plagiometriona sp. 2: Ecuador: 2; Total **2**
Plagiometriona sp. 3: Ecuador: 3; Total **3**
Plagiometriona sp. 4: Ecuador: 3; Total **3**
Plagiometriona sp. 5: Country?: 3; Total **3**

Genus *Pseudoctenochira* Spaeth, 1926

octolineata Spaeth, 1926: Bolivia: 1; Total **1**

Genus *Rhoia* Spaeth, 1913

clavareau Spaeth, 1913: Brazil: 1; Total **1**

Genus *Saulaspis* Spaeth, 1913

bistrilineata (Boheman, 1854): Argentina: 2, French Guyana (misabeled): 1, Uruguay: 1, no country: 2; Total **6**

quadrivittata (Boheman, 1855): Brazil: 3; Total **3**

trivittata Świętojańska, 2002 c: 580: Paraguay: 1 (1 **HT**); Total **1**

Genus *Scaecassis* Spaeth, 1913

combusta (Boheman, 1854): Brazil: 7; Total **7**

corumbana Spaeth, 1937: Brazil: 1; Total **1**
turbulenta (Boheman, 1862): Brazil: 2; Total **2**
unimaculata Spaeth, 1915: Peru: 1; Total **1**

Genus *Strongylocassis* Hincks, 1950

atripes (Leconte, 1859): USA: 8; Total **8**

Genus *Syngambria* Spaeth, 1911

andreae (Boheman, 1855): Brazil: 4; Total **4**
bisinuata (Boheman, 1855): Bolivia: 1, Brazil: 6, Paraguay: 2; Total **9**

Genus *Tapinaspis* Spaeth, 1936

atroannulus (Champion, 1894): Costa Rica: 1; Total **1**
wesmaeli (Boheman, 1855): Guatemala: 6, Mexico: 3; Total **9**
Tapinaspis sp.: Mexico: 1; Total **1**

Genus *Tetracassis* Spaeth, 1952

lyra (Spaeth, 1898): Brazil: 2; Total **2**

Tribe Delocraniini

Genus *Delocrania* Guérin, 1844

cosyphoides Guérin, 1844: Venezuela: 2; Total **2**
latipennis Champion, 1893: Ecuador: 1, Peru: 1; Total **2**
panamensis Champion, 1893: Costa Rica: 1, Panama: 2; Total **3**

Tribe Dorynotini

Genus *Dorynota* Chevrolat, 1836

Subgenus: *Dorynota* s. str.

aculeata (Boheman, 1854): Haiti: 1, Dominican Rep.: 2; Total **3**
apiculata (Boheman, 1854): Brazil: 1; Total **1**
aurita (Boheman, 1862): Mexico: 2; Total **2**
bidens (Fabricius, 1781): Brazil: 9; Total **9**
cornigera (Boheman, 1854): Brazil: 2, Paraguay: 3; Total **5**
monoceros (Germar, 1824): Argentina: 2, Brazil: 7, Paraguay: 2; Total **11**
nigra (Boheman, 1856): Trinidad: 1, Venezuela: 2; Total **3**
nodosa (Boheman, 1854): Colombia: 1, Panama: 1; Total **2**
ohausi (Spaeth, 1915): Ecuador: 1; Total **1**
pubescens (Blake, 1939): Haiti: 1; Total **1**
pugionata (Germar, 1824): Brazil: 12, Paraguay: 4; Total **16**
rileyi Borowiec, 1994 b: 161: Paraguay: 2 (**2 PT**); Total **2**
yucatanana (Champion, 1893): Mexico: 1; Total **1**
Dorynota s. str. sp. 1: Guyana: 1, Venezuela: 4; Total **5**
Dorynota s. str. sp. 2: Colombia: 1; Total **1**

Subgenus: *Akantaka* Maulik, 1916

aeneocincta (Spaeth, 1913): Brazil: 3; Total **3**
biplagiata (Champion, 1893): Panama: 5; Total **5**
bivittipennis (Boheman, 1856): Brazil: 3, Colombia: 1, Peru: 3; Total **7**
boliviana Borowiec, 2005: Bolivia: 2; Total **2**
carlosi Borowiec & Takizawa, 2011 e: 465: Peru: 1 (**1 PT**); Total **1**

collucens (Spaeth, 1923): Bolivia: 5; Total **5**
dejeanii (Boheman, 1854): French Guyana: 2; Total **2**
distincta (Baly, 1872): Ecuador: 3; Total **3**
electa (Spaeth, 1923): Brazil: 2, Peru: 11; Total **13**
funesta (Boheman, 1854): Brazil: 2, French Guyana: 1; Total **3**
godmani (Baly, 1864): Panama: 1; Total **1**
incapabilis (Spaeth, 1923): Brazil: 8; Total **8**
insidiosa (Boheman, 1854): Colombia: 2, Mexico: 1, no country: 1; Total **4**
kiesenwetteri (Boheman, 1854): Bolivia: 2, Colombia: 1, Peru: 2, Venezuela: 1; Total **6**
matogrossoensis Borowiec, 2005 b: 30: Brazil: 1 (**1 HT**); Total **1**
rufoornata (Baly, 1869): Nicaragua: 1; Total **1**
truncata (Fabricius, 1781): French Guyana: 10, Surinam: 1; Total **11**
viridimetallica Borowiec, 2005 b: 37: Brazil: 2 (**2 PT**); Total **2**
viridisignata (Boheman, 1854): Bolivia: 2, Brazil: 3, Paraguay: 4; Total **9**
Dorynota (*Akantaka*) sp. 1: Colombia: 1, Costa Rica: 1; Total **2**

Genus *Omoteina* Chevrolat, 1836

humeralis (Olivier, 1808): Haiti: 3; Total **3**

Genus *Paranota* Monros and Viana, 1949

ensifera (Boheman, 1854): Argentina: 2, Bolivia: 1, Brazil: 26, Paraguay: 1; Total **30**
minima (Wagner, 1881): Bolivia: 2, Paraguay: 3; Total **5**
parallela (Blanchard, 1837): Bolivia: 1, Brazil: 13, French Guyana (mis-labeled!): 1, Paraguay: 3; Total **18**
spinosa (Boheman, 1854): Brazil: 8; Total **8**

Genus *Paratrikona* Spaeth, 1923

lerouxii (Boheman, 1854): Cuba: 5; Total **5**
rubescens Blake, 1939: Dominican Rep.: 1 (**1 HT** *Paratrikona albomaculata* Borowiec, 2009 g: 567); Total **1**
turrifera (Boheman, 1854): Dominican Rep.: 1, Haiti: 1; Total **2**

Tribe Eugenysini

Genus *Agenysa* Spaeth, 1905

boliviana Spaeth, 1915: Bolivia: 3; Total **3**
caedemadens (Lichtenstein, 1795): Brazil: 15, French Guyana: 13, Peru: 1; Total **29**
connectens (Baly, 1869): Brazil: 1, Colombia: 2, Ecuador: 14, Panama: 1, Peru: 6; Total **24**
crassicornis Spaeth, 1905: Ecuador: 22; Total **22**
desmaresti (Boheman, 1850): no country: 1; Total **1**
guianiensis (Boheman, 1856): Brazil: 6, Surinam: 1, Venezuela: 2; Total **9**
parellina (Boheman, 1850): Argentina: 1, Brazil: 7; Total **8**
peruviana Spaeth, 1915: 275: Peru: 17 (**1 ST**); Total **17**
Agenysa sp.: Ecuador: 2, Panama: 1, Peru: 5; Total **8**

Genus *Eugenysa* Chevrolat, 1836

bacchus (Baly, 1869): Brazil: 6; Total **6**
bohemani (Baly, 1859): Ecuador: 1, Peru: 1; Total **2**
colossa (Boheman, 1862): Ecuador: 1, Peru: 4; Total **5**
colombiana (Boheman, 1850): Colombia: 5, Panama: 2; Total **7**
coscaroni Viana, 1968: Panama: 1; Total **1**
cuprifulgens (Spaeth, 1919): Peru: 4; Total **4**
delicata (Boheman, 1862): Brazil: 6; Total **6**
diringshofeni Viana, 1968: Brazil: 3; Total **3**

divalis (Boheman, 1862): Peru: 14; Total **14**
grossa (Linnaeus, 1758): French Guyana: 17; Total **17**
imperialis (Boheman, 1856): Brazil: 1; Total **1**
jasinskii Borowiec & Dąbrowska, 1997 b: 673: Ecuador: 4 (**1 HT**); Total **1**
lata (Boheman, 1850): Bolivia: 4; Total **4**
martae Borowiec, 1987 a: 883: Colombia: 2 (**1 HT, 1 PT**); Total **2**
regalis (Boheman, 1850): Brazil: 2; Total **2**
subopaca (Spaeth, 1919): Ecuador: 1; Total **1**
unicolor Borowiec & Dąbrowska, 1997 b: 675: Ecuador: 9 (**1 HT, 4 PT**); Total **9**
venosa (Fabricius, 1798): French Guyana: 11; Total **11**
willineri Viana, 1968: Peru: 1; Total **1**

Genus *Miocalaspis* Weise, 1899

alurna (Westwood, 1842): Colombia: 1, Ecuador: 1; Total **2**
biplagosa Spaeth, 1905: Bolivia: 1; Total **1**
ecuadorica Borowiec, 2000 g: 231: Ecuador: 1 (**1 HT**); Total **1**
flavofasciata Borowiec, 2000 g: 229: Peru: 1 (**1 HT**); Total **1**
gentilis (Erichson, 1847): Bolivia: 2, Peru: 7; Total **9**

Tribe Goniocheniini

Genus: *Chlamydocassis* Spaeth, 1952

Subgenus *Chlamydocassis* s. str.

cribripennis (Boheman, 1850): Argentina: 13, Brazil: 15 (**1 ST** of *Desmonota refractaria* Boheman, 1862: 66), Paraguay: 30, Peru: 1, Paraguay: 1; Total **60**
intermedia (Boheman, 1850): Brazil: 1; Total **1**
metallica (Klug, 1829): Argentina: 7, Brazil: 6; Total **13**
perforata (Boheman, 1850): Bolivia: 7; Total **7**
runderaria (Erichson, 1847): Bolivia: 1, Peru: 4; Total **5**

Subgenus *Ceratocassis* Viana, 1964

bicornuta (Boheman, 1850): Brazil: 3; Total **3**
bispinosa (Boheman, 1850): Brazil: 1; Total **1**
laticollis (Boheman, 1850): Argentina: 3, Bolivia: 1, Brazil: 2, Venezuela: 1; Total **7**
subcornuta (Boheman, 1862): Brazil: 14, Surinam: 1; Total **15**

Genus: *Goniochenia* Weise, 1896

Subgenus *Goniochenia* s. str.

difformis (Boheman, 1850): no country: 1; Total **1**
parvula Weise, 1896: Brazil: 25; Total **25**
quadraticollis (Boheman, 1850): Brazil: 6; Total **6**
virgo Weise, 1896: Brazil: 2; Total **2**
Goniochenia sp. 1: Brazil: 1; Total **1**

Subgenus *Baranosa* Weise, 1899

buckley (Baly, 1872): Ecuador: 5; Total **5**
elocata (Boheman, 1850): Colombia: 4, Ecuador: 2; Total **6**
flavosparsa (Boheman, 1856): Bolivia: 4; Total **4**
haroldi (Wagener, 1877): Bolivia: 2, Peru: 9; Total **11**

Genus: *Herissa* Spaeth, 1909

pantherina (Blanchard, 1837): Brazil: 4, Paraguay: 1; Total **5**

Genus: *Polychalma* Barber & Bridwell, 1940

multicava (Latreille, 1811): Colombia: 6, Ecuador: 2, Panama: 1; Total **9**

Genus: *Zeugonota* Spaeth, 1913

quadrinodosa (Boheman, 1856): Brazil: 3; Total **3**

Tribe Hemisphaerotini

Genus: *Hemisphaerota* Chevrolat, 1836

cyanea (Say, 1823): USA: 6; Total **6**

gundlachi (Boheman, 1862): Cuba: 20; Total **20**

palmarum (Boheman, 1856): Dominican Rep.: 13; Total **13**

Genus: *Spaethiella* Barber & Bridwell, 1940

atrata Sekerka & Windsor in litt.: Panama: 2 (**2 PT**); Total **2**

bromelioides Sekerka & Windsor in litt.: Panama: 1 (**1 PT**); Total **1**

circumdata (Boheman, 1850): Colombia: 1, Ecuador: 1, French Guyana: 1, Peru: 1, Surinam: 1; Total **5**

costipennis (Boheman, 1850): Brazil: 6, French Guyana: 3, Peru: 1; Total **10**

crassicornis (Spaeth, 1910): Brazil: 2, Paraguay: 4; Total **6**

cruenta Sekerka & Windsor in litt.: Panama: 1 (**1 PT**); Total **1**

erhardti (Boheman, 1862: 36): Surinam: 2 (**1 PT**); Total **2**

heliconiae Sekerka & Windsor in litt.: Panama: 11 (**11 PT**); Total **11**

intricata (Boheman, 1850): Brazil: 1; Total **1**

miniata (Boheman, 1856): Colombia: 2 (**1 ST** of *Porphyraspis panamae* Spaeth, 1901 a: 334), Panama: 2; Total **4**

montanella Sekerka & Windsor in litt.: Panama: 1 (**1 PT**); Total **1**

quadrata (Spaeth, 1902): Peru: 1; Total **1**

robusta (Spaeth, 1910): Colombia: 2; Total **2**

rotundior (Spaeth, 1929): Bolivia: 1; Total **1**

sanguinea (Fabricius, 1801): French Guyana: 11, Peru: 1, Surinam: 1; Total **13**

speculicollis (Spaeth, 1928): Brazil: 1, Paraguay: 1; Total **2**

sublaevis (Spaeth, 1901): Colombia: 2; Total **2**

submetallica (Spaeth, 1928: 29): Bolivia: 2 (**1 ST**), Brazil: 1; Total **3**

tristis (Boheman, 1850): Brazil: 5, Ecuador: 1, Peru: 1; Total **7**

valida (Spaeth, 1901): Colombia: 1; Total **1**

sp. nov. 1 – Costa Rica: 4; Total **4**

sp. nov. 2 – Costa Rica: 3; Total **3**

sp. nov. 3 – Costa Rica: 1; Total **1**

sp. nov. 4 – Ecuador: 2; Total **2**

sp. nov. 5 – Guatemala: 1; Total **1**

Tribe Ischyrosomychini

Genus: *Asteriza* Chevrolat, 1836

darlingtoni Blake, 1939: Dominican Rep.: 1; Total **1**

flavicornis (Olivier, 1790): Dominican Rep.: 10; Total **10**

Genus: *Cistudinella* Champion, 1894

apiata (Boheman, 1854): Brazil: 2; Total **2**

foveolata Champion, 1894: Costa Rica: 1, Ecuador: 2, Panama: 1, Venezuela: 1; Total **5**

inanis (Boheman, 1854): Brazil: 1, Paraguay: 1; Total **2**

lata Spaeth, 1932: Peru: 1; Total **1**

lateripunctata Spaeth, 1905: Argentina: 6, Brazil: 1, Paraguay: 10; Total **17**

notata (Boheman, 1854): Argentina: 1, Brazil: 20; Total **21**

obducta (Boheman, 1854): Argentina: 4, Brazil: 33, Paraguay: 9, Peru: 1, Uruguay: 1; Total **48**
parva (Wagener, 1881): Ecuador: 4; Total **4**
peruana Spaeth, 1905: Ecuador: 1, Peru: 5; Total **6**
plagicollis Spaeth, 1905: Peru: 4; Total **4**
punctipennis (Boheman, 1854): Brazil: 6; Total **6**
rufitarsis Spaeth, 1905: Brazil: 4; Total **4**

Genus: *Enagria* Spaeth, 1913

ovata (Boheman, 1854): Mexico: 5; Total **5**

Genus: *Eurypedus* Gistel, 1834

nigrosignatus (Boheman, 1854): Colombia: 11, Guatemala: 3, Panama: 1; Total **15**
peltoides (Boheman, 1854): Argentina: 4, Brazil: 5, Paraguay: 1, no country: 1; Total **11**

Genus: *Eurypepla* Boheman, 1854

brevilineata Boheman, 1854: Mexico: 2; Total **2**
calochroma (Blake, 1965): USA: 12; Total **12**

Genus: *Physonota* Boheman, 1854

alutacea Boheman, 1854: Costa Rica: 9, Ecuador: 2, El Salvador: 2, Guatemala: 1, Mexico: 5, Trinidad: 2, USA: 1, Venezuela: 1; Total **23**
arizonae Schaeffer, 1925: Mexico: 4, USA: 5; Total **9**
attenuata Boheman, 1854: Guatemala: 2, Mexico: 3, Nicaragua: 5; Total **10**
calcarata (Boheman, 1854): Mexico: 6; Total **6**
caudata Boheman, 1854: Mexico: 7; Total **7**
cerea Boheman, 1854: Mexico: 3; Total **3**
citrina Boheman, 1854: Mexico: 5; Total **5**
convexa Borowiec, 1995 f: 430: Mexico: 1 (**1 PT**); Total **1**
dilatata Kirsch, 1876: Peru: 2; Total **2**
disjuncta (Chevrolat, 1834): Mexico: 13; Total **13**
gigantea Boheman, 1854: Chile (mislabelled!): 1, El Salvador: 2, no country: 1; Total **4**
helianthi (Randall, 1838): USA: 13; Total **13**
humilis Boheman, 1856: Mexico: 8; Total **8**
incrustedata Boheman, 1854: Mexico: 1; Total **1**
limoniata Boheman, 1862: Guatemala: 1, Mexico: 5; Total **6**
lutarella Boheman, 1856: Colombia: 3, Vietnam (mislabelled!): 6, no country: 5; Total **14**
maculiventris Boheman, 1854: Mexico: 5; Total **5**
mexicana Boheman, 1854: Mexico: 5; Total **5**
nitidicollis Boheman, 1854: Mexico: 1; Total **1**
ovalis Boheman, 1854: Mexico: 3; Total **3**
pacifica Spaeth, 1932: Mexico: 2; Total **2**
pellucida Wagener, 1887: Colombia: 3; Total **3**
perampla Champion, 1894: Colombia: 2, Costa Rica: 1; Total **3**
picticollis Boheman, 1854: Mexico: 7; Total **7**
puncticollis Borowiec, 1995 f: 434: Mexico: 1 (**1 PT**); Total **1**
stigmatilis Boheman, 1854: Brazil (mislabelled!): 1, Guatemala: 1, Mexico: 3; Total **5**
sublaevigata Spaeth, 1915: Mexico: 5; Total **5**
translucida Boheman, 1854: Mexico: 3; Total **3**
unipunctata (Say, 1823): USA: 12; Total **12**

Genus: *Platycycla* Boheman, 1854

deruta Boheman, 1854: Mexico: 6, no country: 1; Total **7**

Tribe Mesomphaliini

Genus: *Acromis* Chevrolat, 1836

nebulosa (Boheman, 1854): Brazil: 1; Total **1**

sparsa (Boheman, 1854): Bolivia: 6, Brazil: 6, Panama: 4, Peru: 3; Total **19**

spinifex (Linnaeus, 1763): Argentina: 11, Brazil: 16, French Guyana: 2, Paraguay: 3, Peru: 2, Surinam: 1, Trinidad: 2; Total **37**

venosa Erichson, 1847: Ecuador: 5; Peru: 8; Total **13**

Genus: *Amythra* Spaeth, 1913

valida (Boheman, 1855): Brazil: 4; Total **4**

Genus: *Anacassis* Spaeth, 1913

candida (Boheman, 1854): Brazil: 1; Total **1**

cribrum (Klug, 1829): Brazil: 8; Total **8**

dubia (Boheman, 1854): Brazil: 1; Total **1**

exarata (Boheman, 1854): Argentina: 13; Total **13**

fuscata (Klug, 1829): Argentina: 8, Brazil: 16, Paraguay: 1; Total **25**

languida (Boheman, 1854): Brazil: 1; Total **1**

nigroscutata Świętojańska, 2003 b: 512: Uruguay: 1 (**1 HT**); Total **1**

phaeopoda Buzzi, 1976: Brazil: 5; Total **5**

prolixa (Boheman, 1854): Argentina: 11, Brazil: 2, Chile: 1, no country: 1; Total **15**

punctatissima Spaeth, 1913: no country: 1; Total **1**

punctulata (Klug, 1829): Brazil: 6; Total **6**

rubronotata (Boheman, 1854): Brazil: 4; Total **4**

sulcipennis (Boheman, 1854): Argentina: 2, Uruguay: 1; Total **3**

testudinaria (Boheman, 1850): Brazil: 10; Total **10**

xerampelina (Boheman, 1854): Brazil: 5; Total **5**

Anacassis sp.: Brasil: 2, Guatemala (probably mislabeled): 2, Uruguay: 1, no country: 2; Total **7**

Genus: *Anepsiomorpha* Spaeth, 1913

deplanata (Boheman, 1850): Argentina: 5, Bolivia: 5, Brazil: 4; Total **14**

Genus: *Botanochara* Dejean, 1836

angulata (Germar, 1824): Argentina: 1, Brazil: 22; Total **23**

boggianii (Spaeth, 1900): Paraguay: 3; Total **3**

boliviaca (Spaeth, 1926): Bolivia: 2; Total **2**

bonariensis (Boheman, 1850): Argentina: 16, Brazil: 6; Total **22**

cancellata (Boheman, 1850): Argentina: 13, Brazil: 1; Total **14**

convexiuscula (Spaeth, 1922): Peru: 5; Total **5**

decempustulata (Boheman, 1850): Argentina: 17; Total **17**

decemverrucata (Boheman, 1850): Argentina: 1, Bolivia: 1, Brazil: 9, Peru: 1; Total **12**

duodecimnotata (Boheman, 1850): Bolivia: 1, Brazil: 2, Paraguay: 4; Total **7**

duodecimverrucata (Boheman, 1850): Argentina: 18; Total **18**

encausta (Boheman, 1850): Brazil: 2; Total **2**

guttipennis (Spaeth, 1915: 278): Argentina: 6 (**1 ST**), Paraguay: 1; Total **7**

haematodes (Perty, 1834): Brazil: 2, Peru: 1; Total **3**

impressa (Panzer, 1798): Argentina: 9, Brazil: 37, Paraguay: 1; Total **47**

impressipennis (Boheman, 1850): Brazil: 5; Total **5**

intricata (Boheman, 1850): Argentina: 5, Bolivia: 1, Brazil: 1; Total **7**

lurida Spaeth, 1940: Paraguay: 1; Total **1**

macularia (Boheman, 1850): Argentina: 25, Brazil: 1; Total **26**

miniata (Kolbe, 1887): Paraguay: 4; Total **4**
nigropicta (Boheman, 1850): Brazil: 5; Total **5**
octoplagiata (Spaeth, 1909): Argentina: 12; Total **12**
octopustulata (Klug, 1829): Argentina: 2, Brazil: 3; Total **5**
ordinata (Boheman, 1850): Colombia: 1, Panama: 8, Venezuela: 7; Total **16**
ornata (Boheman, 1850): Brazil: 5; Total **5**
pantherina (Klug, 1829): Argentina: 11, Paraguay: 1; Total **12**
pavonia (Boheman, 1850): Uruguay: 1; Total **1**
planipennis (Spaeth, 1899): Argentina: 12, Bolivia: 2; Total **14**
praeifica (Spaeth, 1940): Paraguay: 5, Peru: 2; Total **7**
quinquefasciata (Perty, 1834): Brazil: 3; Total **3**
regina (Boheman, 1862): Brazil: 1, Peru: 2; Total **3**
ricardo (Buzzi, 1998): Brazil: 2 (**2 PT** of *Botanochara matogrossoensis* Borowiec, 2006 a: 128); Total **2**
rubroguttata (Boheman, 1850): Brazil: 4; Total **4**
ruforeticulata (Boheman, 1850): Brazil: 4; Total **4**
sanguinea (Spaeth, 1909): Bolivia: 1, Brazil: 3, Paraguay: 2; Total **6**
sedecimpustulata (Fabricius, 1781): Argentina: 7 (**1 PT** of *Poecilaspis sedecimpustulata* ssp. *argentina* Spaeth, 1940: 160), Bolivia: 1, Brazil: 13; Total **21**
segnis (Boheman, 1862): Brazil: 3, Paraguay: 6; Total **9**
sigillata (Spaeth, 1940): Argentina: 1, Brazil: 1; Total **2**
subnervosa (Boheman, 1850): Argentina: 1, Brazil: 5; Total **6**
tesselata (Burmeister, 1870): Argentina: 2, Brazil: 4, Paraguay: 1; Total **7**
vianai Borowiec, 1989 a: 725: Argentina: 1 (**1 HT**), Brazil: 2; Total **3**

Genus: *Chelymorpha* Chevrolat, 1836

aculeata Borowiec, 2000: Brazil: 2; Total **2**
alternans (Boheman, 1850): Colombia: 4, Costa Rica: 1, Panama: 9, Venezuela: 5; Total **19**
andicola (Spaeth, 1928): Ecuador: 2, Peru: 1; Total **3**
areata (Erichson, 1847): Argentina: 10, Bolivia: 2, Ecuador: 1, Brazil: 2, Peru: 4; Total **19**
boliviana Boheman, 1854: Argentina: 6, Bolivia: 1; Total **7**
bullata Boheman, 1854: Brazil: 7, Venezuela: 2; Total **9**
cassidea (Fabricius, 1775): Cuba: 1, Mexico: 1, USA: 17; Total **19**
cavata Boheman, 1854: Brazil: 1, Venezuela: 4; Total **5**
cingulata Boheman, 1854: Peru: 4; Total **4**
circumpunctata (Klug, 1829): Brazil: 7; Total **7**
clathrata Spaeth, 1909: Brazil: 2; Total **2**
clivosa Boheman, 1854: Brazil: 8; Total **8**
cobaltina Boheman, 1854: Brazil: 2; Total **2**
comata Boheman, 1854: Belize: 1, Cuba: 10, El Salvador: 1, Guatemala: 2, Mexico: 8, Panama: 1; Total **23**
commutabilis Boheman, 1854: Brazil: 2; Total **2**
constellata (Klug, 1829): Brazil: 17, Peru: 2; Total **19**
costaricensis (Spaeth, 1922): Costa Rica: 2; Total **2**
cribraria (Fabricius, 1775): Argentina: 6, Bolivia: 2, Brazil: 29, Dominican Rep.: 2, Ecuador: 1, French Guyana: 1, Paraguay: 10, Peru: 4, Venezuela: 5, no country: 1; Total **61**
gressoria Boheman, 1862: Costa Rica: 2, El Salvador: 1, Guatemala: 7, Nicaragua: 2; Total **12**
hoepfneri Boheman, 1854: Mexico: 5; Total **5**
indigesta Boheman, 1854: Argentina: 4; Total **4**
infecta Boheman, 1854: Argentina: 10; Total **10**
infirma Boheman, 1854: Colombia: 3; Total **3**
inflata Boheman, 1854: Argentina: 15, Brazil: 40, Paraguay: 1; Total **56**
insignis (Klug, 1829): Argentina: 1, Brazil: 1, no country: 2; Total **4**
klugii Boheman, 1854: Argentina: 2; Total **2**

limbatipennis Spaeth, 1926: Brazil: 1; Total **1**
marginata (Linnaeus, 1758): Argentina: 2, Bolivia: 1, Colombia: 1, French Guyana: 12, Brazil: 5, Paraguay: 1, Venezuela: 1; Total **23**
militaris Boheman, 1862: Costa Rica: 3, Ecuador: 1; Total **4**
multipecta Boheman, 1862: Brazil: 1; Total **1**
nigricollis Boheman, 1854: Brazil: 5; Total **5**
orthogonia Boheman, 1854: Argentina: 1, Brazil: 7, Paraguay: 4; Total **12**
partita Boheman, 1854: Brazil: 1; Total **1**
personata Boheman, 1854: Argentina: 3; Total **3**
peruana Spaeth, 1902: Ecuador: 2, Peru: 3; Total **5**
phytophagica Crotch, 1873: USA: 9; Total **9**
praetextata Boheman, 1854: Argentina: 2, Colombia: 1, Ecuador: 1, Peru: 11; Total **15**
pubescens Boheman, 1854: Mexico: 5; Total **5**
punctigera Boheman, 1854: Brazil: 2; Total **2**
reimoseri Spaeth, 1928: Argentina: 23, Bolivia: 1, Brazil: 3; Total **27**
rosarioensis Buzzi, 2000: Brazil: 1; Total **1**
rufoguttata Spaeth, 1909: Peru: 4; Total **4**
sericea Boheman, 1862: Costa Rica: 4; Total **4**
socia Boheman, 1854: Brazil: 4; Total **4**
subpunctata Boheman, 1854: Brazil: 4; Total **4**
tessellata Spaeth, 1928: Paraguay: 4; Total **4**
texta Boheman, 1862: Paraguay: 2; Total **2**
trinitatis Spaeth, 1926: Trinidad: 2; Total **2**
varians (Blanchard, 1851): Argentina: 19, Chile: 12, Bolivia: 12, Brazil: 1; Total **44**
variolosa (Olivier, 1790): Brazil: 2, French Guyana: 23, Guyana: 1; Total **26**
vermiculata Boheman, 1854: Brazil: 4; Total **4**
wollastoni Boheman, 1854: Surinam: 1; Total **1**

Genus: *Cyrtonota* Chevrolat, 1836

abrili Borowiec & Świątojańska, 2015 d: 88: Colombia: 1 (**1 PT**); Total **1**
adpersa (Boheman, 1850): Brazil: 5; Total **5**
assimilis (Boheman, 1850): Brazil: 3; Total **3**
balyi (Kirsch, 1883): Ecuador: 2 (**1 HT** of *Cyrtonota christophori* Borowiec, 1998 d: 219); Total **2**
banghaasi (Spaeth, 1902): Peru: 5; Total **5**
bergeali Borowiec & Sassi, 1999 e: 89: Colombia: 8 (**8 PT**); Total **8**
bistigma (Boheman, 1856): Bolivia: 1, Peru: 3; Total **4**
bondari (Spaeth, 1928): Brazil: 1; Total **1**
botanocharoides Borowiec, 1989 c: 89: Bolivia: 2 (**2 PT**); Total **2**
caudata (Boheman, 1850): Colombia: 4; Total **4**
compulsa (Spaeth, 1909): Colombia: 2; Total **2**
conglomerata (Boheman, 1862): Argentina: 16, Brazil: 7, Paraguay: 4; Total **27**
cyanea (Linnaeus, 1758): Brazil: 75; Total **75**
deliciosa (Baly, 1872): Ecuador: 4; Total **4**
dimidiata (Boheman, 1850): Brazil: 4; Total **4**
dissecta (Boheman, 1850): Colombia: 15; Total **15**
elongata (Spaeth, 1932): Ecuador: 4; Total **4**
gibbera Borowiec, 1989 c: 28 Colombia: 1 (**1 HT**); Total **1**
goryi (Boheman, 1850): Colombia: 8; Total **8**
inspicata (Spaeth, 1913): Bolivia: 2; Total **2**
jekeli (Boheman, 1856): Brazil: 2; Total **2**
kolbei (Spaeth, 1907): Ecuador: 2; Total **2**
lateralis (Linnaeus, 1758): Brazil: 1, French Guyana: 8, Guyana: 1, Surinam: 1, no country: 1; Total **12**

machupicchu Borowiec & Sassi, 1999 e: 95: Peru: 9 (**1 HT, 2 PT**); Total **9**
marginata (Kirsch, 1883): Ecuador: 1; Total **1**
moderata (Spaeth, 1913): Colombia: 2; Total **2**
montana Borowiec, 2000 b: Ecuador: 2 (**1 HT, 1 PT**); Total **2**
nitida Borowiec & Sassi, 1999 e: 97: Paraguay: 11 (**1 HT, 1 PT**); Total **11**
paupertina (Boheman, 1862): Brazil: 1 (probably mislabeled), Ecuador: 6; Total **7**
plumbea (Germar, 1824): Brazil: 7; Total **7**
poecilaspoides (Baly, 1872): Ecuador: 4; Total **4**
punctatissima (Spaeth, 1901): Venezuela: 4; Total **4**
quadrinotata (Boheman, 1862): Costa Rica: 3; Total **3**
ruforeticulata Borowiec, 2007 f: 478: Peru: 2 (**1 HT, 1 PT**); Total **2**
serinus (Erichson, 1847): Bolivia: 3, Brazil: 4, Ecuador: 4, Peru: 7; Total **18**
sexpustulata Fabricius, 1781): Brazil: 24; Total **24**
similata (Boheman, 1850): Brazil: 3; Total **3**
smaragdina (Boheman, 1856): Peru: 4, Ecuador: 1; Total **5**
steinheili (Wagener, 1877): Colombia: 4; Total **4**
textilis (Boheman, 1850): Colombia: 7; Total **7**
thalassina (Boheman, 1850): Brazil: 12; Total **12**
tigrina (Boheman, 1850): Argentina: 1; Total **1**
trigonata (Spaeth, 1901): Ecuador: 5; Total **5**
tristigma (Boheman, 1850): Guatemala: 14, Mexico: 10; Total **24**
verrucosa (Boheman, 1856): Brazil: 3; Total **3**
vulnerata (Boheman, 1850): Brazil: 6; Total **6**
zikani (Spaeth, 1932): Brazil: 2; Total **2**
Cyrtonota sp. nov.: Bolivia: 1; Total **1**
Cyrtonota sp.: no country: 1; Total **1**

Genus: *Echoma* Chevrolat, 1836

Subgenus *Echoma* s. str.:

anaglypta (Boheman, 1862): Colombia: 1; Total **1**
anaglyptoides Borowiec, 1998 e: 376: Colombia: 4 (**1 HT, 3 PT**), French Guyana: 3 (**1 PT**); Total **7**
biplagosa (Boheman, 1854): Brazil: 5; Total **5**
bonfilsii (Boheman, 1854): Ecuador: 10; Total **10**
clypeata (Panzer, 1798): Bolivia: 2, Brazil: 1, French Guyana: 32, Peru: 14; Total **49**
quadristillata (Boheman, 1862): Costa Rica: 3; Total **3**
signata (Panzer, 1798): Brazil: 1; Total **1**

Subgenus *Pseudomoplata* Spaeth, 1952

weyenberghi (Dohrn, 1878): Argentina: 8; Total **8**

Genus: *Elytrogona* Chevrolat, 1836

bacca Boheman, 1850: Dominican Rep.: 2; Total **2**
quatuordecimmaculata (Latreille, 1802): Dominican Rep.: 8; Total **8**

Genus: *Eutheria* Spaeth, 1909

piperata (Burmeister, 1870): Uruguay: 4; Total **4**

Genus: *Hilarocassis* Spaeth, 1913

albida (Germar, 1824): Brazil: 5, Paraguay: 1; Total **6**
bordoni Borowiec, 2002 a: 128: Venezuela: 4 (**2 PT**); Total **4**
evanida (Boheman, 1850): Argentina: 1, Bolivia: 3, Brazil: 1; Total **5**
exclamationis (Linnaeus, 1767): Costa Rica: 1, Mexico: 3, Venezuela: 6; Total **10**

maculicollis Świętojańska, 2003 b: Brazil: 2 (**1 HT, 1 PT**); Total **2**
quinquelineata Spaeth, 1913: Brazil: 4; Total **4**
suturella (Boheman, 1850): Brazil: 1; Total **1**
venusta (Boheman, 1854): Brazil: 1, Paraguay: 1; Total **2**

Genus: *Mesomphalia* Hope, 1839

denudata Boheman, 1850: Brazil: 2; Total **2**
gibbosa (Fabricius, 1781): Brazil: 11; Total **11**
latipennis Boheman, 1856: Brazil: 3; Total **3**
retipennis Boheman, 1850: Brazil: 6; Total **6**
sexmaculosa Boheman, 1850: Brazil: 1; Total **1**
sublaevis Boheman, 1850: Brazil: 2; Total **2**
turrita (Illiger, 1801): Brazil: 12; Total **12**
variolaris Boheman, 1850: Brazil: 4; Total **4**
Mesomphalia sp. nov.: Brazil: 4; Total **4**

Genus: *Nebraspis* Spaeth, 1913

corticina (Boheman, 1850): Brazil: 1, Paraguay: 1; Total **2**
rubricollis (Boheman, 1856): Brazil: 2; Total **2**
viridimetallica Borowiec, 1999 i: 427: Argentina: 5 (**1 HT, 4 PT**); Total **5**

Genus: *Ogdoecosta* Spaeth, 1909

biannularis (Boheman, 1854): Mexico: 10; Total **10**
catenulata (Boheman, 1854): Guatemala: 4, Mexico: 6; Total **10**
decemstillata (Boheman, 1856): Mexico: 2; Total **2**
epilachnoides (Champion, 1893): Mexico: 13; Total **13**
fasciata (Boheman, 1856): Mexico: 5; Total **5**
guttifera (Boheman, 1854): Guatemala: 5, Mexico: 1; Total **6**
juvenca (Boheman, 1854): Costa Rica: 1, Mexico: 3; Total **4**
mexicana (Champion, 1893): Mexico: 2; Total **2**
obliterata (Champion, 1883): Mexico: 2; Total **2**
Ogdoecosta sp.: Mexico: 1; Total **1**

Genus: *Omaspides* Chevrolat, 1836

Subgenus *Omaspides* s. str. (34):

abbreviata Baly, 1872: Ecuador: 1, Peru: 6 (**2 PT** of *Omaspides andicola* Spaeth, 1909: 386); Total **7**
augusta Boheman, 1856: Peru: 12; Total **12**
basilica Boheman, 1856: Brazil: 4; Total **4**
bioculata Boheman, 1862: Brazil: 1; Total **1**
bistriata Boheman, 1862: Costa Rica: 1, Panama: 1; Total **2**
boliviana Borowiec, 2003 h: 397: Bolivia: 1 (**1 PT**); Total **1**
clathrata (Linnaeus, 1758): French Guyana: 15; Total **15**
confusa Borowiec, 2010 f: 483: Ecuador: 9 (**1 HT, 8 PT**); Total **9**
conjugens Spaeth, 1937: Peru: 1; Total **1**
convexicollis Spaeth, 1909: Costa Rica: 1; Total **1**
erichsoni Spaeth, 1937: Bolivia: 5, Peru: 1; Total **6**
flavofasciata Spaeth, 1909: 388: Peru: 2 (**1 ST**); Total **2**
iheringi (Spaeth, 1909): Brazil: 2, Peru: 1; Total **3**
lunulata Boheman, 1854: Brazil: 2; Total **2**
nigrolineata (Boheman, 1854): Brazil: 1; Total **1**
nitidicollis Spaeth, 1937: Colombia: 5; Total **5**

pallidipennis (Boheman, 1854): Brazil: 12; Total **12**
picaflorensis Borowiec, 2010 f: 484: Peru: 5 (**1 HT, 4 PT**); Total **5**
planicollis Sekerka & Windsor in litt.: Amerique du Sud: 1 (**1 PT**), Ecuador: 2 (**2 PT**), Panama: 1 (**1 PT**), Peru: 1 (**1 PT**); Total **5**
pulchella Baly, 1859: Brazil: 5, Peru: 1; Total **6**
quadrifenestrata Spaeth, 1907: Argentina: 2, Peru: 2; Total **4**
scutaria Spaeth, 1937: Brazil: 2; Total **2**
specularis Erichson, 1847: Peru: 3; Total **3**
trichroa (Boheman, 1854): Bolivia: 4, Brazil: 15; Total **19**
tricolorata (Boheman, 1854): Brazil: 3; Total **3**
trifasciata (Fabricius, 1787): French Guyana: 1, Surinam: 1; Total **2**
unicolor Borowiec, 1998 d: 218: Ecuador: 1 (**1 HT**); Total **1**
Omaspides sp. 1: Peru: 1; Total **1**
Omaspides sp. 2: Peru: 1; Total **1**
Omaspides sp. 3: Peru: 1; Total **1**
Omaspides sp. 4: Brazil: 1; Total **1**
Omaspides sp. 5: Peru: 1; Total **1**
Omaspides sp. 6: Brazil: 2; Total **2**
Subgenus *Parechoma* Spaeth, 1913
semilineata (Boheman, 1854): Brazil: 4; Total **4**
Subgenus *Paromaspides* Spaeth, 1913
brunneosignata (Boheman, 1854): Brazil: 4; Total **4**
haematidea (Boheman, 1854): Brazil: 2; Total **2**
sobrina (Boheman, 1854): Brazil: 9; Total **9**
squalida (Boheman, 1854): Brazil: 6; Total **6**
vexabilis (Boheman, 1862): Brazil: 2; Total **2**

Genus: *Paraselenis* Spaeth, 1913
Subgenus *Paraselenis* s. str.
albida Spaeth, 1913: Brazil: 3; Total **3**
axillaris (Sahlberg, 1823): Brazil: 10; Total **10**
Subgenus *Pseudechoma* Spaeth, 1913
marginipennis (Spaeth, 1907): Ecuador: 3, Peru: 4; Total **7**
Subgenus *Spaethiechoma* Hincks, 1950:
amplicollis (Spaeth, 1907): Argentina: 5, Bolivia: 1, Brazil: 1, Paraguay: 1; Total **8**
aulica (Boheman, 1854): Brazil: 6; Total **6**
brunnidorsis (Spaeth, 1902): Bolivia: 4, Peru: 1; Total **5**
collata (Boheman, 1854): Brazil: 4; Total **4**
convolvuli Villamil, Borowiec & Świętojańska, 2023: 702: Argentina: 10 (**1 HT, 9 PT**); Total **10**
decipiens (Boheman, 1854): Brazil: 8; Total **8**
dichroa (Germar, 1824): Brazil: 19; Total **19**
filia (Spaeth, 1907): Paraguay: 17; Total **17**
flava (Linnaeus, 1758): Paraguay: 9; Total **9**
flavata (Boheman, 1854): Brazil: 3; Total **3**
flavopunctata Borowiec, 2003 i: 407: Argentina: 1 (**1 PT**), Brazil: 1 (**1 HT**); Total **2**
generosa (Boheman, 1854): Brazil: 2; Total **2**
hyalina (Boheman, 1854): Argentina: 8, Bolivia: 1; Total **9**
jugata (Boheman, 1854): Brazil: 1; Total **1**
nigropunctata Borowiec, 2003 i: 404: Bolivia: 8 (**8 PT**); Total **8**
normalis (Germar, 1824): Brazil: 16; Total **16**
nupta (Boheman, 1854): Brazil: 7; Total **7**
puncticollis (Spaeth, 1907): Bolivia: 4, Brazil: 1, Peru: 3; Total **8**

saltaensis Borowiec, 2002 a: 134: Argentina: 47 (**47 PT**); Total **47**
scapulosa (Boheman, 1854): Argentina: 2, Brazil: 1, Paraguay: 4; Total **7**
solieri (Boheman, 1854): Franch Guyana: 2, Guyana: 4, Surinam: 7, Brazil: 1; Total **14**
suspecta (Spaeth, 1907): Uruguay: 2; Total **2**
tersa (Boheman, 1854): Panama: 4, Peru: 1; Total **5**
transversalis (Boheman, 1854): Brazil: 1; Total **1**
Paraselenis sp. 1: Ecuador: 1; Total **1**
Paraselenis sp. 2: Brazil: 1; Total **1**
Paraselenis sp. 3: Brazil: 1; Total **1**

Genus: *Phytodectoidea* Spaeth, 1909

duodecimsignata (Boheman, 1854): Mexico: 11; Total **11**
quatuordecimpunctata (Boheman, 1854): Mexico: 11; Total **11**

Genus: *Poecilaspidella* Spaeth, 1913

bruchi (Spaeth, 1909): Paraguay: 2; Total **2**
duplicata (Boheman, 1850): Brazil: 1; Total **1**
histrion (Boheman, 1850): Brazil: 4, Paraguay: 1; Total **5**
proxima (Boheman, 1850): Argentina: 7, Brazil: 1, Uruguay: 1; Total **9**
vittifera (Boheman, 1850): Uruguay: 6; Total **6**

Genus: *Stoiba* Spaeth, 1909

angusticollis (Suffrian, 1868): Cuba: 1; Total **1**
flavicollis (Klug, 1829): Cuba: 8; Total **8**
indivisa Blake, 1930: Cuba: 1; Total **1**
swartzii (Thunberg, 1817): Jamaica: 2; Total **2**

Genus: *Stolas* Billberg, 1820

acuminata (Boheman, 1850): Brazil: 1; Total **1**
acuta (Boheman, 1850): Brazil: 1; Total **1**
aenea (Olivier, 1790): Brazil: 13; Total **13**
aeneovittata (Champion, 1894): Panama: 2; Total **2**
agenysiformis Borowiec, 2011 c: 486: Brazil: 1 (**1 PT**); Total **1**
amoena (Spaeth, 1917): Ecuador: 2; Total **2**
anceps (Boheman, 1850): Brazil: 3; Total **3**
antiqua (Sahlberg, 1823): Brazil: 18; Total **18**
areolata (Germar, 1824): Brazil: 7; Total **7**
armirantensis Borowiec, 1999 i: 432: Peru: 1 (**1 HT**); Total **1**
asema (Boheman, 1862): Brazil: 1; Total **1**
atalayaensis Borowiec & Takizawa, 2011 e: 477: Ecuador: 2 (**1 PT**); Total **2**
augur (Boheman, 1856): Brazil: 1; Total **1**
bilineata (Boheman, 1850): Brazil: 1; Total **1**
bioculata (Boheman, 1850): Bolivia: 2, Peru: 2; Total **4**
blanda (Boheman, 1850): Ecuador: 1; Total **1**
boliviana (Spaeth, 1909): Bolivia: 2; Total **2**
brachiata (Fabricius, 1798): Brazil: 5, French Guyana: 6, Paraguay: 1; Total **12**
brevicuspis (Spaeth, 1922): Brazil: 2; Total **2**
callizona (Spaeth, 1909: 372): Peru: 1 (**1 ST**); Total **1**
cassandra (Spaeth, 1911): Peru: 2; Total **2**
castigata (Boheman, 1856): Bolivia: 7; Total **7**
chalybaea (Germar, 1824): Argentina: 5, Brazil: 20, Paraguay: 4; Total **29**
coalita (Boheman, 1862): Ecuador: 24; Total **24**

coerulescens (Boheman, 1850): Bolivia: 10, Brazil: 1; Total **11**
comis (Spaeth, 1932): Peru: 3; Total **3**
confusa (Boheman, 1850): French Guyana: 4; Total **4**
consanguinea (Kirsch, 1876): Peru: 6; Total **6**
consociata (Baly, 1872): Ecuador: 2; Total **2**
conspersa (Germar, 1824): Brazil: 21; Total **21**
contubernalis (Boheman, 1856): Brazil: 1; Total **1**
croceovittata (Spaeth, 1901): Bolivia: 3; Total **3**
cruentata (Erichson, 1847): Bolivia: 26, Peru: 6; Total **32**
cucullata (Boheman, 1862): Costa Rica: 5; Total **5**
decemguttata (Sturm, 1828): Brazil: 16; Total **16**
deleta (Boheman, 1850): Brazil: 12; Total **12**
demissa (Boheman, 1856): Brazil: 1; Total **1**
discooides (Linnaeus, 1758): Brazil: 8, Colombia: 1, Ecuador: 7, French Guyana: 10, Peru: 8, Trinidad: 2; Total **36**
diversa (Boheman, 1850): Brazil: 2; Total **2**
echoma Borowiec, 1998 b: 221: Ecuador: 1 (**1 HT**); Total **1**
ephippium (Lichtenstein, 1795): Colombia: 5, Ecuador: 2, Nicaragua: 1, Panama: 2; Total **10**
erectepilosa Borowiec, 1998 d: 222: Ecuador: 1 (**1 HT**); Total **1**
erichsoni (Weise, 1902): Bolivia: 6; Total **6**
eugenea (Boheman, 1862): Peru: 2; Total **2**
excelsa (Spaeth, 1917): Ecuador: 2; Total **2**
extricata (Boheman, 1850): Colombia: 2, Panama: 1; Total **3**
famula (Spaeth, 1928): Bolivia: 1; Total **2**
fasciculata (Boheman, 1850): Brazil: 1; Total **1**
festiva (Klug, 1829): Argentina: 9, Brazil: 13, Paraguay: 1, Uruguay: 1; Total **24**
flavipennis (Boheman, 1850): Peru: 1; Total **1**
flavofenestrata (Boheman, 1862): Venezuela: 1; Total **1**
flavoguttata (Boheman, 1850): Brazil: 4; Total **4**
flavomarginata Borowiec, 1999 i: 427: Peru: 1 (**1 HT**), Brazil: 1; Total **2**
flavonotata (Boheman, 1856): Brazil: 1; Total **1**
flavoreticulata (Boheman, 1856): Bolivia: 2, Peru: 1; Total **3**
floccosa (Erichson, 1847): Bolivia: 5, Peru: 1; Total **6**
funebri (Boheman, 1850): Bolivia: 2, Peru: 12; Total **14**
godeti (Boheman, 1850): Brazil: 1; Total **1**
haematites (Lichtenstein, 1795): Colombia: 1, Ecuador: 1, Panama: 2; Total **4**
hameli Borowiec, 2007 j: 662: Bolivia: 1 (**1 PT**); Total **1**
hermanni (Spaeth, 1911): Argentina: 1; Total **1**
huanocensis (Spaeth, 1901): Bolivia: 3, Peru: 2; Total **5**
hypocrita (Boheman, 1862): Ecuador: 3; Total **3**
ignita (Boheman, 1850): Brazil: 7; Total **7**
iheringi (Spaeth, 1913): Brazil: 1; Total **1**
illustris (Chevrolat, 1834): Guatemala: 2, Mexico: 4; Total **6**
imitatrix Borowiec, 1999 i: 430: Peru: 1 (**1 HT**); Total **1**
imparilis (Boheman, (1862): Brazil: 3; Total **3**
imperialis (Spaeth, 1898): Brazil: 6; Total **6**
implexa (Boheman, 1850): Brazil: 11; Total **11**
impluviata (Boheman, 1850): Brazil: 3; Total **3**
inaequalis (Linnaeus, 1758): Brazil: 22; Total **22**
inaurata (Burmeister, 1870): Argentina: 6, Bolivia: 2; Total **8**
inca (Spaeth, 1901): Peru: 4; Total **4**
indigacea (Boheman, 1850): Argentina: 12, Bolivia: 2; Total **14**
inexculata (Boheman, 1862): Ecuador: 20; Total **20**

informis (Boheman, 1862): Ecuador: 1; Total **1**
intermedia Borowiec, 1999 i: 434: Peru: 4 (**1 HT**); Total **4**
isthmica (Champion, 1893): Colombia: 1, Costa Rica: 16; Total **17**
kraatzi (Boheman, 1862): Peru: 2; Total **2**
lacordairei (Boheman, 1850): Argentina: 14, Brazil: 9, no country: 4; Total **27**
lacunosa (Boheman, 1850): Brazil: 3; Total **3**
latevittata (Boheman, 1862): Ecuador: 3; Total **3**
lebasii (Boheman, 1850): Costa Rica: 4, Guatemala: 5, Mexico: 3, Panama: 1; Total **13**
lenis (Boheman, 1850): Brazil: 5; Total **5**
lineaticollis (Boheman, 1850): Brazil: 9; Total **9**
mannerheimi (Boheman, 1850): Peru: 11; Total **11**
modica (Boheman, 1850): Brazil: 7; Total **7**
napoensis Borowiec, 1998 d: 225: Colombia: 1 (**1 PT**), Ecuador: 5 (**4 PT**); Total **6**
nigrolineata (Champion, 1893): Costa Rica: 5; Total **5**
niobe (Spaeth, 1919): Ecuador: 1; Total **1**
nudicollis (Boheman, 1850): Brazil: 10; Total **10**
oblita (Boheman, 1850): Brazil: 9; Total **9**
obvolvata (Boheman, 1862): Bolivia: 1, Paru: 3; Total **4**
octosignata (Spaeth, 1913): Brazil: 12; Total **12**
omaspidiformis Borowiec, 2007 j: 665: Peru: 1 (**1 HT**); Total **1**
paranensis (Spaeth, 1928): Brazil: 10; Total **10**
pascoei (Baly, 1872): Ecuador: 2; Total **2**
paulista (Spaeth, 1935): Brazil: 2; Total **2**
pectinata (Baly, 1872): Ecuador: 1; Total **1**
pellicula (Spaeth, 1915): Ecuador: 12; Total **12**
perezi Borowiec, 1998 d: 227: Ecuador: 2 (**1 PT**); Total **2**
perfuga (Spaeth, 1926): Brazil: 1; Total **1**
perjucunda (Baly, 1872): Ecuador: 6; Total **6**
pertusa (Boheman, 1850): Argentina: 2, Bolivia: 1, French Guyana: 3, Brazil: 4, Colombia: 1, Ecuador: 1, Paraguay: 1, no country: 3, Venezuela: 3; Total **19**
placida (Spaeth, 1911): Peru: 3; Total **3**
plagicollis (Boheman, 1850): Brazil: 13; Total **13**
pleurosticha (Erichson, 1847): Bolivia: 5, Ecuador: 1, Peru: 8; Total **14**
praecalva (Spaeth, 1942): Peru: 1; Total **1**
praetoria (Spaeth, 1928): Ecuador: 1; Total **1**
puberula (Boheman, 1856): Colombia: 1, Ecuador: 11; Total **12**
pucallpaensis Borowiec & Takizawa, 2011 e: 480: Peru: 1 (**1 PT**); Total **1**
punicea (Boheman, 1850): Guatemala: 3, Mexico: 5; Total **8**
quatuordecimsignata (Boheman, 1850): Bolivia: 2; Total **2**
quinquefasciata (Wagner, 1877): Peru: 1; Total **1**
redtenbacheri (Boheman, 1850): Argentina: 1, Brazil: 10, Paraguay: 5; Total **16**
reticularis (Linnaeus, 1758): French Guyana: 10; Total **10**
rubroreticulata (Boheman, 1856): Colombia: 4; Total **4**
sanguineovittata Borowiec, 1998 d: 228: Ecuador: 2 (**1 HT**); Total **2**
sanramonensis Borowiec, 2005 e: Bolivia: 1 (**1 PT**); Total **1**
saundersi (Boheman, 1856): Ecuador: 7; Total **7**
schaumi (Boheman, 1850): French Guyana: 3; Total **3**
scoparia (Erichson, 1847): Bolivia: 11, Peru: 3; Total **14**
selecta (Spaeth, 1928): Brazil: 3; Total **3**
sexplagiata (Boheman, 1850): Brazil: 3; Total **3**
sexsignata (Boheman, 1850): Brazil: 7; Total **7**
similis Sekerka & Windsor in litt.: Panama: 1 (**1 PT**); Total **1**

sommeri (Boheman, 1850): Brazil: 5; Total **5**
stevensi (Boheman, 1856): Brazil: 1; Total **1**
stolida (Spaeth, 1917): Colombia: 2, Ecuador: 7 (**1 HT, 3 PT** *Stolas stolida* ssp. *jadwiszczaki* Borowiec, 1998 d: 231); Total **9**
stygia (Spaeth, 1932): Bolivia: 4; Total **4**
submetallica (Weise, 1902): Peru: 3; Total **3**
subreticulata (Boheman, 1850): Brazil: 8; Total **8**
suturalis (Fabricius, 1777): Brazil: 6; Total **6**
tachiraensis Borowiec, 2009 i: 715: Colombia: 2, Venezuela: 9 (**1 HT, 8 PT**); Total **11**
thoreyi (Boheman, 1862): Brazil: 4; Total **4**
tumulus (Boheman, 1850): Brazil: 2; Total **2**
uniformis Borowiec & Świętojańska, 2010 e: Peru: 1 (**1 HT**); Total **1**
verecunda (Boheman, 1850): Bolivia: 2, Brazil: 1, Peru: 2; Total **5**
vetula (Boheman, 1850): Brazil: 1; Total **1**
vicina (Boheman, 1850): Brazil: 4; Total **4**
vorax (Weise, 1902): Peru: 1; Total **1**
warchalowskii Borowiec, 2007 j: 667: Paraguay: 1 (**1 HT**); Total **1**
zonata (Spaeth, 1902): Peru: 13; Total **13**
zumbaensis Borowiec, 1998 d: 232: Ecuador: 2 (**1 HT**); Total **2**
Stolas sp. 1: French Guyana: 1; Total **1**
Stolas sp. 2: Ecuador: 2; Total **2**
Stolas indet.: **27**

Genus: *Terpsis* Spaeth, 1913

quadrivittata (Champion, 1893): Costa Rica: 2, Panama: 3; Total **5**

Genus: *Trilaccodea* Spaeth, 1902

ecuadorica Borowiec, 2007 a: Ecuador: 1 (**1 HT**); Total **1**
langei (Spaeth, 1902): Peru: 2; Total **2**
meridionalis Borowiec, 2004 a: Bolivia: 1 (**1 HT**); Total **1**
schneideri (Spaeth, 1909): Peru: 1; Total **1**
tomentosa (Boheman, 1850): Venezuela: 14; Total **14**

Genus: *Xenicomorpha* Spaeth, 1913

scapularis (Boheman, 1854): Bolivia: 2; Total **2**

Genus: *Zatrephina* Spaeth, 1909

atrорubra (Guérin, 1844): Argentina: 3, Bolivia: 6; Total **9**
lineata (Fabricius, 1787): Argentina: 5, Brazil: 10, Venezuela: 3; Total **18**
princeps (Boheman, 1854): Argentina: 7, French Guyana (mislabeled?): 1; Total **8**
sexlunata (Klug, 1829): Argentina: 5, Brazil: 1, Paraguay: 7; Total **13**
Zatrephina sp. 1: Paraguay: 3; Total **3**

Tribe Notosacanthini

Genus: *Herminella* Spaeth, 1913

flavocostata Borowiec & Świętojańska, 2014 d: 259: Namibia: 1 (**1 HT**); RSA: 1 (**1 PT**); Total **2**
liliputana Spaeth, 1926: RSA: 1; Total **1**
marshalli Spaeth, 1913: RSA: 1; Total **1**
quadrimaculata Borowiec & Świętojańska, 2014 d: 266: RSA: 1 (**1 HT**); Total **1**

Genus: *Hermosacantha* Borowiec & Świętojańska, 2014

madagascari Borowiec & Świętojańska, 2014 d: 270: Madagascar: 1 (**1 HT**); Total **1**

Genus: *Notosacantha* Chevrolat, 1837

alberti (Spaeth, 1913): Zambia: 14; Total **14**

alluaudi (Spaeth, 1913): Madagascar: 1; Total **1**

ambrica (Spaeth, 1913): Madagascar: 1; Total **1**

appropinquans (Spaeth, 1913): Malaysia: 1; Total **1**

armigera (Olivier, 1808): Madagascar: 2; Total **2**

badia (Boheman, 1850): RSA: 7; Total **7**

bakeri (Weise, 1915): Philippines: 2; Total **2**

balyi (Spaeth, 1913): Madagascar: 6; Total **6**

banaszkiewiczae Świętojańska & Stach, 2011: 421: Vietnam: 1 (**1 HT**); Total **1**

benguetina (Spaeth, 1921): Philippines: 5; Total **5**

bertounesquei Borowiec & Le Tirant, 2013 f: 289: India: 1 (**1 HT**); Total **1**

bezdeki Świętojańska, 2002 a: 39: Laos (**1 HT**): 1; Total **1**

biaculeata (Boheman, 1856): Madagascar: 1; Total **1**

bifenestrella (Boheman, 1856): Sri Lanka: 1; Total **1**

bioculata (Wagener, 1877): Indonesia: 5; Total **5**

bipunctata (Spaeth, 1907): Philippines: 2; Total **2**

birmanica (Spaeth, 1913): Laos: 4, Thailand: 2; Total **6**

biroi (Spaeth, 1903): Papua N.G.: 3; Total **3**

blanda (Spaeth, 1932): Indonesia: 1; Total **1**

bohemani (Weise, 1910): Madagascar: 1; Total **1**

brevicollis (Spaeth, 1913): Madagascar: 1; Total **1**

buergersi (Spaeth, 1925): Papua N.G.: 3; Total **3**

butuana (Spaeth, 1925): Philippines: 2; Total **2**

calligera (Spaeth, 1921): Malaysia (Borneo): 2; Total **2**

capitata (Spaeth, 1913): Madagascar: 1; Total **1**

castanea (Spaeth, 1913): India: 2, Taiwan: 1; Total **3**

cavillata (Spaeth, 1933): Philippines: 3; Total **3**

centinodia (Spaeth, 1913): Laos: 6; Total **6**

chandrapurensis Świętojańska, Ghate & Marathe, 2001: 489: India: 1 (**1 HT**); Total **1**

cincta (Spaeth, 1925): Papua N.G.: 2; Total **2**

circumdata (Wagener, 1881): Indonesia: 2, Malaysia (Borneo): 1; Total **3**

circumminiata (Spaeth, 1913): Madagascar: 1; Total **1**

collarti (Spaeth, 1931: 103): Dem. Rep. of Congo: 2 (**1 ST**); Total **2**

corneola (Spaeth, 1913): India: 1; Total **1**

dammaropsi Sekerka, 2009: Papua N.G.: 1; Total **1**

dapitana (Spaeth, 1921): Philippines: 3; Total **3**

darjeelingensis Borowiec & Takizawa, 1991 d: 646: India: 1 (**1 HT**); Total **1**

decorsei (Spaeth, 1913): Madagascar: 1; Total **1**

delicatula (Spaeth, 1925): Philippines: 1; Total **1**

delutata (Spaeth, 1931): Zambia: 1; Total **1**

diabolica Świętojańska, 2001 b: 341: Malaysia (Borneo): 3 (**1 HT, 2 PT**); Total **3**

dohrni Świętojańska, 2001 a: 54: Indonesia: 2 (**2 PT**); Total **2**

donckieri (Spaeth, 1913): Madagascar: 1; Total **1**

dorsalis (Waterhouse, 1877): Australia: 1; Total **1**

echinata (Fabricius, 1801): Madagascar: 2; Total **2**

elisabethae (Spaeth, 1913): Dem. Rep. of Congo: 2, Zambia: 2; Total **4**

excelsa (Spaeth, 1940): Malaysia (Borneo): 1; Total **1**

fabricii (Weise, 1910): Madagascar: 1; Total **1**

filiola (Spaeth, 1913): RSA: 1; Total **1**
flavicornis (Spaeth, 1913): Indonesia: 1; Total **1**
flaviventris Borowiec & Takizawa, 2013 in Borowiec, Takizawa & Świętojańska, 2013: 162: Indonesia: 1 (**1 PT**);
 Total **1**
gemmata (Klug, 1833): Madagascar: 7; Total **7**
goudoti (Weise, 1910): Madagascar: 7; Total **7**
guerini (Weise, 1910): Madagascar: 1; Total **1**
gutticollis (Spaeth, 1932): Madagascar: 1; Total **1**
halmaherana Świętojańska, 2006: 364: Indonesia: 1 (**1 HT**); Total **1**
hebe (Spaeth, 1913): Madagascar: 3; Total **3**
hispoides (Spaeth, 1913): Madagascar: 6; Total **6**
holubi (Spaeth, 1905): Dem. Rep. of Congo: 1, Zimbabwe: 1; Total **2**
horni (Spaeth, 1914): Sri Lanka: 2; Total **2**
horrida (Boheman, 1850): Madagascar: 2; Total **2**
horrifica (Boheman, 1856): India: 2; Total **2**
hystrix (Spaeth, 1913): Madagascar: 4; Total **4**
ihai Chûjô, 1958: Japan: 1; Total **1**
impicta (Spaeth, 1925): Philippines: 6; Total **6**
jacobsoni (Spaeth, 1926): Indonesia: 4; Total **4**
jammuensis Borowiec & Takizawa, 1991 d: 647: India: 1 (**1 HT**); Total **1**
junodi (Spaeth, 1913): RSA: 2, Zimbabwe: 1; Total **3**
kantneri Świętojańska & Borowiec, 1999 h: 422: Thailand: 1 (**1 HT**); Total **1**
kinabaluensis Świętojańska, 2001 b: 343: Malaysia: 1 (**1 HT**); Total **1**
kohlschuetteri (Weise, 1903): Dem. Rep. of Congo: 6, Zimbabwe: 1; Total **7**
komiya Dąbrowska & Borowiec, 1996 c: 452: Laos: 1, Malaysia: 2, Thailand: 1 (**1 HT**); Total **4**
laticollis (Boheman, 1862): RSA: 6; Total **6**
lenta (Spaeth, 1913 b: 519): India: 5 (**1 ST**); Total **5**
leplaei (Spaeth, 1913): Dem. Rep. of Congo: 1; Total **1**
loochooana Chûjô, 1961: Japan: 2; Total **2**
luctuosa (Boheman, 1850): Madagascar: 1; Total **1**
lumbwa (Spaeth, 1924): Zimbabwe: 1; Total **1**
luteocincta (Spaeth, 1913): Madagascar: 1; Total **1**
maculipennis (Boheman, 1856): India: 3, Nepal: 4; Total **7**
maeander (Spaeth, 1925): Papua N.G.: 11; Total **11**
malaysiana Świętojańska, 2004: 242: Indonesia: 1, Malaysia: 1 (**1 HT**); Total **2**
marginata (Boheman, 1850): Madagascar: 1; Total **1**
matsuzawai Świętojańska, 2001 a: 51: Indonesia: 1 (**1 PT**); Total **1**
mentawaiensis Świętojańska & Stach, 2011: 423: Indonesia: 1 (**1 HT**); Total **1**
minutissima Świętojańska, 2000 a: 210: Philippines: 2 (**1 HT, 1 PT**); Total **2**
mocquerisi (Spaeth, 1913): Madagascar: 1; Total **1**
moderata Chen & Zia, 1964: Thailand: 4; Total **4**
molucana (Boheman, 1850): Australia: 1, Indonesia: 2, Papua N.G.: 6; Total **9**
monilicornis (Spaeth, 1913): Australia: 1; Total **1**
moultoni (Spaeth, 1912): Malaysia (Borneo): 1; Total **1**
multicostata Świętojańska, 2000 b: 593: Madagascar: 1 (**1 HT**); Total **1**
multimaculata Świętojańska, 2000 a: 214: Philippines: 2 (**2 PT**); Total **2**
myanmarensis Świętojańska, 2004: 244: Myanmar: 1 (**1 HT**); Total **1**
nathani Borowiec & Takizawa, 1991: 649: India: 4 (**1 HT**); Total **4**
negrosia (Spaeth, 1925): Philippines: 5; Total **5**
nervosa (Spaeth, 1913): Dem. Rep. of Congo: 1, Zambia: 1; Total **2**
nickerli (Spaeth, 1905: 79): Zimbabwe: 3 (**1 ST**); Total **3**
nigra (Duvivier, 1891): Madagascar: 1; Total **1**

nigriclavis (Spaeth, 1913): Madagascar: 1; Total **1**
nigritula (Spaeth, 1913): Madagascar: 1; Total **1**
nigrodorsata Chen & Zia, 1961: Laos: 1; Total **1**
obscura (Wagener, 1881): Indonesia: 2; Total **2**
octomaculata Świętojańska, 2000 a: 212: Philippines: 2 (**2 PT**); Total **2**
odiosa (Boheman, 1856): Indonesia: 1; Total **1**
ovoidea Borowiec & Świętojańska, 2013 in Borowiec, Takizawa & Świętojańska, 2013 a: 170: Malaysia (Borneo):
1 (**1 PT**); Total **1**
pakistanica Borowiec, Sultan & Rafi, 2008 in Borowiec, Sultan, Rafi, Naz, 2008 c: 288: Pakistan: 4 (**1 HT, 3 PT**);
Total **4**
paradoxa (Boheman, 1850): Madagascar: 3; Total **3**
parallela (Boheman, 1850): RSA: 2; Total **2**
parvula (Spaeth, 1925): Madagascar: 1; Total **1**
pauli (Weise, 1899): Tanzania: 1; unknown country: 1; Total **2**
pauliani Hincks, 1962: Madagascar: 1; Total **1**
pavida (Spaeth, 1913): Madagascar: 1; Total **1**
permira (Fairmaire, 1895): Madagascar: 2; Total **2**
perrieri (Spaeth, 1913): Madagascar: 1; Total **1**
pochoni (Spaeth, 1940): Madagascar: 2; Total **2**
pulchra (Spaeth, 1913): Madagascar: 11; Total **11**
quadra (Spaeth, 1915): Malaysia (Borneo): 2; Total **2**
quadricolorata Hincks, 1962: Madagascar: 1; Total **1**
quadrimaculata Świętojańska, 2000 a: 208: Philippines: 1 (**1 HT**); Total **1**
quadrituberculata (Guérin, 1844): Madagascar: 1; Total **1**
reinecki (Spaeth, 1913): Indonesia: 1; Total **1**
riedeli Świętojańska, 2002 b: 353: Indonesia: 1 (**1 HT**); Total **1**
roseola (Spaeth, 1915): Dem. Rep. of Congo: 1, Ghana: 1, Rep. of Central Africa: 2; Total **4**
2 PT); Total **3**
rufa (Wagener, 1881): Indonesia: 6, Malaysia: 2; Total **8**
sandakanensis Świętojańska, 2001 b: 338: Malaysia: 1 (**1 HT**); Total **1**
sarawaiensis Świętojańska, 2001 b: 335: Indonesia: 1 (**1 HT**); Total **1**
sauteri (Spaeth, 1914): China: 3, Laos: 2, Taiwan: 1; Total **6**
schoutedeni (Spaeth, 1913): Dem. Rep. of Congo: 6, Zambia: 1; Total **7**
semiviridis (Guérin, 1844): Madagascar: 5; Total **5**
severini (Spaeth, 1913): India: 3; Total **3**
shibatai Kimoto, 1981: Taiwan: 1; Total **1**
shishona Chen & Zia, 1964: India: 1, Laos: 1, Thailand: 1; Total **3**
sibuyana (Spaeth, 1925): Philippines: 1; Total **1**
socia (Spaeth, 1932): Madagascar: 1; Total **1**
spiculata (Spaeth, 1918): Madagascar: 1; Total **1**
striolata Hincks, 1962: Madagascar: 1; Total **1**
subparallela Hincks, 1962: 239: Madagascar: 1 (**1 PT**); Total **1**
sulawesica Borowiec, 1999 b: 8: Indonesia (**1 HT**): 1; Total **1**
sumatrana (Spaeth, 1940): Indonesia: 1; Total **1**
sumbawaensis Świętojańska & Borowiec, 1999 h: 424: Indonesia (**1 HT**): 1; Total **1**
taeniata (Fabricius, 1801): Philippines: 24; Total **24**
tenuicula (Spaeth, 1913): India: 2; Total **2**
tenuispina (Spaeth, 1913): Madagascar: 1; Total **1**
tessmanni (Spaeth, 1925): Dem. Rep. of Congo: 1, Total **1**
thiemei (Weise, 1910): Madagascar: 1; Total **1**
torrida (Spaeth, 1934): Madagascar: 1; Total **1**
vandepolli (Spaeth, 1913): Zambia: 1; Total **1**

undulata (Wagener, 1881): Philippines: 4; Total **4**
venus (Spaeth, 1926): Madagascar: 1; Total **1**
vicaria (Spaeth, 1913): India: 3, Sri Lanka: 2; Total **5**
vilis (Spaeth, 1913): Madagascar: 1; Total **1**
viridipennis Świętojańska, 2003 a: 393: Madagascar: 1 (**1 HT**); Total **1**
viridipicta (Boheman, 1862): Madagascar: 1; Total **1**
viriditincta (Boheman, 1856): Madagascar: 1; Total **1**
vittata (Wagener, 1881): Philippines: 2; Total **2**
vogeli (Weise, 1903): Dem. Rep. of Congo: 1, Tanzania: 1, Zambia: 1; Total **3**
warchalowskii Świętojańska & Sekerka, 2007: 681: India: 7 (**1 HT, 6 PT**); Total **7**
weyersi (Spaeth, 1900): Indonesia: 3; Total **3**
 sp. nov. 1_MADAG: Madagascar: 1; Total **1**
 sp. nov. 2_MADAG: Madagascar: 3; Total **3**
 sp. nov. 3_MADAG: Madagascar: 1; Total **1**
 sp. nov. 4_MADAG: Madagascar: 1; Total **1**
 sp. nov. 5_MADAG: Madagascar: 2; Total **2**
 sp. nov. 6_MADAG: Madagascar: 2; Total **2**
 sp. nov. 7_MADAG: Madagascar: 1; Total **1**
 sp. nov. 8_MADAG: Madagascar: 2; Total **2**
 sp. nov. 9_MADAG: Madagascar: 2; Total **2**
 sp. nov. 10_MADAG: Madagascar: 1; Total **1**
 sp. nov. 11_MADAG: Madagascar: 1; Total **1**
 sp. nov. 12_MADAG: Madagascar: 1; Total **1**
 sp. nov. 13_MADAG: Madagascar: 1; Total **1**
 sp. nov. 14_MADAG: Madagascar: 1; Total **1**
 sp. nov. 15_MADAG: Madagascar: 1; Total **1**
 sp. nov. 16_MADAG: Madagascar: 1; Total **1**
 sp. nov. 17_MADAG: Madagascar: 1; Total **1**
 sp. nov. 18_MADAG: Madagascar: 1; Total **1**
 sp. nov. 19_MADAG: Madagascar: 1; Total **1**
 sp. nov. 20_MADAG: Madagascar: 1; Total **1**
 sp. nov. 21_MADAG: Madagascar: 1; Total **1**
 sp. nov. 22_MADAG: Madagascar: 1; Total **1**
 sp. nov. 23_MADAG: Madagascar: 1; Total **1**
 sp. nov. 24_MADAG: Madagascar: 1; Total **1**
 sp. nov. 25_MADAG: Madagascar: 1; Total **1**
 sp. nov. 26_MADAG: Madagascar: 1; Total **1**
 sp. nov. 27_MADAG: Madagascar: 1; Total **1**
 sp. nov. 28_MADAG: Madagascar: 1; Total **1**
 sp. nov. 29_MADAG: Madagascar: 1; Total **1**
 sp. nov. 30_MALAWI: Madagascar: 1; Total **1**
 sp. nov. 31_ZAMBIA: Zambia: 1; Total **1**
 sp. nov. 32_UGANDA: Uganda: 1; Total **1**
 sp. nov. 33_ZAMBIA: Zambia: 1; Total **1**
 sp. nov. 34_RSA: RSA: 1; Total **1**
 sp. nov. 35_RWA+ZAI: Dem. Rep. of Congo: 1, Rwanda: 4; Total **5**
 sp. nov. 36_IND: India: 1; Total **1**
 sp. nov. 37_MAL: Malaysia (Borneo): 1; Total **1**

Tribe Omocerini

Genus: *Canistra* Erichson, 1847

Subgenus: *Canistra* s. str.

bitorosa Weise, 1900: Brazil: 1; Total **1**

dohrni Spaeth, 1905: Brazil: 13; Total **13**

irrorata (Guérin, 1844): Bolivia: 5, Paraguay: 1, Brazil: 2; Total **8**

osculatii Guérin, 1844: Colombia: 1, Ecuador: 5, no country: 1; Total **7**

procera Boheman, 1850: Bolivia: 2, Peru 4, no country: 1; Total **7**

varicosa Erichson, 1847: Peru: 1; Total **1**

sp. 1: Bolivia: 1; Total **1**

Subgenus: *Canistrella* Spaeth, 1913

calamitosa Boheman, 1850: Brazil: 1; Total **1**

rubiginosa (Guérin, 1844): Brazil: 28; Total **28**

Genus: *Carlobruchia* Spaeth, 1911

Subgenus: *Carlobruchia* s. str.

boliviana Borowiec, 1995 e: 423: Bolivia: 1 (**1 PT**); Total **1**

tricostata (Spaeth, 1907): Argentina: 6; Total **6**

Subgenus: *Smodingonota* Spaeth, 1913

carbonaria (Klug, 1829): Argentina: 1, Brazil: 5, Paraguay: 1; Total **7**

dorsuosa (Boheman, 1862): Brazil: 2; Total **2**

Genus: *Cassidinoma* Hincks, 1950

denticulata (Boheman, 1850): Paraguay: 6, no country: 1; Total **7**

Genus: *Cyclosoma* Guérin, 1835

Subgenus: *Cyclosoma* s. str.

anomala (Boheman, 1862): Ecuador: 3; Total **3**

palliatata (Fabricius, 1787): French Guyana: 19; Total **19**

tristis (Guérin, 1844): Bolivia: 3, Peru: 10; Total **13**

Subgenus: *Dolichotoma* Hope, 1839

fuscopunctata (Spaeth, 1919): Brazil: 4; Total **4**

germari (Spaeth, 1913): Brazil: 20; Total **20**

inepta (Boheman, 1850): Brazil: 3; Total **3**

lugens (Germar, 1824): Brazil: 2; Total **2**

spurca (Boheman, 1856): Brazil: 3; Total **3**

Subgenus: *Monrosiacassis* Viana, 1964

bicostata Borowiec, 1999 d: 82: Brazil: 6 (**5 PT**); Total **6**

nigritarsis (Boheman, 1856): Brazil: 1; Total **1**

puberula (Boheman, 1850): Venezuela: 1; Total **1**

sericata (Guérin, 1844): Bolivia: 5, Brazil: 15; Total **20**

strigata (Panzer, 1798): Brazil: 15; Total **15**

Subgenus: *Proglima* Hincks, 1950

gloriosa (Baly, 1859): Bolivia: 1, no country: 1; Total **2**

mirabilis (Boheman, 1856): Peru: 1; Total **1**

satanas (Weise, 1896): Brazil: 1; Total **1**

Genus: *Discomorpha* Chevrolat, 1836

Subgenus: *Discomorpha* s. str.

amazona (Spaeth, 1940): Brazil: 1; Total **1**

atahualpai Borowiec, 2006 j: 47: Peru: 4 (**1 HT, 1 PT**); Total **4**

batesi (Boheman, 1856): Brazil: 2; Total **2**
bernhaueri (Spaeth, 1909: 364): Colombia: 3 (**1 ST**); Total **3**
biplagiata (Guérin, 1844): Brazil: 1, Colombia: 17; Total **18**
bisbiplagiata (Boheman, 1856): Guatemala: 3; Total **3**
carlobrivioi Borowiec & Sassi, 2003 f: 231: Bolivia: 2 (**2 PT**); Total **2**
davidsoni Borowiec & Dąbrowska, 1996 d: 463: Brazil: 2 (**2 PT**); Total **2**
depilata (Boheman, 1850): Venezuela: 4; Total **4**
distincta (Baly, 1869): ?Brazil: 1 (probably mislabeled), Costa Rica: 2, Nicaragua: 2; Total **5**
dromedarius (Boheman, 1850): Brazil: 6; Total **6**
ducalis (Boheman, 1850): Brazil: 18; Total **18**
ganglbaueri (Spaeth, 1909: 366): Peru: 1 (**1 ST**); Total **1**
giganteensis Borowiec, 2006 j: 52: Colombia: 3 (**1 HT, 2 PT**); Total **3**
heikertingeri (Spaeth, 1920): Ecuador: 2; Total **2**
howdenorum Borowiec, 2006 j: 46: Venezuela: 1 (**1 PT**); Total **1**
instabilis (Baly, 1872): Colombia: 1, Ecuador: 3, Venezuela: 5, Paru: 2; Total **11**
languinosa (Boheman, 1850): Venezuela: 4; Total **4**
metallica (Guérin, 1844): Bolivia: 1, Brazil: 1; Total **2**
miniata (Boheman, 1850): Colombia: 4; Total **4**
mirifica (Boheman, 1856): Mexico (probably mislabeled): 1, Venezuela: 5; Total **6**
moczarskii (Spaeth, 1919): Brazil: 10; Total **10**
nigropunctata (Boheman, 1850): Colombia: 1, Ecuador: 1; Total **2**
nigrosparisa (Wagner, 1877): Colombia: 2; Total **2**
peruviana (Boheman, 1850): Ecuador: 5, Peru: 1; Total **6**
putamayoensis Borowiec, 2006 j: 50: Peru: 3 (**1 HT, 2 PT**); Total **3**
salvini (Baly, 1864): Panama: 4; Total **4**
santaremi Borowiec & Dąbrowska, 1996 d: 459: Brazil: 6 (**3 PT**); Total **6**
schusteri (Spaeth, 1907): Brazil: 11; Total **11**
variegata (Linnaeus, 1758): French Guyana: 12; Total **12**
waehneri (Spaeth, 1940): Argentina: 2, Colombia: 3, Peru: 2; Total **7**
wingelmuelleri (Spaeth, 1907): Colombia: 1; Total **1**
 sp. 1: Colombia: 2; Total **2**
 sp. 2: Peru: 3; Total **3**
Subgenus: *Paravulpia* Spaeth, 1940
sericea (Baly, 1872): Ecuador: 3; Total **3**
Subgenus: *Vulpia* Spaeth, 1913
fryi (Spaeth, 1928): Peru: 3; Total **3**
mandli (Spaeth, 1909): Brazil: 1; Total **1**
onorei Borowiec, 1998 d: 215: Ecuador: 5 (**5 PT**); Total **5**
secky Borowiec, 2001: Peru: 1; Total **1**
tigreensis Borowiec & Takizawa, 2011 e: 463: Peru: 1 (**1 PT**); Total **1**

Genus: *Omocerus* Chevrolat, 1835

Subgenus: *Omocerus* s. str.

arietinus (Boheman, 1862: 53): Brazil: 2 (**1 ST**), Surinam: 1; Total **3**
aureicornis (Blanchard, 1837): Brazil: 26; Total **26**
azureicornis (Chevrolat, 1835): Brazil: 1, Guatemala: 1, Mexico: 2; Total **4**
bicornis (Linnaeus, 1763): Brazil: 4, French Guyana: 6; Total **10**
clavareau (Spaeth, 1913): Ecuador: 4, no country: 1; Total **5**
ignava (Boheman, 1850): Brazil: 1; Total **1**
insculptus (Kirsch, 1876): Brazil: 4; Total **4**
masoni (Spaeth, 1926): Bolivia: 6; Total **6**
purpureus (Spaeth, 1912): Peru: 1; Total **1**

reichei (Boheman, 1850): Bolivia: 5; Total **5**
similis Borowiec, 2000 f: 219: Brazil: 16 (**16 PT**), no country: 1; Total **17**
taurus (Fabricius, 1787): Guyana: 1, Trinidad: 4, Venezuela: 1; Total **6**
tenebrosus Borowiec & Takizawa, 2011 e: 474: Peru: 1 (**1 PT**); Total **1**

Subgenus: *Nebroma* Spaeth, 1913

gibberus (Boheman, 1850): Brazil: 1; Total **1**
humerosus (Spaeth, 1913): Brazil: 1; Total **1**
klugi (Spaeth, 1913): Brazil: 5, Columbia (probably mislabeled): 1, Paraguay: 2; Total **8**
malachiticus (Germar, 1824): Brazil: 4; Total **4**
scabrosus (Boheman, 1850): Brazil: 8; Total **8**

Subgenus: *Paratauroma* Spaeth, 1913

angulicollis Borowiec, 2000 f: 223: Peru: 1 (**1 HT**); Total **1**
burakowskii Dąbrowska & Borowiec, 1995 d: 419: Brazil: 2 (**1 HT**); Total **2**
caeruleopunctatus (Boheman, 1862): Costa Rica: 1; Total **1**
casta (Boheman, 1862): Costa Rica: 2, Ecuador: 2; Total **4**
creberrimus (Boheman, 1850): Colombia: 6; Total **6**
doeberli Dąbrowska & Borowiec, 1995 d: 415: Brazil: 4 (**1 HT**); Total **4**
incertus Sekerka & Windsor in litt.: Panama: 1 (**1 PT**); Total **1**
janthinus (Boheman, 1862): Brazil: 1; Total **1**
reticulatus (Kirsch, 1876): Peru: 2; Total **2**
rugosicollis Borowiec, 2000 f: 225: Brazil: 1 (**1 HT**); Total **1**
smaragdinus (Boheman, 1850): Brazil: 25; Total **25**

Subgenus: *Platytauroma* Spaeth, 1913

anchoralis (Boheman, 1850): Brazil: 2; Total **2**
coracinus (Boheman, 1850): Brazil: 1; Total **1**
cornutus (Boheman, 1850): Brazil: 1; Total **1**
truncatus (Boheman, 1850): Brazil: 7; Total **7**

Genus: *Polychalca* Chevrolat, 1836

Subgenus: *Polychalca* s. str.

cariosa (Boheman, 1850): Brazil: 1; Total **1**
dentipennis (Boheman, 1850): Brazil: 4; Total **4**
nickerli (Spaeth, 1907): Brazil: 2; Total **2**
perforata (Boheman, 1850): Colombia: 3; Total **3**
punctatissima (Wolf, 1818): Brazil: 15; Total **15**

Subgenus: *Desmonota* Hope, 1840

aerea (Boheman, 1850): Brazil: 5; Total **5**
duponti (Boheman, 1850): Brazil: 4; Total **4**
gravida (Boheman, 1850): Bolivia: 2; Total **2**
platynota (Germar, 1824): Brazil: 11; Total **11**
salebrosa (Boheman, 1850): Bolivia: 8, Brazil: 1, Paraguay: 2; Total **11**

Tribe Spilophorini

Genus: *Calyptocephala* Chevrolat, 1836

attenuata Spaeth, 1919: Colombia: 1, Mexico: 1, Panama: 3; Total: **5**
brevicornis Boheman, 1850: Brazil: 4, Paraguay: 1; Total: **5**
discoidea Champion, 1893: Costa Rica: 1, Panama: 3; Total: **4**
gerstaeckeri Boheman, 1862: Belize: 1, Costa Rica: 1; Mexico: 2; Total: **4**
lutea Boheman, 1850: Brazil: 3, Surinam: 1; Total: **4**
marginipennis Boheman, 1850: Mexico: 3; Total: **3**

nigricornis (Germar, 1824): Brazil: 4; Total: **4**
punctata Boheman, 1850: Bolivia: 3, Brazil: 1, Surinam: 1; Total: **5**
ruficornis Spaeth, 1937: Brazil: 1; Total: **1**

Genus: *Cladispa* Baly, 1858

amboroensis Sekerka & Windsor in Sekerka, Windsor & Dury, 2014: 8: Bolivia: 1 (**1 PT**); Total **1**

Genus: *Spilophora* Boheman, 1850

aequatoriensis Spaeth, 1905: Ecuador: 5; Total **5**
bifasciata Spaeth, 1937: Brazil: 1, Ecuador: 3; Total **4**
cuneata Borowiec, 2003 j: 701: Ecuador: 3 (**1 HT, 2 PT**); Total **3**
lacrimata Borowiec, 2003 j: 702: Ecuador: 1 (**1 HT**), Peru: 1 (**1 PT**); Total **2**
nigriceps Spaeth, 1928: Ecuador: 1, Peru: 1; Total **2**
peruana (Erichson, 1847): Peru: 2; Total **2**
pulchra Boheman, 1856: Peru: 1; Total **1**
romani Weise, 1921: Brazil: 1; Total **1**
sellata Boheman, 1856: Brazil: 1; Total **1**
speciosa Baly, 1859: Brazil: 1, French Guyana: 1, Surinam: 1; Total **3**
tetraspilota Baly, 1859: Ecuador: 2; Total **2**
trigemina (Guérin, 1844): French Guyana: 1; Total **1**

Genus: *Spilophoroides* Sekerka in litt.

jucundus (Weise, 1910): Ecuador: 1, Panama: 1; Total **2**
marginatus (Weise, 1910): Panama: 1; Total **1**
panamensis Sekerka & Windsor in litt.: Panama: 2 (**2 PT**); Total **2**

Doublets and undetermined material: 243 ex.

Number of species: 2 491

Number of specimens: 21 292

Number of primary types: 229

Number of secondary types (paratypes + syntypes + paralectotypes): 904 + 71 + 4 = 979

Appendix

List of all taxa of Cassidinae described by Lech Borowiec

Genera and subgenera

1. *Afroaspidimorpha* Borowiec, 1997
2. *Aslamidium* Borowiec, 1984
3. *Aspidocassis* Borowiec, 1997
4. *Chapadacassis* Sekerka & Borowiec, 2015
5. *Hermosacantha* Borowiec & Świętojańska, 2014
6. *Neoaslamidium* Borowiec, 1998
7. *Neoaspidimorpha* Borowiec, 1992
8. *Paratorquispa* Borowiec, Świętojańska & Sekerka, 2019
9. *Pseudostilpnaspis* Borowiec, 2000
10. *Semiaspidimorpha* Borowiec, 1997
11. *Spaethiomorpha* Borowiec, 1997
12. *Wanatispa* Borowiec, Świętojańska & Sekerka, 2019

Species

1. *Acrocassis intermedia* Borowiec, 2002
2. *Acrocassis undulatipennis* Borowiec, 2002

3. *Aethiopocassis angulicollis* Borowiec & Świetojańska, 2018
4. *Aethiopocassis dewittei* Borowiec & Świetojańska, 2018
5. *Aethiopocassis garambana* Borowiec & Świetojańska, 2018
6. *Aethiopocassis guineensis* Borowiec & Świetojańska, 2018
7. *Aethiopocassis huilaensis* Borowiec & Świetojańska, 2018
8. *Aethiopocassis longidoana* Borowiec & Świetojańska, 2018
9. *Aethiopocassis transvaalensis* Borowiec & Świetojańska, 2018
10. *Agroiconota atromaculata* Borowiec, 2005
11. *Agroiconota atropunctata* Borowiec, 2005
12. *Agroiconota carlobrivioi* Borowiec, 2005
13. *Agroiconota gibbipennis* Borowiec, 2005
14. *Agroiconota paraguayana* Borowiec, 2005
15. *Agroiconota sanareensis* Borowiec, 2005
16. *Aslamidium bolivianum* Borowiec & Sassi, 2001
17. *Aslamidium coca* Borowiec, 1998
18. *Aslamidium ecuadoricum* Borowiec, 1998
19. *Aslamidium lescheni* Borowiec, 1998
20. *Aslamidium pichinchaensis* Borowiec, 1998
21. *Aspidomorpha andrei* Borowiec, 1997
22. *Aspidomorpha angoramensis* Borowiec, 1992
23. *Aspidomorpha atrodorsata* Borowiec, 1997
24. *Aspidomorpha bertiae* Borowiec, 1997
25. *Aspidomorpha collarti* Borowiec, 1997
26. *Aspidomorpha corrugata* Borowiec, 1997
27. *Aspidomorpha equatoriensis* Borowiec, 1986
28. *Aspidomorpha fampanamboensis* Borowiec, 1997
29. *Aspidomorpha filiola* Borowiec, 1997
30. *Aspidomorpha flaviceps* Borowiec, 1997
31. *Aspidomorpha gruevi* Borowiec, 1985
32. *Aspidomorpha heroni* Borowiec, 1997
33. *Aspidomorpha hiekei* Borowiec, 1997
34. *Aspidomorpha incerta* Borowiec, 1997
35. *Aspidomorpha karamojana* Borowiec & Sekerka, 2010
36. *Aspidomorpha levissima* Borowiec, 1997
37. *Aspidomorpha maffinbayensis* Borowiec, 1992
38. *Aspidomorpha mirabilis* Borowiec, 1997
39. *Aspidomorpha montanella* Borowiec, 1997
40. *Aspidomorpha muehlei* Borowiec, 1997
41. *Aspidomorpha obuduensis* Borowiec, 1997
42. *Aspidomorpha proshynskii* Borowiec, 1985
43. *Aspidomorpha pseudoareata* Borowiec, 1997
44. *Aspidomorpha pseudochlorina* Borowiec, 1986
45. *Aspidomorpha rainoni* Borowiec & Coache, 2018
46. *Aspidomorpha rubroornata* Borowiec, 1997
47. *Aspidomorpha salazarensis* Borowiec, 1997
48. *Aspidomorpha sankuruensis* Borowiec, 1997
49. *Aspidomorpha sassii* Borowiec, 1997
50. *Aspidomorpha sculpturata* Borowiec, 1997
51. *Aspidomorpha setosa* Borowiec, 1997
52. *Aspidomorpha silfverbergi* Borowiec, 1985
53. *Aspidomorpha sumbavaensis* Świetojańska & Borowiec, 2013
54. *Aspidomorpha tanolaensis* Borowiec, 1997

55. *Aspidimorpha tibetana* Świętojańska & Borowiec, 2006
56. *Aspidimorpha tuberosa* Borowiec, 1997
57. *Aspidimorpha uluguruensis* Borowiec, 1997
58. *Aspidimorpha zambiana* Borowiec, 2006
59. *Basiprionota flavicornis* Borowiec, 1993
60. *Basiprionota lomholdti* Borowiec, 1991
61. *Basiprionota sumba* Borowiec, 2006
62. *Basipta sebastiani* Borowiec, 2002
63. *Botanochara matogrossoensis* Borowiec, 2006
64. *Botanochara vianai* Borowiec, 1989
65. *Bradycassis agroiconotoides* Borowiec, 2005
66. *Bradycassis matogrossoensis* Świętojańska & Borowiec, 1996
67. *Brontispa caledonica* Borowiec, Świętojańska & Sekerka, 2019
68. *Caledonispā bivittata* Borowiec, Świętojańska & Sekerka, 2019
69. *Caledonispā panieensis* Borowiec, Świętojańska & Sekerka, 2019
70. *Caledonispā spinosa* Borowiec, Świętojańska & Sekerka, 2019
71. *Calliaspis cerdai* Borowiec, 2003
72. *Calliaspis sachaensis* Borowiec & Stojczew, 1998
73. *Calliaspis surinamensis* Borowiec, 2000
74. *Calliaspis wegrzynowiczi* Borowiec & Stojczew, 1998
75. *Carlobruchia boliviana* Borowiec, 1995
76. *Cassida acutangula* Borowiec, 1999
77. *Cassida ambrica* Borowiec, 1999
78. *Cassida andapaensis* Borowiec, 1988
79. *Cassida andohahelana* Borowiec & Świętojańska, 2013
80. *Cassida angulicollis* Borowiec & Świętojańska, 2013
81. *Cassida anosyennesensis* Borowiec & Świętojańska, 2013
82. *Cassida antoni* Borowiec & Świętojańska, 1997
83. *Cassida atroannulus* Borowiec & Świętojańska, 2013
84. *Cassida atrofemorata* Borowiec & Sassi, 2002
85. *Cassida atromarginata* Borowiec & Świętojańska, 2013
86. *Cassida atropunctata* Borowiec & Świętojańska, 2013
87. *Cassida atrorubra* Borowiec, 1999
88. *Cassida atrosignata* Sekerka & Borowiec, 2008
89. *Cassida bamendana* Borowiec & Świętojańska, 2022
90. *Cassida benaadirensis* Borowiec & Świętojańska, 2022
91. *Cassida beondrokana* Borowiec & Świętojańska, 2013
92. *Cassida beniowskii* Borowiec, 1988
93. *Cassida bezdeki* Borowiec, 2002
94. *Cassida brooksi* Borowiec, 1992
95. *Cassida bulirschi* Borowiec & Świętojańska, 2013
96. *Cassida butterwecki* Borowiec, 2007
97. *Cassida capensis* Borowiec, 2005
98. *Cassida Chiangmaiensis* Borowiec, 2001
99. *Cassida chrysanthemoides* Borowiec & Świętojańska, 2001
100. *Cassida coelebs* Borowiec, 1999
101. *Cassida crucipennis* Borowiec, 2003
102. *Cassida dehradunensis* Borowiec & Takizawa, 1991
103. *Cassida dembickyi* Borowiec, 2001
104. *Cassida densestriata* Borowiec & Świętojańska, 2013
105. *Cassida devalaensis* Borowiec & Takizawa, 1991
106. *Cassida diversepunctata* Borowiec & Świętojańska, 2001

107. *Cassida dolens* Borowiec, 1999
108. *Cassida drakensbergensis* Borowiec, 2005
109. *Cassida elgonensis* Borowiec & Świętojańska, 2022
110. *Cassida emontinensis* Borowiec & Świętojańska, 2022
111. *Cassida eximia* Borowiec & Ghate, 2004
112. *Cassida flavooculata* Borowiec & Świętojańska, 2013
113. *Cassida foveolatipennis* Borowiec & Świętojańska, 2001
114. *Cassida fuscomacula* Borowiec, 1988
115. *Cassida garambana* Borowiec & Świętojańska, 2022
116. *Cassida heroni* Borowiec & Świętojańska, 2022
117. *Cassida hovacassiformis* Borowiec, 1999
118. *Cassida ictericiformis* Borowiec, 2009
119. *Cassida impressipennis* Borowiec & Świętojańska, 2013
120. *Cassida inopinata* Sassi & Borowiec, 2006
121. *Cassida isarogensis* Borowiec, 2009
122. *Cassida johnsoni* Borowiec, 1988
123. *Cassida kantnerorum* Borowiec & Świętojańska, 2022
124. *Cassida kinabaluensis* Borowiec, 1999
125. *Cassida koreana* Borowiec & Cho, 2011
126. *Cassida laccopteroides* Borowiec & Świętojańska, 2013
127. *Cassida langeri* Borowiec, 2009
128. *Cassida laotica* Borowiec, 2002
129. *Cassida lawrencei* Borowiec 1990
130. *Cassida lateritioides* Borowiec & Świętojańska, 2013
131. *Cassida leleupi* Borowiec & Świętojańska, 2022
132. *Cassida liliputana* Borowiec & Świętojańska, 2013
133. *Cassida limpopoana* Borowiec & Świętojańska, 2001
134. *Cassida linnavuori* Borowiec, 1986
135. *Cassida lukasi* Borowiec & Świętojańska, 2013
136. *Cassida lycii* Borowiec & Świętojańska, 2001
137. *Cassida madagascarica* Borowiec, 1999
138. *Cassida malaysiana* Borowiec, 2010
139. *Cassida minutissima* Borowiec & Świętojańska, 2022
140. *Cassida mishmiensis* Sekerka & Borowiec, 2008
141. *Cassida montana* Borowiec, 1999
142. *Cassida monticola* Borowiec, 1988
143. *Cassida morondaviana* Borowiec, 2007
144. *Cassida mroczkowskii* Borowiec & Świętojańska, 1997
145. *Cassida multistrigata* Borowiec & Świętojańska, 2013
146. *Cassida multitiberculata* Borowiec & Świętojańska, 2013
147. *Cassida mysorensis* Borowiec & Świętojańska, 1996
148. *Cassida namibiensis* Borowiec, 2005
149. *Cassida nigeriana* Borowiec & Świętojańska, 2022
150. *Cassida nigroflavens* Borowiec, 1988
151. *Cassida nigrohumeralis* Borowiec & Ghate, 2004
152. *Cassida nigropunctata* Borowiec & Świętojańska, 2013
153. *Cassida nilgiriensis* Borowiec & Takizawa, 1991
154. *Cassida nosybeensis* Borowiec & Świętojańska, 2013
155. *Cassida oromiaensis* Borowiec & Świętojańska, 2022
156. *Cassida overlaeti* Borowiec & Świętojańska, 2022
157. *Cassida oxylepiformis* Borowiec & Świętojańska, 2022
158. *Cassida pacholatkoii* Sekerka & Borowiec, 2008

159. *Cassida paiensis* Borowiec, 2001
160. *Cassida pauliani* Borowiec, 1999
161. *Cassida paveli* Borowiec & Świętojańska, 2013
162. *Cassida polymeriae* Borowiec & Burwell, 2011
163. *Cassida praensis* Borowiec, 2001
164. *Cassida pretiosa* Borowiec, 1988
165. *Cassida pseudolateritia* Borowiec & Świętojańska, 2013
166. *Cassida pseudostrumosa* Borowiec & Świętojańska, 2013
167. *Cassida pseudovicinalis* Borowiec & Świętojańska, 2013
168. *Cassida pubipennis* Borowiec, 1999
169. *Cassida quadricolorata* Borowiec, 1999
170. *Cassida queenslandica* Borowiec, 2006
171. *Cassida rabaiensis* Borowiec & Świętojańska, 2022
172. *Cassida reticulipennis* Borowiec & Świętojańska, 2001
173. *Cassida rogezensis* Borowiec & Świętojańska, 2013
174. *Cassida rogozinskii* Borowiec & Świętojańska, 2022
175. *Cassida rubripennis* Borowiec, 2002
176. *Cassida sabahensis* Świętojańska & Borowiec, 2002 e
177. *Cassida satanas* Borowiec & Świętojańska, 2022
178. *Cassida sculpturipennis* Borowiec & Świętojańska, 2013
179. *Cassida scymnoides* Borowiec, 1999
180. *Cassida sekerkai* Borowiec & Świętojańska, 2017
181. *Cassida senegalensis* Borowiec & Świętojańska, 2022
182. *Cassida seniculoides* Borowiec, 1999
183. *Cassida shimba* Borowiec & Świętojańska, 2022
184. *Cassida silvicola* Borowiec, 1988
185. *Cassida sparteiformis* Borowiec & Świętojańska, 2022
186. *Cassida spatiosiformis* Borowiec & Świętojańska, 2001
187. *Cassida stipidosa* Borowiec & Świętojańska, 2022
188. *Cassida strigaticollis* Borowiec, 1988
189. *Cassida subacuticollis* Borowiec, 1999
190. *Cassida thailandica* Borowiec, 2001
191. *Cassida tianshanica* Borowiec & Świętojańska, 2001
192. *Cassida timorensis* Borowiec, 1995
193. *Cassida transcaucasica* Borowiec & Świętojańska, 2001
194. *Cassida trianguliformis* Borowiec & Świętojańska, 2013
195. *Cassida tryznai* Borowiec & Świętojańska, 2013
196. *Cassida tsaratanana* Borowiec, 1994
197. *Cassida ultima* Borowiec & Świętojańska, 2013
198. *Cassida umbonata* Borowiec, 1999
199. *Cassida unica* Świętojańska & Borowiec, 2002
200. *Cassida villiersi* Borowiec & Świętojańska, 2022
201. *Cassida voiensis* Borowiec & Świętojańska, 2022
202. *Cassida wanati* Borowiec & Świętojańska, 2022
203. *Cassida wittmeri* Borowiec & Świętojańska, 2022
204. *Chapadacassis paradoxa* Sekerka & Borowiec, 2015
205. *Charidotella atromarginata* Borowiec, 2009
206. *Charidotella bordoni* Borowiec, 2002
207. *Charidotella dominicanensis* Borowiec, 2011
208. *Charidotella ecuadorica* Borowiec, 1989
209. *Charidotella kesseli* Borowiec, 1989
210. *Charidotella marginepunctata* Borowiec, 2004

211. *Charidotella moraguesi* Borowiec, 2007
212. *Charidotella nigripennis* Borowiec, 2009
213. *Charidotella pacata* Borowiec, 2007
214. *Charidotella steinhauseni* Borowiec, 1989
215. *Charidotella santaremi* Borowiec, 1995
216. *Charidotis diabolica* Świętojańska Borowiec, 2000
217. *Charidotis tuberculata* Świętojańska Borowiec, 2000
218. *Chelymorpha aculeata* Borowiec, 2000
219. *Chiridopsis atricollis* Borowiec, 2005
220. *Chiridopsis ghati* Borowiec & Świętojańska, 2000
221. *Chiridopsis klapperichi* Świętojańska & Borowiec, 2017
222. *Chiridopsis levis* Borowiec, 2005
223. *Chiridopsis maculata* Borowiec, 2005
224. *Chiridopsis marginepunctata* Borowiec, 2005
225. *Chiridopsis nigropunctata* Borowiec & Ghate, 1999
226. *Chiridopsis nigroreticulata* Borowiec, 2005
227. *Chiridopsis rubromaculata* Borowiec, Ranade, Rane & Ghate, 2001
228. *Chiridopsis tanzaniensis* Świętojańska & Borowiec, 2017
229. *Chiridopsis zambiana* Świętojańska & Borowiec, 2017
230. *Conchyloctenia aspidiformis* Borowiec, 1994
231. *Conchyloctenia capensis* Świętojańska & Borowiec, 2002
232. *Crambelea minuta* Borowiec, 2009
233. *Craspedonta laotica* Świętojańska & Borowiec, 2000
234. *Craspedonta levis* Borowiec, 1993
235. *Cteisella flava* Borowiec, 2004
236. *Cyclocassis secunda* Borowiec, 1998
237. *Cyclosoma bicostata* Borowiec, 1999
238. *Cyrtonota abrili* Borowiec & Świętojańska, 2015
239. *Cyrtonota bergeali* Borowiec & Sassi, 1999
240. *Cyrtonota botanocharoides* Borowiec, 1989
241. *Cyrtonota bugaensis* Borowiec & Sassi, 1999
242. *Cyrtonota christophori* Borowiec, 1998
243. *Cyrtonota gibbera* Borowiec, 1989
244. *Cyrtonota machupicchu* Borowiec & Sassi, 1999
245. *Cyrtonota montana* Borowiec, 2000
246. *Cyrtonota nitida* Borowiec & Sassi, 1999
247. *Cyrtonota ruforeticulata* Borowiec, 2007
248. *Cyrtonota santanderensis* Borowiec, 2009
249. *Demotispa brunneofasciata* Borowiec, 2000
250. *Discomorpha araii* Borowiec & Takizawa, 2011
251. *Discomorpha atahualpai* Borowiec, 2006
252. *Discomorpha carlobrivioi* Borowiec & Sassi, 2003
253. *Discomorpha davidsoni* Borowiec & Dąbrowska, 1996
254. *Discomorpha giganteensis* Borowiec, 2006
255. *Discomorpha howdenorum* Borowiec, 2006
256. *Discomorpha onorei* Borowiec, 1998
257. *Discomorpha putamayoensis* Borowiec, 2006
258. *Discomorpha sanataremi* Borowiec & Dąbrowska, 1996
259. *Discomorpha seckyi* Borowiec, 2001
260. *Discomorpha tigrensensis* Borowiec & Takizawa, 2011
261. *Dorynota boliviana* Borowiec, 2005
262. *Dorynota carlosi* Borowiec & Takizawa, 2011

263. *Dorynota matogrossoensis* Borowiec, 2005
264. *Dorynota rileyi* Borowiec, 1994
265. *Dorynota storki* Borowiec, 2006
266. *Dorynota viridimetallica* Borowiec, 2005
267. *Echoma amapaensis* Borowiec & Takizawa, 2011
268. *Echoma anaglyptoides* Borowiec, 1998
269. *Eugenusa jasiniskii* Borowiec & Dąbrowska, 1997
270. *Eugenysa martae* Borowiec, 1987
271. *Eugenysa unicolor* Borowiec & Dąbrowska, 1997
272. *Herminella flavocostata* Borowiec & Świętojańska, 2014
273. *Herminella quadrimaculata* Borowiec & Świętojańska, 2014
274. *Hermosacantha madagascariensis* Borowiec & Świętojańska, 2014
275. *Hilarocassis bordoni* Borowiec, 2002
276. *Hispa waiensis* Borowiec & Świętojańska, 2007
277. *Hovacassis brunneofasciata* Borowiec, 2002
278. *Hovacassis flavonigra* Borowiec, 2002
279. *Hovacassis formosa* Borowiec, 2002
280. *Hovacassis murzini* Borowiec, 2002
281. *Hovacassis rubromaculata* Borowiec, 2002
282. *Hovacassis rubrovittata* Borowiec, 2002
283. *Hybosa marginepunctata* Borowiec, 2011
284. *Hybosa santaritae* Borowiec & Takizawa, 2011
285. *Hypocassida convexipennis* Borowiec, 2000
286. *Ischnocodia flavofasciata* Borowiec, 1995
287. *Ischyronota brisouti* ssp. *jordanensis* Borowiec, 1986
288. *Isopedhispa costata* Borowiec, Świętojańska & Sekerka, 2019
289. *Isopedhispa latemarginata* Borowiec, Świętojańska & Sekerka, 2019
290. *Laccoptera caduca* Borowiec, 1994
291. *Laccoptera nunbergi* Borowiec, 1994
292. *Laccoptera obscuromaculata* Borowiec, 2010
293. *Laccoptera rotundicollis* Borowiec, 1997
294. *Laccoptera warchalowskii* Borowiec, 1985
295. *Limnocassis compacta* Borowiec & Sekerka, 2015
296. *Limnocassis regularis* Borowiec & Sekerka, 2015
297. *Lorentzocassis riedeli* Borowiec, 2003
298. *Megapyga minima* Borowiec, 1998
299. *Megapyga obscuricollis* Borowiec, 1993
300. *Metrioronella tucumana* Borowiec, 2006
301. *Metrioronella tumacoensis* Borowiec, 2002
302. *Microctenochira belizensis* Borowiec, 2007
303. *Microctenochira brasiliensis* Borowiec & Świętojańska, 1999
304. *Microctenochira chapada* Świętojańska & Borowiec, 1995
305. *Microctenochira danielssoni* Świętojańska & Borowiec, 1995
306. *Microctenochira mapiriensis* Borowiec, 2002
307. *Microctenochira obscurata* Borowiec & Świętojańska, 1999
308. *Microctenochira panamensis* Świętojańska & Borowiec, 1999
309. *Miocalaspis ecuadorica* Borowiec, 2000
310. *Miocalaspis flavofasciata* Borowiec, 2000
311. *Nebraspis viridimetallica* Borowiec, 1999
312. *Notosacantha bertounesquei* Borowiec & Le Tirant, 2013
313. *Notosacantha darjeelingensis* Borowiec & Takizawa, 1991
314. *Notosacantha flaviventris* Borowiec & Takizawa, 2013

315. *Notosacantha jammuensis* Borowiec & Takizawa, 1991
316. *Notosacantha kantneri* Świętojańska Borowiec, 1999
317. *Notosacantha komiyai* Dąbrowska & Borowiec, 1996
318. *Notosacantha nathani* Borowiec & Takizawa, 1991
319. *Notosacantha nepalensis* Borowiec & Takizawa, 1991
320. *Notosacantha nilgiriensis* Borowiec & Takizawa, 1991
321. *Notosacantha obscuricornis* Borowiec & Takizawa, 2013
322. *Notosacantha ovoidea* Borowiec & Świętojańska, 2013
323. *Notosacantha pakistanica* Borowiec, Sultan & Rafi, 2008
324. *Notosacantha sabahensis* Borowiec & Świętojańska, 1999
325. *Notosacantha sulawesica* Borowiec, 1999
326. *Notosacantha sumbawaensis* Świętojańska Borowiec, 1999
327. *Notosacantha timorensis* Świętojańska & Borowiec 2016
328. *Nuzonia amazonica* Borowiec 2000
329. *Nuzonia atromaculata* Borowiec, 2000
330. *Nuzonia brevicornis* Borowiec, 1998
331. *Nuzonia marginepunctata* Borowiec, 2000
332. *Omaspides boliviana* Borowiec, 2003
333. *Omaspides confusa* Borowiec, 2010
334. *Omaspides curiosa* Borowiec & Takizawa, 2011
335. *Omaspides picaflorensis* Borowiec, 2010
336. *Omaspides unicolor* Borowiec, 1998
337. *Omocerus angulicollis* Borowiec, 2000
338. *Omocerus burakowskii* Dąbrowska & Borowiec, 1995
339. *Omocerus doeberli* Dąbrowska & Borowiec, 1995
340. *Omocerus rugosicollis* Borowiec, 2000
341. *Omocerus similis* Borowiec, 2000
342. *Omocerus tenebrosus* Borowiec & Takizawa, 2011
343. *Orexita minima* Borowiec & Świętojańska, 1997
344. *Orexita woytkowskii* Borowiec, 1992
345. *Oxylepus bituberculatus* Borowiec, 2002
346. *Oxylepus boroveci* Borowiec, 2001
347. *Oxylepus convexicollis* Borowiec, 2002
348. *Oxylepus grobbelaarae* Borowiec, 2002
349. *Oxylepus impressipennis* Borowiec, 2002
350. *Oxylepus intermedius* Borowiec, 2002
351. *Oxylepus planicollis* Borowiec, 2002
352. *Oxylepus sextuberculatus* Borowiec, 2002
353. *Paraselenis convolvuli* Villamil, Borowiec & Świętojańska, 2023
354. *Paraselenis flavopunctata* Borowiec, 2003
355. *Paraselenis nigropunctata* Borowiec, 2003
356. *Paraselenis saltaensis* Borowiec, 2002
357. *Paratrikona albomaculata* Borowiec, 2009
358. *Physonota convexa* Borowiec, 1995
359. *Physonota puncticollis* Borowiec, 1995
360. *Plagiometriona costaricensis* Borowiec, 2001
361. *Plagiometriona latemarginata* Borowiec, 2001
362. *Plagiometriona losi* Borowiec, 1998
363. *Plagiometriona obrienorum* Borowiec & Świętojańska, 2013
364. *Pseudostilpnaspis belizensis* Borowiec, 2008
365. *Pseudostilpnaspis costaricana* Borowiec, 2000
366. *Pseudostilpnaspis muzoensis* Borowiec, 2000

367. *Rhytidocassis angulipennis* Borowiec, 2002
368. *Rhytidocassis lopatini* Borowiec & Świętojańska, 2001
369. *Spaethaspis peruviana* Borowiec, 2000
370. *Sphenocassis anosibensis* Borowiec, 2002
371. *Sphenocassis impressipennis* Borowiec, 2002
372. *Sphenocassis rotundella* Borowiec, 2002
373. *Spilophora cuneata* Borowiec, 2003
374. *Spilophora lacrimata* Borowiec, 2003
375. *Stilpnaspis bicolorata* Borowiec, 2000
376. *Stilpnaspis filicornis* Borowiec, 2000
377. *Stilpnaspis impunctata* Borowiec, 2000
378. *Stilpnaspis monteverdensis* Borowiec, 2000
379. *Stilpnaspis panamensis* Borowiec, 2000
380. *Stilpnaspis tambitoensis* Borowiec, 2000
381. *Stolas agenysiformis* Borowiec, 2011
382. *Stolas armirantensis* Borowiec, 1999
383. *Stolas atalayensis* Borowiec & Takizawa, 2011
384. *Stolas echoma* Borowiec, 1998
385. *Stolas erectepilosa* Borowiec, 1998
386. *Stolas flavomarginata* Borowiec, 1999
387. *Stolas flavoradiata* Borowiec & Stach, 2012
388. *Stolas hameli* Borowiec, 2007
389. *Stolas imitatrix* Borowiec, 1999
390. *Stolas intermedia* Borowiec, 1999
391. *Stolas napoensis* Borowiec, 1998
392. *Stolas omaspidiformis* Borowiec, 2007
393. *Stolas perezii* Borowiec, 1998
394. *Stolas pucallpaensis* Borowiec & Takizawa, 2011
395. *Stolas radiata* Borowiec & Stach, 2013
396. *Stolas sanguineovittata* Borowiec, 1998
397. *Stolas sanramonensis* Borowiec, 2005
398. *Stolas stolidi* ssp. *jadwischczaki* Borowiec, 1998
399. *Stolas tachiraensis* Borowiec, 2009
400. *Stolas uniformis* Borowiec & Świętojańska, 2010
401. *Stolas warchalowskii* Borowiec, 2007
402. *Stolas zumbaensis* Borowiec, 1998
403. *Syngambria panamensis* Borowiec, 2006
404. *Teretrispa daccordii* Borowiec, Świętojańska & Sekerka, 2019
405. *Teretrispa longicollis* Borowiec, Świętojańska & Sekerka, 2019
406. *Thlaspida tsoui* Borowiec & Lee, 2009
407. *Thlaspidula riedeli* Borowiec & Świętojańska, 2001
408. *Torquispa convexifrons* Borowiec, Świętojańska & Sekerka, 2019
409. *Trichaspis brevicornis* Borowiec, 2002
410. *Trichaspis erinacea* Borowiec, 2002
411. *Trichaspis louwi* Borowiec, 2002
412. *Trichaspis minutissima* Borowiec, 2002
413. *Trichaspis tomentosa* Borowiec, 2002
414. *Trilaccodea ecuadorica* Borowiec, 2007
415. *Trilaccodea meridionalis* Borowiec, 2004
416. *Wanatispa cylindricollis* Borowiec, Świętojańska & Sekerka, 2019
417. *Wanatispa rutai* Borowiec, Świętojańska & Sekerka, 2019

References

Complete list of papers on tortoise beetles published by Lech Borowiec chronologically—papers with descriptions of new taxa whose types are in Lech Borowiec and MNHW collections marked in bold

- Borowiec, L. (1984) On the synonymy in *Imatidium* sensu lato (Coleoptera, Chrysomelidae, Cassidinae). *Polskie Pismo Entomologiczne*, 54, 411–412.
- Borowiec, L. (1985 a) Asiatic Cassidinae (Coleoptera, Chrysomelidae) in Polish collections. *Polskie Pismo Entomologiczne*, 55, 25–38.
- Borowiec, L. (1985 b) Contribution to the knowledge of African Cassidinae, 1 (Coleoptera, Chrysomelidae). *Polskie Pismo Entomologiczne*, 55, 223–244.**
- Borowiec, L. (1985 c) Contribution to the knowledge of African Cassidinae, 2 (Coleoptera, Chrysomelidae). *Polskie Pismo Entomologiczne*, 55, 439–450.
- Borowiec, L. (1985 d) *Aspidomorpha gruevi* n. sp. (Coleoptera, Chrysomelidae, Cassidinae) from Yemen. *Polskie Pismo Entomologiczne*, 55, 451–456.**
- Borowiec, L. (1986 a) Contribution to the knowledge of African Cassidinae, 3 (Coleoptera, Chrysomelidae). *Polskie Pismo Entomologiczne* [1985], 55, 791–809.**
- Borowiec, L. (1986 b) A new species and a new subspecies of tortoise-beetles from the Middle East (Coleoptera, Chrysomelidae, Cassidinae). *Polskie Pismo Entomologiczne*, 56, 573–576.**
- Borowiec, L. (1987 a) A new species of *Eugenysa* Chevrolat from Colombia (Coleoptera, Chrysomelidae, Cassidinae). *Polskie Pismo Entomologiczne* 1986, 56, 883–885.**
- Borowiec, L. (1987 b) Lectotype designations for some Aspidomorphini described by Weise (Coleoptera, Chrysomelidae, Cassidinae). *Polskie Pismo Entomologiczne*, 57, 413–415.
- Borowiec, L. (1987 c) On the genus *Rhacocassis* Spaeth, 1904 (Coleoptera, Chrysomelidae, Cassidinae). *Polskie Pismo Entomologiczne*, 57, 453–460.
- Borowiec, L. (1988). Review of the genus *Parachirida* Hincks, 1952 (Coleoptera, Chrysomelidae, Cassidinae). *Polskie Pismo Entomologiczne*, 58, 301–308.
- Borowiec, L. (1988 b) Ten new species of *Cassida* L. from Madagascar (Coleoptera, Chrysomelidae, Cassidinae). *Polskie Pismo Entomologiczne*, 58, 547–569.**
- Borowiec, L. (1989 a) A new species of *Botanochara* Dejean, 1837, from Argentina (Coleoptera, Chrysomelidae, Cassidinae). *Polskie Pismo Entomologiczne* [1988], 58, 725–727.**
- Borowiec, L. (1989 b) Three new species of *Charidotella* Weise (Coleoptera, Chrysomelidae, Cassidinae), with checklist of the genus. *Polskie Pismo Entomologiczne*, 59, 203–222.**
- Borowiec, L. (1989 c) Two new species of *Cyrtonota* Chevrolat (Coleoptera, Chrysomelidae, Cassidinae). *Elytron*, 3, 25–30.**
- Borowiec, L. (1990 a) New records and new synonyms of Asiatic Cassidinae (Coleoptera, Chrysomelidae). *Polskie Pismo Entomologiczne*, 59, 677–711.
- Borowiec, L. (1990 b) A review of the genus *Cassida* L. of the Australian Region and Papuan Subregion (Coleoptera, Chrysomelidae, Cassidinae). *Genus*, 1, 1–51.
- Borowiec, L. (1990 c) A new species of *Basiprionota* Chevrolat from Philippines (Coleoptera, Chrysomelidae, Cassidinae). *Polskie Pismo Entomologiczne*, 60, 167–172.
- Borowiec, L. (1991 a) On the genus *Malayocassis* Spaeth, 1952 (Coleoptera, Chrysomelidae, Cassidinae). *Polskie Pismo Entomologiczne* [1990], 60, 3–7.
- Borowiec, L. (1991 b) On the genera *Emdenia* Spaeth, 1915 and *Austrosecadia* Hincks, 1950 (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 2, 1–10.
- Borowiec, L. (1991 c) On the genus *Lorentzocassis* Spaeth, 1913 (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 2, 11–17.
- Borowiec, L. & Takizawa H. (1991 d) Notes on chrysomelid beetles (Coleoptera) of India and its neighboring areas. Part 10. *Japanese Journal of Entomology*, 59, 637–654.**
- Borowiec, L. (1992 a) A new species of *Cassida* L. from Madagascar (Coleoptera: Chrysomelidae: Cassidinae). *Genus* [1991], 2, 283–285.**
- Borowiec, L. (1992 b) A new species of *Orexita* Spaeth from Peru (Coleoptera: Chrysomelidae: Cassidinae). *Genus* [1991], 2, 287–289.
- Borowiec, L. (1992 c) A review of the tribe Aspidomorphini of the Australian Region and Papuan Subregion (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 3, 121–184.**
- Borowiec, L. (1993) Three new species of Basiprionotini from Oriental Region (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 4, 79–85.**
- Borowiec, L. (1994 a) New synonyms in the Cassidinae (Coleoptera: Chrysomelidae). *Genus*, 5, 153–159.
- Borowiec, L. (1994 b) *Dorynota rileyi* n. sp. from Paraguay (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 5, 161–164.**
- Borowiec, L. (1994 c) *A monograph of the Afrotropical Cassidinae (Coleoptera: Chrysomelidae). Part I. Introduction, morphology, key to the genera, and reviews of the tribes Epistictinini, Basiprionotini and Aspidimorphini (except the***

genus *Aspidimorpha*. Genus (suppl.), Biologica Silesiae, Wrocław, 276 pp.

- Borowiec, L. (1995 a) Cassidinae (Coleoptera: Chrysomelidae) of Namibia. *Mitteilungen aus dem Zoologischen Museum in Berlin*, 71, 435–438.
<https://doi.org/10.1002/mmnz.19950710215>
- Borowiec, L. (1995 b) A new species of *Ischnocodia* Spaeth, 1942 from Brazil (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 6, 91–94.
- Borowiec, L. (1995 c) *Cassida timorensis* n. sp. from Timor Is. (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 6, 95–98.
- Dąbrowska, A. & Borowiec, L. (1995 d) Two new species of *Omocerus (Paratauroma)* from Brazil (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 6, 415–422.
- Borowiec, L. (1995 e) *Carlobruchia* (s. str.) *boliviana* n. sp. from Bolivia (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 6, 423–427.
- Borowiec, L. (1995 f) Two new species of *Physonota* Boh. from Mexico, and notes on *Ph. cerea* Boh. (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 6, 429–439.
- Borowiec, L. (1995 g) *Charidotella (Metrionaspis) santaremi* n. sp., from Brazil (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 6, 441–446.
- Świętojańska, J. & Borowiec, L. (1995 h) Two new species of *Microctenochira* Spaeth from Brazil and Peru (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 6, 447–454.
- Borowiec, L. (1995 i) Tribal classification of the cassidoid Hispinae (Coleoptera: Chrysomelidae). In: Pakaluk, J. & Ślipinski S.A. (Eds.), *Biology, Phylogeny, and Classification of Coleoptera: Papers Celebrating the 80th Birthday of Roy A. Crowson*. Warszawa, pp. 541–558.
- Borowiec, L. (1996 a) New records of Asiatic Cassidinae (Coleoptera: Chrysomelidae). *Annals of the Upper Silesian Museum in Bytom (Entomology)*, 6-7, 5–47.
- Borowiec, L. (1996 b) Faunistic records of Neotropical Cassidinae (Coleoptera: Chrysomelidae). *Polskie Pismo Entomologiczne*, 65, 119–251.
- Dąbrowska, A. & Borowiec, L. (1996 c) *Notosacantha komiyai* n. sp. from Thailand, with notes on another two species (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 7, 451–458.
- Borowiec, L. & Dąbrowska, A. (1996 d) Two new species of *Discomorpha* Chevrolat, 1837 from Brazil (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 7, 459–466.
- Borowiec, L. & Świętojańska, J. (1996 e) *Cassida mysorensis* n. sp. from India (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 7, 467–470.
- Świętojańska, J. & Borowiec, L. (1996 f) A new species of *Bradycassis* Spaeth from Brazil (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 7, 471–474.
- Borowiec, L. (1997 a) New synonyms and notes on some Cassidinae species, with description of *Laccoptera rotundicollis* n. sp. from West Africa (Coleoptera: Chrysomelidae). *Genus*, 8, 659–672.
- Borowiec, L. & Dąbrowska, A. (1997 b) Two new species of *Eugenysa* Chevrolat, 1837 from Ecuador and Peru (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 8, 673–678.
- Borowiec, L. & Świętojańska, J. (1997 c) A new species of *Orexita* Spaeth from Brazil and Bolivia (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 8, 679–683.
- Borowiec, L. & Świętojańska, J. (1997 d) Two new species of *Cassida* L. (Coleoptera: Chrysomelidae: Cassidinae) from the Oriental Region. *Annales Zoologici*, Warszawa, 47, 275–278.
- Borowiec, L. & Świętojańska, J. (1997 e) *Cassida leucanthemi* Bordy i *C. bergali* Bordy (Coleoptera, Chrysomelidae), gatunki nowe dla fauny Polski. *Wiadomości entomologiczne*, 15, 237–240 [*Cassida leucanthemi* Bordy and *C. bergali* Bordy (Coleoptera, Chrysomelidae), new to the Polish fauna]
- Borowiec, L., Chikatunov, V. & Halperin, J. (1997 f) Cassidinae (Coleoptera: Chrysomelidae) of Israel. *Israel Journal of Entomology*, 31, 147–152.
- Borowiec, L. (1997 g) *A monograph of the Afrotropical Cassidinae (Coleoptera: Chrysomelidae). Part II. Revision of the tribe Aspidimorphini 2, the genus Aspidimorpha Hope*. Genus (suppl.), Biologica Silesiae, Wrocław, 595 pp.
- Borowiec, L. (1998 a) *Megapyga minima*, n. sp. from Borneo (Coleoptera: Chrysomelidae: Cassidinae). *Serangga*, 3, 103–106.
- Borowiec, L. & Stojczew, A. (1998 b) Two new species of *Calliaspis* Boheman, 1850 from Ecuador (Coleoptera: Chrysomelidae: Hispinae). *Annales Zoologici*, Warszawa, 48, 325–328.
- Borowiec, L. (1998 c) Four new species of *Aslamidium* Borowiec, with description of *Neoaslamidium* new subgenus (Coleoptera: Chrysomelidae: Hispinae). *Genus*, 9, 367–374.
- Borowiec, L. (1998 d) Review of the Cassidinae of Ecuador, with a description of thirteen new species (Coleoptera: Chrysomelidae). *Genus*, 9, 155–246.
- Borowiec, L. (1998 e) *Echoma anaglypta* (Boheman, 1862) and *E. anaglyptoides* n. sp. (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 9, 375–385.
- Borowiec, L. (1998 f) *Agroiconota judaica* (Fabricius) and *A. inedita* (Boheman) - distinct species (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 9, 387–393.
- Borowiec, L. (1999 a) *Cassida kinabaluensis* n. sp. from Sabah, Malaysia (Coleoptera: Chrysomelidae: Cassidinae).

Serangga, 4, 1–5.

- Borowiec, L. (1999 b) A new species of *Notosacantha* Chevrolat from Sulawesi, Indonesia (Coleoptera: Chrysomelidae: Cassidinae). *Serangga*, 4, 7–10.
- Borowiec, L. & Świętojańska, J. (1999 c) A new species of *Notosacantha* Chevrolat from Sabah, Malaysia (Coleoptera: Chrysomelidae: Cassidinae). *Serangga*, 4, 185–188.
- Borowiec, L. (1999 d) *Cyclosoma* (*Monrosiacassis*) *bicostata* n. sp. and notes on *C. (M.) puberula* (Boh.) (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 10, 81–87.
- Borowiec, L. & Sassi, D. (1999 e) Four new species of *Cyrtanota* Chevrolat (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 10, 89–101.
- Borowiec, L. & Ghate, H.V. (1999 f) *Chiridopsis nigropunctata* n. sp. and notes on *Ch. selecta* (Weise) (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 10, 103–108.
- Świętojańska, J. & Borowiec, L. (1999 g) Three new species of *Microctenochira* Spaeth from Brazil and Panama (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 10, 109–116.
- Świętojańska, J. & Borowiec, L. (1999 h) Two new species of *Notosacantha* Chevrolat from the Oriental Region (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 10, 421–426.
<https://doi.org/10.3161/000345411X584870>
- Borowiec, L. (1999 i) New species of the genera *Stolas* Billb. and *Nebraspis* Spaeth (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 10, 427–438.
- Borowiec, L. (1999 j) New species of *Cassida* L. from Madagascar (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 10, 439–477.
- Borowiec, L. (1999 k) *A world catalogue of the Cassidinae* (Coleoptera: Chrysomelidae). Biologica Silesiae, Wrocław, 476 pp.
- Świętojańska, J. & Borowiec, L. (2000 a) *Craspedonta laotica* n. sp. from Laos (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 11, 45–48.
- Borowiec, L. (2000 b) *Cyrtanota montana*, a new species from Ecuador (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 11, 49–52.
- Borowiec, L. & Świętojańska, J. (2000 c) *Chiridopsis ghatei*, a new species from India (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 11, 53–57.
- Borowiec, L. (2000 d) Three new species of *Nuzonia* Spaeth, 1912 (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 11, 59–67.
- Borowiec, L. (2000 e) Notes on the genus *Stilpnaspis* Weise, with a description of *Pseudostilpnaspis*, new genus and eleven new species of the tribe Imatidiini (Coleoptera: Chrysomelidae: cassidoid Hispinae). *Genus*, 11, 147–195.
- Borowiec, L. (2000 f) Three new species of *Omocerus* Chevrolat, 1835 (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 11, 219–228.
- Borowiec, L. (2000 g) Two new species of *Miocalaspis* Weise, 1899 from Ecuador and Peru (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 11, 229–233.
- Borowiec, L. (2000 h) *Chelymorpha aculeata*, a new species from Bolivia (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 11, 597–599.
- Borowiec, L. (2000 i) *Hypocassida convexipennis*, a new species from Iran (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 11, 601–605.
- Świętojańska, J. & Borowiec, L. (2000 j) Two new species of *Charidotis* Boheman from Ecuador and Brazil (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 11, 607–612.
- Borowiec, L. & Świętojańska, J. (2001 a) Revision of the *Cassida litigiosa* group from southern Africa (Coleoptera: Chrysomelidae: Cassidinae). *Annales Zoologici*, Warszawa, 51, 153–184.
- Borowiec, L. & Świętojańska, J. (2001 b) The Palearctic species of the genus *Rhytidocassis* Spaeth, 1941 (Coleoptera: Chrysomelidae: Cassidinae). *Annales Zoologici*, Warszawa, 51, 325–329.
- Borowiec, L. & Świętojańska, J. (2001 c). Revision of the genus *Thlaspidula* Spaeth, 1901 (Coleoptera: Chrysomelidae: Cassidinae). *Annales Zoologici*, Warszawa, 51, 331–343.
- Borowiec, L. (2001 d) *Discomorpha* (*Vulpia*) *secky*, a new species from Peru (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 12, 59–61.
- Borowiec, L. & Świętojańska, J. (2001 e) *Cassida undecimnotata* - a species complex (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 12, 63–79.
- Borowiec, L. (2001 f) Redescription of *Macromonycha anatolica* (Weise, 1900) (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 12, 81–86.
- Borowiec, L. (2001 g) *Oxylepus boroveci*, a new species from Tunisia (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 12, 349–352.
- Borowiec, L. (2001 h) Two new species of *Plagiometriona* Spaeth from Costa Rica and Panama. (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 12, 353–359.
- Borowiec, L., Ranade, S., Rane, N. & Ghate, H.V. (2001 i) *Chiridopsis rubromaculata* n. sp. from India, and notes on its bionomy and immature stages (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 12, 361–371.
- Borowiec, L. & Sassi, D. (2001 j) *Aslamidium* (s. str.) *bolivianum*, a new species from Bolivia (Coleoptera: Chrysomelidae: Hispinae). *Genus*, 12, 483–487.

- Borowiec, L. (2001 k) New records of Asian and Australopapuan Cassidinae, with a description of five new species of *Cassida* L. from Thailand (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 12, 493–562.
- Borowiec, L. (2002 a) New records of Neotropical Cassidinae, with description of three new species (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 13, 43–138.
- Borowiec, L. (2002 b) *Metrionella tumacoensis*, a new species from Colombia (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 13, 139–142.
- Borowiec, L. & Sassi D. (2002 c) A new species of *Cassida* L. from Palaearctic China (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 13, 143–147.
- Świętojańska, J. & Borowiec, L. (2002 d) A new species of *Conchyloctenia* Spaeth from South Africa and redescription of *Aspidimorpha kilimana* Weise, bona species (Coleoptera: Chrysomelidae: Cassidinae: Aspidimorphini). *Genus*, 13, 211–217.
- Świętojańska, J. & Borowiec, L. (2002 e) Two new species of *Cassida* L. from Borneo (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 13, 357–363.
- Borowiec, L. (2002 f) A new species of *Microctenochira* Spaeth from Bolivia (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 13, 365–369.
- Borowiec, L. & Świętojańska, J. (2002 g) Redescription of two neglected species: *Charidotis obtusa* (Boh.) and *Ch. plicatula* Boh. (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 13, 371–377.
- Borowiec, L. (2002 h) Three new species of *Cassida rati* group from Laos (Coleoptera: Chrysomelidae: Cassidinae). *Annales Zoologici*, Warszawa, 52, 567–572.
- Borowiec, L. (2002 i) *A monograph of the Afrotropical Cassidinae (Coleoptera: Chrysomelidae). Part III. Revision of the tribe Cassidini 1, except the genera Aethiopocassis Sp., Cassida L., and Chiridopsis Sp.* Genus, supplement, Biologica Silesiae, Wrocław, 292 pp. + 17 pl.
- Borowiec, L. (2003 a) A new species of *Calliaspis* Dejean, 1837 (Coleoptera: Chrysomelidae: Cassidinae) from French Guyana. *Zootaxa*, 148 (1), 1–6.
<https://doi.org/10.11646/zootaxa.148.1.1>
- Borowiec, L. & Świętojańska, J. (2003 b) The first instar larva of *Cassida nebulosa* L. (Coleoptera: Chrysomelidae: Cassidinae) - a model description. *Annales Zoologici*, Warszawa, 53, 325–329.
- Borowiec, L. (2003 c) A new species of *Cassida variabilis* group from Laos (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 14, 121–124.
- Chaboo, C.S. & Borowiec, L. (2003 d) Annotated checklist of tortoise beetles of Trinidad and Tobago. *Coleopterists Bulletin*, 57, 71–78.
[https://doi.org/10.1649/0010-065X\(2003\)057\[0071:ACOTBO\]2.0.CO;2](https://doi.org/10.1649/0010-065X(2003)057[0071:ACOTBO]2.0.CO;2)
- Borowiec, L. (2003 e) Nowe stanowiska gatunków z rodzaju *Cassida* L. (Coleoptera: Chrysomelidae). *Wiadomości entomologiczne*, 22, 62–63. [New records of *Cassida* species from Poland]
- Borowiec, L. & Sassi D. (2003 f) *Discomorpha carlobrivioi*, a new species from Bolivia (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 14, 231–234.
- Borowiec, L. (2003 g) A new species of *Lorentzocassis* Spaeth from New Guinea (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 14, 235–239.
- Borowiec, L. (2003 h) *Omaspides boliviana* n. sp., with a key to *Omaspides* s. str. (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 14, 397–402.
- Borowiec, L. (2003 i) Two new species of the genus *Paraselenis* Spaeth, 1913 (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 14, 403–411.
- Borowiec, L. (2003 j) Two new species of *Spilophora* Boheman from Ecuador and Peru (Coleoptera: Chrysomelidae: Cassidinae). *Annales Zoologici*, Warszawa, 53, 701–704.
- Ghate, H.V., Borowiec, L., Rane, N.S., Ranade, S.P. & Pandit, S. (2003 k) Tortoise beetles and their host plants from Pune (Maharashtra State, India) and nearby places (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 14, 519–539.
- Borowiec, L. (2004 a) *Trilaccodea meridionalis* n. sp., with a key to the genus *Trilaccodea* (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 15, 109–112.
- Borowiec, L. & Skuza, M. (2004 b) The structure of spermatheca in the genus *Chelymorpha* Chevrolat, 1837 (Coleoptera: Chrysomelidae: Cassidinae) and its taxonomic significance. *Annales Zoologici*, Warszawa, 54, 439–451.
- Borowiec, L. & Ghate H.V. (2004 c) Two new species of *Cassida* Linnaeus from southern India (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 15, 247–252.
- Borowiec, L. (2004 d) *Charidotella (Philaspis) marginepunctata*, a new species from Brazil (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 15, 253–256.
- Borowiec, L. (2004 e) *Cteisella flava*, a new species from Mato Grosso, Brazil (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 15, 387–390.
- Borowiec, L. (2005 a) Six new species of the genus *Agroiconota* Spaeth, 1913 (Coleoptera: Chrysomelidae: Cassidinae), with a key to the genus. *Annales Zoologici*, Warszawa, 55, 61–74.
- Borowiec, L. (2005 b) Three new species of the genus *Dorynota* sgen. *Akantaka* (Coleoptera: Chrysomelidae: Cassidinae: Dorynotini). *Genus*, 16, 29–41.
- Borowiec, L. (2005 c) A new species of the genus *Bradycassis* Spaeth, 1952 (Coleoptera: Chrysomelidae: Cassidinae).

Genus, 16, 43–47.

- Borowiec, L. & Moragues, G. (2005 d) Tortoise beetles of the French Guyana - a faunistic review (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 16, 247–278.
- Borowiec, L. (2005 e) *Stolas sanramonensis*, a new species from Bolivia (Coleoptera: Chrysomelidae: Cassidinae: Stolaini). *Genus*, 17, 279–283.
- Borowiec, L. (2005 f) New species and new records of Cassidinae from Southern Africa (Coleoptera: Chrysomelidae). *Mitteilungen aus dem Museum für Naturkunde in Berlin. Zoologische Reihe*, 81, 115–130.
<https://doi.org/10.1002/mmz.200410007>
- Borowiec, L. (2005 g) Revision of Madagascan species of the genus *Chiridopsis* Spaeth, 1922 (Coleoptera: Chrysomelidae: Cassidinae). *Annales Zoologici*, Warszawa, 55, 383–393.
- Borowiec, L. (2006 a) A new species of *Botanochara* Dejean, 1837 from Brazil (Coleoptera: Chrysomelidae: Cassidinae: Stolaini). *Genus*, 17, 127–131.
- Borowiec, L. (2006 b) A new species of *Basiprionota* Chevrolat from Sumba, Indonesia (Coleoptera: Chrysomelidae: Cassidinae: Basiprionotini). *Genus*, 17, 245–248.
- Borowiec, L. (2006 c) A new species of *Aspidimorpha* sgen. *Spaethia* from Zambia (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 17, 249–252.
- Borowiec, L. (2006 d) *Dorynota* (*Akantaka*) *storki*, a new species from Peru, and notes on *Dorynota* (*Akantaka*) *boliviana* Borowiec, 2005 (Coleoptera: Chrysomelidae: Cassidinae: Dorynotini). *Genus*, 17, 369–372.
- Borowiec, L. (2006 e) A new species of *Cassida* Linnaeus from Australia (Coleoptera: Chrysomelidae: Cassidinae: Cassidini). *Genus*, 17, 373–376.
- Borowiec, L. (2006 f) *Syngambria panamensis*, a new species from Panama (Coleoptera: Chrysomelidae: Cassidinae: Cassidini). *Genus*, 17, 377–379.
- Świętojańska, J. & Borowiec, L. (2006 g) *Aspidimorpha* (s. str.) *tibetana*, a new species from China (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 17, 541–544.
- Sassi, D. & Borowiec, L. (2006 h) *Cassida inopinata*, a new species from Italy and Balkan Region (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 17, 545–560.
- Borowiec, L. (2006 i) *Metrionella tucumana*, a new species from Argentina (Coleoptera: Chrysomelidae: Cassidinae: Cassidini). *Genus*, 17, 567–570.
- Borowiec, L. (2006 j) Four new species of *Discomorpha* Chevrolat, 1837 (Coleoptera: Chrysomelidae: Cassidinae: Omocerini), with key to *Discomorpha* s. str. *Zootaxa*, 1357 (1), 45–60.
<https://doi.org/10.11646/zootaxa.1357.1.4>
- Borowiec, L. (2007 a) *Trilaccodea ecuadorica*, a new species from Ecuador (Coleoptera: Chrysomelidae: Cassidinae: Stolaini). *Genus*, 18, 103–106.
- Borowiec, L. & Świętojańska, J. (2007 b) Cassidinae collected during Czech-Polish Expedition to Maharashtra, India (Coleoptera: Chrysomelidae). *Genus*, 18, 279–295.
- Świętojańska, J. & Borowiec, L. (2007 c) Comparative description of last instar larva of *Cassida informis* Boheman, 1862 and *Cassida varians* Herbst, 1799 (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 18, 297–314.
- Borowiec, L. (2007 d) Two new species of *Cassida* Linnaeus, 1758 (Coleoptera: Chrysomelidae: Cassidinae) from Madagascar and notes on subgenera of the genus *Cassida*. *Zootaxa*, 1586 (1), 47–58.
<https://doi.org/10.11646/zootaxa.1586.1.5>
- Borowiec, L. (2007 e) Two new species of *Charidotella* Weise (Coleoptera: Chrysomelidae: Cassidinae: Cassidini), with a key to *Charidotella sexpunctata* group. *Zootaxa*, 1586 (1), 59–66.
<https://doi.org/10.11646/zootaxa.1586.1.6>
- Borowiec, L. (2007 f) A new species of *Cyrtanota* Chevrolat from Peru and note on *Cyrtanota balyi* (Kirsch, 1883) (Coleoptera: Chrysomelidae: Cassidinae: Stolaini). *Genus*, 18, 477–485.
- Borowiec, L. (2007 g) A new species of *Microctenochira* Spaeth from Belize (Coleoptera: Chrysomelidae: Cassidinae: Cassidini). *Genus*, 18, 487–492.
- Borowiec, L. & Opalińska S. (2007 h) The structure of spermathecae of selected genera of Stolaini and Eugenysini (Coleoptera: Chrysomelidae: Cassidinae) and its taxonomic significance. *Annales Zoologici*, Warszawa, 57, 463–479.
- Świętojańska, J. & Borowiec, L. (2007 i) Redescriptions of last instar larvae of *Ischyronota conicicollis* (Weise, 1890) and *Ischyronota desertorum* (Gebler, 1833) (Coleoptera: Chrysomelidae: Cassidinae). *Zootaxa*, 1651 (1), 43–56.
<https://doi.org/10.11646/zootaxa.1651.1.2>
- Borowiec, L. (2007 j) Three new species of the genus *Stolas* Billberg, 1820 (Coleoptera: Chrysomelidae: Cassidinae: Cassidinae). *Genus*, 18, 661–670.
- Borowiec, L. (2007 k) Cassidinae (Coleoptera: Chrysomelidae) collected in Turkey during expeditions of the Upper Silesian Museum, Bytom, Poland. *Annals of the Upper Silesian Museum (Entomology)*, 14–15, 5–10.
- Borowiec, L. (2008 a) *Pseudostilpnaspis belizensis*, a new species of the tribe Cephaloleinini (Coleoptera: Chrysomelidae: Cassidinae: Cassidinae). *Genus*, 19, 89–92.
- Sultan, A., Borowiec, L., Rafi, M.A., Ilyas, M., Naz, F. & Shehzad, A. (2008 b) Tortoise beetles of Rawalpindi-Islamabad area, Pakistan and their host plants (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 19, 93–102.
- Borowiec, L., Sultan, A., Rafi, M.A. & Naz, F. (2008 c) *Notosacantha pakistanica*, a new species From Pakistan (Coleoptera:

- Chrysomelidae: Cassidinae: Notosacanthini). *Genus*, 19, 285–290.**
- Sekerka, L. & Borowiec, L. (2008 d) Three new species of *Cassida* Linne, 1758 from India and note on *Thlaspidia obenbergeri* Spaeth, 1928 (Coleoptera: Chrysomelidae: Cassidinae: Cassidini). *Annales Zoologici*, Warszawa, 58, 611–620.
<https://doi.org/10.3161/000345408X364454>
- Borowiec, L. & Lee, C-F. (2008 e) Redescription of *Cassida insulana* Gressitt, 1952 and notes on some other *Cassida* species from Taiwan (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 19, 699–707.
- Borowiec, L. (2009 a) A new species of *Cyrtanota* Chevrolat from Colombia (Coleoptera: Chrysomelidae: Cassidinae: Mesomphaliini). *Genus*, 20, 111–116.
- Borowiec, L. (2009 b) Revision of the genus *Crambelea* Spaeth, 1913 (Coleoptera: Chrysomelidae: Cassidinae: Cassidini). *Genus*, 20, 117–123.**
- Borowiec, L. & Pomorska, J. (2009 c) The structure of the spermathecae of the genus *Stolas* (Coleoptera: Chrysomelidae: Cassidinae: Mesomphaliini) and its taxonomic significance. *Annales Zoologici*, Warszawa, 59, 201–221.
<https://doi.org/10.3161/000345409X464038>
- Borowiec, L. (2009 d) Two new species of *Charidotella* Weise with black dorsal pattern (Coleoptera: Chrysomelidae: Cassidinae: Cassidini). *Genus*, 20, 341–347.**
- Borowiec, L. & Lee C-F. (2009 e) A new species of *Thlaspidia* Weise from Taiwan, and notes on distribution and host plant of *Cassida insulana* Gressitt (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 20, 349–353.**
- Borowiec, L. (2009 f) New records of Asian and Australopapuan tortoise beetles (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 20, 435–484.**
- Borowiec, L. (2009 g) *Paratrikona albomaculata*, a new species from the Dominican Republic (Coleoptera: Chrysomelidae: Cassidinae: Dorynotini). *Genus*, 20, 567–570.**
- Flinte, V., Borowiec, L., de Freitas, S., Viana, J.H., Fernandes, F.R., Nogueira de Sa, F., de Macedo, M.V. & Monteiro, R.F. (2009 h) Tortoise beetles of the State of Rio de Janeiro, Brazil (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 20, 571–614.
- Borowiec, L. (2009 i) New records of Neotropical tortoise beetles (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 20, 615–722.**
- Borowiec, L. & Sekerka, L. (2010 a) *Aspidimorpha* (s. str.) *karamojana*, a new species from Central Africa (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 21, 271–277.**
- Borowiec, L. (2010 b) A new species of *Cassida* L. from Malaysia (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 21, 279–286.**
- Borowiec, L. (2010 c) *Laccoptera* (*Orphnodella*) *obscurumaculata*, a new species from Central Africa (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 21, 287–292.**
- Borowiec, L. (2010 d) A monograph of the Afrotropical Cassidinae. Supplement 1 (Coleoptera: Chrysomelidae). *Genus*, 21, 535–578.
- Borowiec, L. & Świętojańska, J. (2010 e) A new species of *Stolas* Billberg from Peru (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 21, 579–582.**
- Borowiec, L. (2010 f) Revision of the *Omaspides augusta* group (Coleoptera: Chrysomelidae: Cassidinae: Mesomphaliini). *Annales Zoologici*, Warszawa, 60, 481–492.
<https://doi.org/10.3161/000345410X550373>**
- Borowiec, L. & Sekerka, L. (2010 g) Subfamily Cassidinae Gyllenhal, 1813. In: Löbl, I. & Smetana, A. (Eds.). *Catalogue of Palaearctic Coleoptera. Volume 6. Chrysomeloidea*. Apollo Books, pp.64–65, 368–390.
- Borowiec, L. & Burwell, C.J. (2011 a) A new species of *Cassida* L. from Australia (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 22, 285–290.**
- Borowiec, L. & Cho, H-W. (2011 b) On the subgenus *Lasiocassis* Gressitt (Coleoptera: Chrysomelidae: Cassidinae), with description of a new species from South Korea. *Annales Zoologici*, Warszawa, 61, 445–451.
<https://doi.org/10.3161/000345411X603319>**
- Borowiec, L. (2011 c) Two new species of Cassidinae from Rondonia, Brazil (Coleoptera: Chrysomelidae). *Genus*, 22, 485–491.**
- Borowiec, L. (2011 d) A new species of *Charidotella* Weise from Hispaniola (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 22, 493–497.**
- Borowiec, L. & Takizawa, H. (2011 e) Neotropical tortoise beetles in the Amazon Insectarium, Tokyo, Japan with description of nine new species (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 22, 427–484.**
- Sekerka, L. & Borowiec, L. (2011 f) New synonymy in Palearctic Cassidini (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 22, 505–509.
- Borowiec, L., Ścibior, R. & Kubisz, D. (2011 g) Critical check-list of the Polish Chrysomeloidea, except Cerambycidae (Coleoptera: Phytophaga). *Genus*, 22, 579–608.
- Borowiec, L. & Świętojańska, J. (2011 h) *The Tortoise beetles of Madagascar (Coleoptera: Chrysomelidae: Cassidinae). Part 1: Basiprionotini, Aspidimorphini and Cassidini (except the genus Cassida)*. Polish Taxonomical Monographs vol. 18, Biologica Silesiae, Wrocław, 246 pp.
- Borowiec, L. (2012 a) On the synonymy of two neglected species of *Cassida* L. described by Ritter Böber, 1793 (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 23, 115–117.**

- Borowiec, L. & Stach, M. (2012 b) A new species of *Stolas* Billberg with a radial pattern on the explanate margin of elytra (Coleoptera: Chrysomelidae: Cassidinae: Mesomphaliini). *Genus*, 23, 577–584.
- Świętojańska, J. & Borowiec, L. (2012 c) Cassidinae (Coleoptera: Chrysomelidae) from Socotra Island. *Acta Entomologica Musei Nationalis Pragae*, 52 (Suppl. 2), 381–394.
- Borowiec, L., Takizawa, H. & Świętojańska, J. (2013 a) Five new species of *Notosacantha* Chevrolat (Coleoptera: Chrysomelidae: Cassidinae) from Borneo, with a key to the Bornean species and new faunistic data. *Zootaxa*, 3608 (3), 161–177.
<https://doi.org/10.11646/zootaxa.3608.3.1>**
- Fernandes, F.R. & Borowiec, L. (2013 b) Review of the genus *Cassidinoma* Hincks (Coleoptera: Chrysomelidae: Cassidinae: Omocerini). *Zootaxa*, 3620 (4), 544–552.
<https://doi.org/10.11646/zootaxa.3620.4.4>
- Borowiec, L. & Świętojańska, J. (2013 c) *Plagiometriona obrienorum*, a new species from Rondonia, Brazil (Coleoptera: Chrysomelidae: Cassidinae: Cassidini). *Genus*, 24, 109–114.
- Borowiec, L. & Stach, M. (2013 d) *Stolas radiata*, a new name for *Stolas flavoradiata* Borowiec & Stach, 2012 not *Stolas placida* var. *flavoradiata* Hincks, 1956 (Coleoptera: Chrysomelidae: Cassidinae: Mesomphaliini). *Genus*, 24, 115.
- Świętojańska, J. & Borowiec, L. (2013 e) A new species of *Aspidimorpha* Hope from Sumbawa, Indonesia (Coleoptera: Chrysomelidae: Cassidinae: Aspidimorphini). *Genus*, 24, 217–221.**
- Borowiec, L. & Le Tirant, S. (2013 f) A new species of *Notosacantha* Chevrolat (Coleoptera: Chrysomelidae: Cassidinae: Notosacanthini) from southern India. *Zootaxa*, 3694 (3), 289–395.
<https://doi.org/10.11646/zootaxa.3694.3.9>**
- Świętojańska, J., Moradian, H., Borowiec, L. & Ostovan, H. (2013 g) Description of larvae of two closely related species *Cassida palaestina* Reiche, 1858 and *Cassida rubiginosa* Müller, 1776 (Coleoptera: Chrysomelidae: Cassidinae). *Zootaxa*, 3741 (4), 511–537.
<https://doi.org/10.11646/3894>
- Borowiec, L. & Świętojańska, J. (2013 h) *The Tortoise beetles of Madagascar (Coleoptera: Chrysomelidae: Cassidinae). Part 2: Cassidini, the genus Cassida*. Polish Taxonomical Monographs vol. 20, Biologica Silesiae, Wrocław, 294 pp.**
- Borowiec, L. & Sekerka, L. (2014 a) A new species of *Aspidimorpha* (Coleoptera: Chrysomelidae: Cassidinae: Aspidimorphini) from Uganda. *Zootaxa*, 3796 (1), 197–200.
<https://doi.org/10.11646/zootaxa.3796.1.11>**
- Świętojańska, J., Borowiec, L. & Stach, M. (2014 b) Redescription of immatures and bionomy of the Palaearctic species *Dicladispa testacea* (Linnaeus, 1767) (Coleoptera: Chrysomelidae: Cassidinae: Hispini), a leaf-mining hispine beetle. *Zootaxa*, 3811 (1), 1–33.
<https://doi.org/10.11646/zootaxa.3811.1.1>
- Cho, H-W. & Borowiec, L. (2014 c) Three *Cassida* species new to Korea, with additional faunistic data and key to all Korean species (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 25, 481–492.
- Borowiec, L. & Świętojańska, J. (2014 d) A revision of the genus *Herminella* Spaeth (Chrysomelidae: Cassidinae: Notosacanthini), with a description of a new related genus and species from Madagascar. *Zootaxa*, 3895 (2), 257–272.
<https://doi.org/10.11646/zootaxa.3895.2.7>**
- Borowiec, L. & Świętojańska, J. (2014) Cassidinae Gyllenhal, 1813. In: Leschen, R.B., Beutel, R.G. & Lawrence, J.F. (Eds.), *Handbook of Zoology. Vol. 3. Arthropoda: Insecta, Coleoptera, Beetles, Morphology and Systematics*. De Gruyter, Berlin/Boston, pp. 198–217.
- Świętojańska, J., Massuda, K., Stach, M. & Borowiec, L. (2015 a) Description of immatures of *Chelymorpha reimoseri* Spaeth, 1928 (Coleoptera: Chrysomelidae: Cassidinae: Mesomphaliini). *Zootaxa*, 3949 (4), 515–539.
<https://doi.org/10.11646/zootaxa.3949.4.3>
- Sekerka, L. & Borowiec, L. (2015 b) Subgenera of *Charidotella* Weise with description of a new subgenus and species from Brazil (Coleoptera, Chrysomelidae, Cassidinae, Cassidini). *ZooKeys* 506, 61–74.
<https://doi.org/10.3897/zookeys.506.8770>**
- Borowiec, L. & Sekerka, L. (2015 c) On the genus *Limnocassis* Spaeth, with description of two new species from South Africa (Coleoptera: Chrysomelidae: Cassidinae). *Zootaxa*, 3999 (2), 272–278.
<https://doi.org/10.11646/zootaxa.3999.2.6>**
- Borowiec, L. & Świętojańska, J. (2015 d) Checklist of tortoise beetles (Coleoptera, Chrysomelidae, Cassidinae) from Colombia with new data and description of a new species. *ZooKeys*, 518, 87–127.
<https://doi.org/10.3897/zookeys.518.9350>**
- Sekerka, L., Jia, F., Pang, H. & Borowiec, L. (2016 a) Cassidinae (Coleoptera: Chrysomelidae) types deposited at Sun Yat-sen University, Guangzhou, China. *Zootaxa*, 4084 (1), 50–78.
<https://doi.org/10.11646/zootaxa.4084.1.2>
- Maes, J-M., Gómez-Zurita, J., Riley, E.G., Windsor, D., Borowiec, L. & Chaboo, C. (2016 b) Chrysomelidae (Coleoptera) de Nicaragua, Parte VIII, Cassidinae sensu stricto (tortoise beetles). *Revista Nicaraguense de Entomología*, 76 (VIII), 1–193.
- Świętojańska, J. & Borowiec, L. (2016 c) A new species of *Notosacantha* Chevrolat (Coleoptera: Chrysomelidae: Cassidinae) from Timor-Leste with a key to the *Notosacantha vicaria*-group. *Zootaxa*, 4175 (3), 274–280.

<https://doi.org/10.11646/zootaxa.4175.3.6>

- Borowiec, L. & Świętojańska, J. (2017 a) *Cassida sekerkai* sp. nov. (Coleoptera: Chrysomelidae: Cassidinae), a new species from Madagascar. *Zootaxa*, 4268 (3), 448–450.
<https://doi.org/10.11646/zootaxa.4268.3.10>
- Borowiec, L. & Świętojańska, J. (2017 b) A monograph of the Afrotropical Cassidinae (Coleoptera: Chrysomelidae). Part 4. Revision of the genus *Chiridopsis* Spaeth. *Zootaxa*, 4316 (1), 1–85.
<https://doi.org/10.11646/zootaxa.4316.1.1>**
- Świętojańska, J., Sekerka, L., Borowiec, L., Kwiatek, A. & Windsor, D. (2018 a) Description and comparison of immature stages of four Ischyrosynychini Chapuis, 1875 (Coleoptera: Chrysomelidae: Cassidinae) Species. *Annales Zoologici*, Warszawa, 68 (2), 317–343.
<https://doi.org/10.3161/00034541ANZ2018.68.2.007>
- Borowiec, L. & Coache, A. (2018 b) *Aspidimorpha (Afroaspidimorpha) rainoni*, a new species from Benin (Coleoptera, Chrysomelidae, Cassidinae), Greece. *Faunitaxys*, 6 (11), 1–5.**
- Borowiec, L. & Świętojańska, J. (2018 c) A monograph of the Afrotropical Cassidinae (Coleoptera: Chrysomelidae). Part 5. Revision of the genus *Aethiopocassis* Spaeth. *Zootaxa*, 4488 (1), 1–99.
<https://doi.org/10.11646/zootaxa.4488.1.1>**
- Borowiec, L., Świętojańska, J. & Sekerka L. (2019) Revision of the tribe Cryptonychini (Coleoptera: Chrysomelidae: Cassidinae) of New Caledonia. *Zootaxa*, 4690 (1), 1–71.
<https://doi.org/10.11646/zootaxa.4690.1.1>**
- Borowiec, L. & Świętojańska, J. (2022) A monograph of the Afrotropical Cassidinae (Coleoptera: Chrysomelidae). Part 6. Revision of the tribe Cassidini 3, the genus *Cassida* Spaeth. *Zootaxa*, 5171 (1), 1–250.
<https://doi.org/10.11646/zootaxa.5171.1.1>**
- Villamil, S.C., Borowiec, L. & Świętojańska, J. (2023) *Paraselenis (Spaethiechoma) convolvuli* (Coleoptera: Chrysomelidae: Cassidinae: Mesomphaliini), a new species from Argentina, with a description of its immature stages and notes on its bionomy. *Annales Zoologici*, 73, 701–725.
<https://doi.org/10.3161/00034541ANZ2023.73.4.011>**

A list of papers with descriptions of new taxa whose types are preserved in the Lech Borowiec collection

- Boheman, C.H. (1850) *Monographia Cassididarum. Tomus primus*. Holmiae, 452 pp. + 4 tab.
<https://doi.org/10.5962/bhl.title.8973>
- Boheman, C.H. (1855) *Monographia Cassididarum. Tomus tertius*. Holmiae, 543 pp. + 1 tab.
- Boheman, C.H. (1862) *Monographia Cassididarum. Tomus quartus*. Holmiae, 504 pp.
- Bordy, B. (1995 a) *Cassida leucanthemi* n. sp., espece jumelle de *C. sanguinosa* Suffrian, 1844, (Coleoptera, Chrysomelidae). *Bulletin de la Société entomologique de France*, 100, 377–383.
<https://doi.org/10.3406/bsef.1995.17173>
- Bordy, B. (1995 b) *Cassida bergeali*, espece nouvelle du groupe de *Cassida vibex* L. 1767 (Col. Chrysomelidae). *Nouvelle Revue d'Entomologie* (N.S.), 12, 243–248.
- Bordy, B. (2009) Identification des especes du complexe *Hypocassida subferruginea* (Schränk, 1776) (Coleoptera, Chrysomelidae). *Nouvelle Revue d'Entomologie* (N.S.) [2008], 25, 45–59.
- Dohrn, C. (1880) Exotisches. *Stettiner Entomologische Zeitung*, 41, 367–382.
- Fairmaire, M.L. (1898) Matériaux pour la Faune coléoptérique de la région Malgache, 5 note. *Annales de la Société royale belge d'entomologie*, 42, 222–260.
- Fairmaire, M.L. (1899 a) Matériaux pour la faune coléoptérique de la région Malgache, 8 note. *Annales de la Société royale belge d'entomologie*, 43, 511–558.
- Fairmaire, M.L. (1899 b) Matériaux pour la faune coléoptérique de la région Malgache, 9 note. *Annales de la Société entomologique de France*, 1899, 466–507.
<https://doi.org/10.1080/21686351.1899.12279744>
- Fairmaire, L. (1900) Descriptions de Coléopteres recueillis par M. H. Perrier de la Bâthie a Madagascar. *Bulletin de la Société entomologique de France*, 1900, 85–90.
<https://doi.org/10.3406/bsef.1900.22548>
- Fairmaire, L. (1901) Descriptions de quelques Coléopteres recueillis par M. le D. Decorse dans le sud de Madagascar, Plateau de l'Androy. *Notes Leyden Museum*, 23, 65–84.
- Fairmaire, L. (1903) Description de Coléopteres nouveaux de Madagascar. *Naturaliste*, 25, No 380, 9–11.
- Hincks, W.D. (1962) Madagascar Cassidinae (Col., Chrysomelidae) Part I. *Naturalista Malgache*, 13, 225–250.
- Matis, E.G. (1974) Novii vid roda *Cassida* (Coleoptera, Chrysomelidae) iz Primoria. *Zoologicheskij Zhurnal*, 53, 1100–1102.
- Maulik, S. (1918) Two new Indian Cassidinae beetles. *The Annals and Magazine of Natural History*, ser. 9, 1, 322–325.
<https://doi.org/10.1080/00222931808562321>
- Medvedev, L.N. & Eroshkina, G.A. (1982) Fauna i ekologija Cassidinae (Coleoptera, Chrysomelidae) plato Taj-Nguien. *In:*

- Zhivotnii mir Vietnama*. Nauka, Moskva, pp. 103–115.
- Medvedev, L.N. & Eroshkina, G.A. (1988) Revizya listoyedov-shchitonosok (Chrysomelidae, Cassidinae) fauni Vietnama. In: *Fauna i ekologiya nasekomi Vietnama*. Nauka, Moskva, pp. 105–142.
- Sekerka, L. (2005) *Cassida olympica*, a new species from Greece (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 16, 285–289.
- Sekerka, L. (2006) A new species of *Cassida* Linné, 1758 from Cyprus (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 17, 253–262.
- Sekerka, L. (2011) *Cassida stevensi*, a new species from India (Coleoptera: Chrysomelidae: Cassidinae: Cassidini). *Genus*, 22, 499–504.
- Sekerka, L. (2014) Review of Imatidiini genera (Coleoptera: Chrysomelidae: Cassidinae). *Acta Entomologica Musei Nationalis Pragae*, 54, 257–314.
- Sekerka, L. & Windsor, D.M., in litt. Unpublished monograph of Panamanian Cassidinae.
- Spaeth, F. (1901 a) Beschreibung neuer Cassididen nebst synonymischen Bemerkungen. IV. *Verhandlungen der Zoologisch-Botanischen Gesellschaft in Wien*, 51, 333–350.
- Spaeth, F. (1901 b) Ueber *Chelysida* und *Oxylepus* als zwei verschiedene Cassiden-Gattungen. *Verhandlungen der Zoologisch-Botanischen Gesellschaft in Wien*, 51, 750–756.
- Spaeth, F. (1901 c) Neue Cassiden aus Sumatra gesammelt von Dr. H. Dohrn. *Stettiner Entomologische Zeitung*, 62, 3–13.
- Spaeth, F. (1905) Beschreibung neuer Cassididen nebst synonymischen Bemerkungen. V. *Verhandlungen der Zoologisch-Botanischen Gesellschaft in Wien*, 55, 79–118.
- Spaeth, F. (1909) Beschreibung neuer Cassididen nebst synonymischen Bemerkungen. VII. *Verhandlungen der Zoologisch-Botanischen Gesellschaft in Wien*, 59, 364–397.
- Spaeth, F. (1911) Beschreibung neuer Cassididen nebst synonymischen Bemerkungen. VIII. *Verhandlungen der Zoologisch-Botanischen Gesellschaft in Wien*, 61, 239–277.
- Spaeth, F. (1912) Neue Cassiden aus Columbien, Peru, Bolivien und Ecuador. *Stettiner Entomologische Zeitung*, 73, 3–16.
- Spaeth, F. (1913 a) Kritische Studien über den Umfang und die Begrenzung mehrerer Cassiden-Gattungen nebst Beschreibung neuer amerikan. Arten. *Archiv für Naturgeschichte*, 79, 126–164.
- Spaeth, F. (1913 b) Studien über die Gattung *Hoplionota* Hope und Beschreibung einer verwandten neuen Gattung. *Verhandlungen der Zoologisch-Botanischen Gesellschaft in Wien*, 63, 381–534.
- Spaeth, F. (1914 a) Neue Cassididen aus Paraguay und Goyaz (Col.). *Entomologische Mitteilungen*, 3, 166–168.
<https://doi.org/10.5962/bhl.part.5083>
- Spaeth, F. (1914 b) Zur Kenntnis der indischen Cassidinen. *Deutsche Entomologische Zeitschrift*, 1914, 542–568.
<https://doi.org/10.1002/mmnd.48019140506>
- Spaeth, F. (1914 c) Neue Cassidinen aus Yünnan (Col.). *Entomologische Mitteilungen*, 3, 226–230.
<https://doi.org/10.5962/bhl.part.5085>
- Spaeth, F. (1915) Neue Cassidinen (Coleoptera). *Stettiner Entomologische Zeitung*, 76, 265–290.
- Spaeth, F. (1917) Neuer Beitrag zur Kenntnis der Ost- und Zentralafrikanischen Cassidinen. *Annales historico-naturales Musei Nationalis Hungarici*, 15, 422–444.
- Spaeth, F. (1919) Neue Cassidinae aus der Sammlung von Dr. K. Brancsik, dem Ungarischen National-Museum und meiner Sammlung. *Annales historico-naturales Musei Nationalis Hungarici*, 17, 184–204.
- Spaeth, F. (1922) Chrysomérides, Cassidines. In: *Voyage de M. Le Baron Maurice de Rothschild en Ethiopie & en Afrique Orientale Anglaise (1904-1905)*. Imprimerie Nationale, Paris, pp. 997–1004.
- Spaeth, F. (1924) Cassidinae. In: *Voyage de CH. Alluaud & R. Jeannel en Afrique orientale (1911-1912). Résultats scientifiques. Coleoptera XVIII*. A. Schultz, Paris, pp. 275–363.
- Spaeth, F. (1926 a) Monographie der zur Gruppe der Coptocyclitae gehörigen amerikanischen Cassidinen (Col.): I. Die Gattungen mit gekammten Klauen. *Supplementa Entomologica*, 13, 1–108.
- Spaeth, F. (1926 b) Beschreibung neuer Cassidinen. *Bulletins mensuels (Société des naturalistes luxembourgeois) N.S.*, 20, 11–24, 47–60.
- Spaeth, F. (1926 c) Mitteilungen ueber die Cassidinen des Prager Nationalmuseums (Col. Chrysomelidae). *Sborník Entomologického Oddělení Národního Muzea v Praze*, 4, 81–96.
- Spaeth, F. (1926 d) Neue Cassidinen des Rijksmuseums in Leiden, des British Museums und meiner Sammlung. *Zoologische Medelingen*, 9, 1–15.
- Spaeth, F. (1928) Mitteilungen ueber die Cassidinen des Nationalmuseum in Prag. II. (Col., Chrysomelidae). *Sborník Entomologického Oddělení Národního Muzea v Praze*, 6, 50, pp. 29–47.
- Spaeth, F. (1931) Neue Beiträge zur Kenntnis der afrikanischen Cassidinen. (Col. Chrys.). *Revue de Zoologie & de Botanique Africaines*, 21, 100–112.
- Spaeth, F. (1932) Neue Beiträge zur Kenntnis der Afrikanischen Cassidinen (Col. Chrys.). *Revue de Zoologie & de Botanique Africaine*, 22, pp. 2–22, 227–241.
- Spaeth, F. (1933) Neue Beiträge zur Kenntnis der Afrikanischen Cassidinen (Col. Chrys.). *Revue de Zoologie & de Botanique Africaine*, 22, 345–359.
- Spaeth, F. (1935) V. Cassidinen. In: *Über die Coleopteren-Ausbeute der II. Schweizer wissenschaftlichen Expedition nach Angola 1932/33. Stettiner Entomologische Zeitung*, 96, 172–176.

- Spaeth, F. (1936 a) Die Untergattungen von *Coptocycla* mit gerieften Klauen. Festschrift zum 60. Geburtstage von Professor Dr. Embrik Strand, Vol. I, Riga, 17 X 1936: 251–261.
- Spaeth, F. (1936 b) Neue Cassidinen und Hispinen (Col.) aus dem British Museum. *Proceedings of the Royal Entomological Society of London*, ser. B 5, 8–11.
<https://doi.org/10.1111/j.1365-3113.1936.tb01303.x>
- Spaeth, F. (1937) Neue Cassidinen des Rijksmuseums in Leiden und meiner Sammlung. *Temminckia*, 2, 135–158.
- Spaeth, F. (1938) Über die von Herrn R. Malaise 1934 in Burma gesammelten Cassidinen. *Entomologisk Tidskrift*, 59, 228–236.
- Spaeth, F. (1940) Ueber die Gattung *Poecilaspis* Hope (Coleoptera, Cassidinae). *Temminckia*, 5, 143–162.
- Świętojańska, J. (2000 a) Five new species of *Notosacantha* Chevrolat from the Philippines (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 11, 205–217.
- Świętojańska, J. (2000 b) *Notosacantha multicostata*, a new species from Madagascar (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 11, 593–596.
- Świętojańska, J. (2001 a) Two new species of *Notosacantha* Chevrolat from Sumatra (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 12, 51–57.
- Świętojańska, J. (2001 b) Four new species of *Notosacantha* Chevrolat from Borneo (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 12, 335–347.
- Świętojańska, J. (2001 c) A revision of the tribe *Aspidimorphini* of the Oriental Region (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, suppl., 318 pp. + 18 pl.
- Świętojańska, J. (2002 a) *Notosacantha bezdeki*, a new species from Laos (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 13, 39–42.
- Świętojańska, J. (2002 b) *Notosacantha riedeli*, a new species from Sulawesi (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 13, 353–356.
- Świętojańska, J. (2002 c) Revision of the genera *Aporocassida* Spaeth, 1952 and *Saulaspis* Spaeth, 1913 (Coleoptera: Chrysomelidae: Cassidinae). *Annales Zoologici*, Warszawa, 52, 573–581.
- Świętojańska, J. (2003 a) *Notosacantha viridipennis*, a new species from Madagascar (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 14, 393–396.
- Świętojańska, J. (2003 b) Two new species of the tribe *Stolaini* (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 14, 511–518.
- Świętojańska, J. (2004) Two new species of *Notosacantha* Chevrolat from Malaysia and Myanmar (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 15, 241–246.
- Świętojańska, J. (2006) A new species of *Notosacantha* Chevrolat from Halmahera, Indonesia (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 17, 363–367.
- Świętojańska, J. & Sekerka, L. (2007) *Notosacantha warchalowskii*, a new species from India (Coleoptera: Chrysomelidae: Cassidinae: Notosacanthini). *Genus*, 18, 681–685.
- Świętojańska, J. & Stach, M. (2011) Two new species of *Notosacantha* Chevrolat (Coleoptera: Chrysomelidae: Cassidinae) from the Oriental region. *Annales Zoologici*, Warszawa, 61, 421–426.
<https://doi.org/10.3161/000345411X584870>
- Świętojańska, J., Ghate, H.V. & Marathe, R. (2001) *Notosacantha chandrapurensis* n. sp. from India (Coleoptera: Chrysomelidae: Cassidinae). *Genus*, 12, 489–492.
- Weise, J. (1897) Kritisches Verzeichniss der von Mr. Andrewes eingesandten Cassidinen und Hispinen aus Indien. *Deutsche Entomologische Zeitschrift*, 1897, 97–150.
<https://doi.org/10.1002/mmnd.48018970120>
- Weise, J. (1900 b) Einige neue Hispinen und Cassidinen von Paul Weise in Usambara gesammelt. *Deutsche Entomologische Zeitschrift*, 1900, 213–217.
<https://doi.org/10.1002/mmnd.48019000128>

Reference cited in Introduction

- Sekerka, L. & Świętojańska, J. (2024) Cassidinae. In: Bezděk, J. & Sekerka L. (Eds.), *Catalogue of Palaearctic Coleoptera, Volume 6/2/1, updated and revised second edition, Chrysomeloidea II (Orsodacnidae, Megalopodidae, Chrysomelidae)*. Brill, Leiden & Boston, pp. 34–59, 233–272.